

Search for the onset of baryon anomaly at RHIC-PHENIX


Tatsuya Chujo
for the PHENIX collaboration



Introduction

~ Baryon Anomaly at RHIC ~

PHENIX: PRL 91, 172301 (2003), PRC 69, 034909 (2004)




- In Au+Au $\sqrt{s_{NN}} = 200$ GeV central collisions:
 - Pions: Strong suppression of yields above $p_T \sim 2$ GeV/c, due to jet quenching effect.
 - Protons: No suppression for at intermediate p_T (2-5 GeV/c).
- **Called “Baryon anomaly at RHIC”.**
- **Quark recombination models** reproduce the data qualitatively.

Nuclear Modification Factor

$$R_{AA}(p_T) = \frac{\text{yield}(AuAu) / N_{\text{coll}}}{\text{yield}(pp)} \sim R_{CP}$$

p/π ratios in Au+Au 200 GeV

More (anti) baryons than pions at moderate p_T (2-5 GeV/c).
Does not look like vacuum jet fragmentation.





Where is the onset of the baryon anomaly at RHIC?

→ New Cu+Cu data at lower $\sqrt{s_{NN}}$ may give an answer...

Data Analysis



- **Data set:**
 - RHIC Run-5 (2005), measured by the PHENIX experiment.
 - **Cu+Cu at 22.5 GeV.**
 - **Cu+Cu at 62.4 GeV.**
- **Detectors:**
 - Drift Chamber, PC1, BBC and TOF for PID charged analysis.
- **Centrality:**
 - Subdivided minimum bias triggered events, based on BBC charge (62 GeV), or the number of PC1 hit (22 GeV).
- **Corrections:**
 - Geometrical acceptance, in flight decay.
 - NOTE: No weak decay feed-down correction applied.



Results



1. p_T spectra in Cu+Cu at $\sqrt{s_{NN}} = 22.5$ and 62.4 GeV.
2. p/π^+ and $\bar{p}/\pi^-$ ratios vs. p_T .
3. Nuclear modification factor: R_{AA} .

p_T spectra for pions





Filled symbols : π^+
Open symbols : π^-

p_T spectra for kaons





Filled symbols : K^+
Open symbols : K^-

p_T spectra for protons





* No weak decay feed-down correction applied

p_T spectra for antiprotons





* No weak decay feed-down correction applied

p/π in Cu+Cu 22.5 GeV





- **Larger p/π^+ ratios** than those in Au+Au 200 GeV (0.5 @ $p_T = 1.5$ GeV).
 - Affected by the participant protons (not produced protons) at lower $\sqrt{s_{NN}}$.
- **$\bar{p}/\pi^-$ ratios are ~ 0.3 - 0.4 at $p_T = 2$ GeV/c**, which is close to the fragmentation expectations in p+p.
 - **Almost no centrality dependence.**

p/π in Cu+Cu 62.4 GeV




- p/π^+ ratios are reduced, compared to those in Cu+Cu 22 GeV.
- $\bar{p}/\pi^-$ ratio in most central collisions reaches $R \sim 0.6$ at $p_T = 2$ GeV/c.
 - Centrality dependence is seen.
 - Peripheral $\bar{p}/\pi^-$: consistent with p+p value.

Beam energy dep. in Cu+Cu



- **p/π^+ ratio** : decreasing as a function of $\sqrt{s}_{NN}$.
- **$\bar{p}/\pi^-$ ratio** : increasing as a function $\sqrt{s}_{NN}$.
 - Cu+Cu 22.5 GeV central data reaches the p+p values.
 - Cu+Cu 62.4 GeV central data is higher than that in 22.5 GeV.


R_{AA} for pions (22 GeV)



• p+p reference: global fit of pion data at $\sqrt{s} \sim 23$ GeV (D. d'Enterria).

- p+p reference: global fit of pion data at $\sqrt{s} \sim 22$ GeV in p+p.
- R_{AA} is ~ 1.5 at $p_T = 2$ GeV/c, **no suppression**.


R_{AA} for pions (62 GeV)



- p+p reference: global fit of pion data at $\sqrt{s} \sim 62$ GeV.
- nucl-ex/ nucl-ex/0411049, D. d'Enterria.

- Similar, might be slightly smaller than that in 22 GeV.


R_{AA} for proton (22 GeV)



• p+p reference: fitted the ISR data in p+p 23 GeV, B. Alper et al, NPB 100, 237 (1975).

- **Larger than pions significantly**, which suggests the large contributions of participant protons.


R_{AA} for proton (62 GeV)



• p+p reference: fitted the ISR data in p+p 63 GeV, B. Alper et al, NPB 100, 237 (1975).

• Similar as in 22.5 GeV, but slightly smaller value.


R_{AA} for antiproton (22 GeV)



• p+p reference: fitted the ISR data in p+p 23 GeV, B. Alper et al, NPB 100, 237 (1975).

- Different p_T dependence from protons.
- $R_{AA} \sim 1$ (similar to pions' R_{AA}), binary collision scaling worked?

R_{AA} for antiproton (62 GeV)



• p+p reference: fitted the ISR data in p+p 63 GeV, B. Alper et al, NPB 100, 237 (1975).

• Similar to that for 22 GeV (and also pion's R_{AA}).

Summary

Measured p_T spectra for $\pi^\pm$, $K^\pm$, p , $\bar{p}$ in Cu+Cu at $\sqrt{s_{NN}} = 22.5$ and 62.4 GeV in PHENIX.

- p/π^+ ratio : **decreasing** as a function of $\sqrt{s_{NN}}$ in Cu+Cu.
 - Suggests a significant contribution of participant protons (not by the produced protons) in lower energies Cu+Cu.
- $\bar{p}/\pi^-$ ratio : **increasing** as a function $\sqrt{s_{NN}}$ in Cu+Cu.
 - Cu+Cu 22.5 GeV shows that there is (almost) **no centrality dependence**, and central Cu+Cu already reaches to the p+p values.
 - Cu+Cu 62.4 GeV central data is higher than that in 22.5 GeV, centrality dependence is seen.
- R_{AA} : depends on particle type.
 - π : no suppression at both 22.5 and 62.4 GeV in Cu+Cu.
 - p : enhanced (mostly due to the participant nucleons).
 - $\bar{p}$: binary scaling, very similar to pion's R_{AA} .
- *New reference p+p data at 62.4 GeV will be taken in June 2006 (2 weeks).*

- University of São Paulo, São Paulo, Brazil
- Academia Sinica, Taipei 11529, China
- China Institute of Atomic Energy (CIAE), Beijing, P. R. China
- Peking University, Beijing, P. R. China
- Charles University, Faculty of Mathematics and Physics, Ke Karlovu 3, 12116 Prague, Czech Republic
- Czech Technical University, Faculty of Nuclear Sciences and Physical Engineering, Brehova 7, 11519 Prague, Czech Republic
- Institute of Physics, Academy of Sciences of the Czech Republic, Na Slovance 2, 182 21 Prague, Czech Republic
- Laboratoire de Physique Corpusculaire (LPC), Université de Clermont-Ferrand, 63 170 Aubière, Clermont-Ferrand, France
- Dapnia, CEA Saclay, Bat. 703, F-91191 Gif-sur-Yvette, France
- IPN-Orsay, Université Paris Sud, CNRS-IN2P3, BP1, F-91406 Orsay, France
- Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS-IN2P3, Route de Saclay, F-91128 Palaiseau, France
- SUBATECH, École des Mines at Nantes, F-44307 Nantes France
- University of Münster, Münster, Germany
- KFKI Research Institute for Particle and Nuclear Physics at the Hungarian Academy of Sciences (MTA KFKI RMKI), Budapest, Hungary
- Debrecen University, Debrecen, Hungary
- Eötvös Loránd University (ELTE), Budapest, Hungary
- Banaras Hindu University, Banaras, India
- Bhabha Atomic Research Centre (BARC), Bombay, India
- Weizmann Institute, Rehovot, 76100, Israel
- Center for Nuclear Study (CNS-Tokyo), University of Tokyo, Tanashi, Tokyo 188, Japan
- Hiroshima University, Higashi-Hiroshima 739, Japan
- KEK - High Energy Accelerator Research Organization, 1-1 Oho, Tsukuba, Ibaraki 305-0801, Japan
- Kyoto University, Kyoto, Japan
- Nagasaki Institute of Applied Science, Nagasaki-shi, Nagasaki, Japan
- RIKEN, The Institute of Physical and Chemical Research, Wako, Saitama 351-0198, Japan
- RIKEN – BNL Research Center, Japan, located at BNL
- Physics Department, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima, Tokyo 171-8501, Japan
- Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan
- University of Tsukuba, 1-1-1 Tennodai, Tsukuba-shi Ibaraki-ken 305-8577, Japan
- Waseda University, Tokyo, Japan
- Cyclotron Application Laboratory, KAERI, Seoul, South Korea
- Kangnung National University, Kangnung 210-702, South Korea
- Korea University, Seoul, 136-701, Korea
- Myong Ji University, Yongin City 449-728, Korea
- System Electronics Laboratory, Seoul National University, Seoul, South Korea
- Yonsei University, Seoul 120-749, Korea
- IHEP (Protvino), State Research Center of Russian Federation "Institute for High Energy Physics", Protvino 142281, Russia
- Joint Institute for Nuclear Research (JINR-Dubna), Dubna, Russia
- Kurchatov Institute, Moscow, Russia
- PNPI, Petersburg Nuclear Physics Institute, Gatchina, Leningrad region, 188300, Russia
- Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Vorob'evy Gory, Moscow 119992, Russia
- Saint-Petersburg State Polytechnical University, Politechnicheskayastr, 29, St. Petersburg, 195251, Russia




13 Countries; 62 Institutions; 550 Participants*

- Lund University, Lund, Sweden
- Abilene Christian University, Abilene, Texas, USA
- Brookhaven National Laboratory (BNL), Upton, NY 11973, USA
- University of California - Riverside (UCR), Riverside, CA 92521, USA
- University of Colorado, Boulder, CO, USA
- Columbia University, Nevis Laboratories, Irvington, NY 10533, USA
- Florida Institute of Technology, Melbourne, FL 32901, USA
- Florida State University (FSU), Tallahassee, FL 32306, USA
- Georgia State University (GSU), Atlanta, GA, 30303, USA
- University of Illinois Urbana-Champaign, Urbana-Champaign, IL, USA
- Iowa State University (ISU) and Ames Laboratory, Ames, IA 50011, USA
- Los Alamos National Laboratory (LANL), Los Alamos, NM 87545, USA
- Lawrence Livermore National Laboratory (LLNL), Livermore, CA 94550, USA
- University of New Mexico, Albuquerque, New Mexico, USA
- New Mexico State University, Las Cruces, New Mexico, USA
- Department of Chemistry, State University of New York at Stony Brook (USB), Stony Brook, NY 11794, USA
- Department of Physics and Astronomy, State University of New York at Stony Brook (USB), Stony Brook, NY 11794, USA
- Oak Ridge National Laboratory (ORNL), Oak Ridge, TN 37831, USA
- University of Tennessee (UT), Knoxville, TN 37996, USA
- Vanderbilt University, Nashville, TN 37235, USA

***as of March 2005**

Backup Slides

p+p fit at $\sqrt{s} = 22.5$ GeV (pions)



p+p $\rightarrow \pi + X$ @ $\sqrt{s} \approx 22.5$ GeV

- p+p $\rightarrow \pi^- + X$ $\sqrt{s} = 21.7$ GeV -- EHS-NA22 [adamus88]
- p+p $\rightarrow \pi^0 + X$ $\sqrt{s} = 21.7$ GeV -- FNAL E-063 [carey76]
- p+p $\rightarrow \pi^0 + X$ $\sqrt{s} = 22.8$ GeV -- FNAL E-063 [carey76]
- p+p $\rightarrow \pi^0 + X$ $\sqrt{s} = 23.0$ GeV -- CERN-WA70 [bonesi89]
- p+p $\rightarrow \pi^0 + X$ $\sqrt{s} = 23.1$ GeV -- FNAL E-063 [carey76]
- p+p $\rightarrow \pi^0 + X$ $\sqrt{s} = 23.8$ GeV -- FNAL E-063 [carey76]
- p+p $\rightarrow \pi^0 + X$ $\sqrt{s} = 23.8$ GeV -- CERN-NA24 [demarzo87]
- p+p $\rightarrow \pi^0 + X$ $\sqrt{s} = 23.8$ GeV -- FNAL-E-268 [donaldson78]
- p+p $\rightarrow \pi^0 + X$ $\sqrt{s} = 23.3$ GeV -- R-107 [lloydowen80]
- p+p $\rightarrow \pi^0 + X$ $\sqrt{s} = 23.5$ GeV -- CCRS [busser76]
- p+p $\rightarrow \pi^0 + X$ $\sqrt{s} = 23$ GeV -- ACHM [eggert75]
- p+p $\rightarrow \pi^+ + X$ $\sqrt{s} = 23$ GeV -- Brit.-Scand.[alper75]
- p+p $\rightarrow \pi^+ + X$ $\sqrt{s} = 23$ GeV -- Brit.-Scand.[alper75]

Compilation and fitting
by D. d'Enterria

R_{AA} for π^0 in Cu+Cu 22.5 GeV PHENIX

PbSc (0-10%)




Figure 10: R_{AA} for the PbSc calorimeter and the Centrality Class 0-10%

PbGl (0-10%)




Figure 14: R_{AA} for the PbGl calorimeter and the Centrality Class 0-10%

SPS and AGS




Fig. 47. Invariant yields of p , $\bar{p}$, and π as function of p_T in central Pb + Pb collisions at the SPS ($\sqrt{s_{NN}} = 17$ GeV) (left panel) and in central Au + Au collisions at the AGS ($\sqrt{s_{NN}} = 5$ GeV) (right panel). The $\bar{p}$ spectrum from the AGS is scaled up by a factor 100. All data are at mid-rapidity ($y - y_{cm} \approx 0$) and are from W98 [163], NA44 [226], NA49 [227], and E866 [228,229].

Nucl. Phys. A 757, 184 (2005),
PHENIX, nucl-ex/0410003

NA49 data






Figure 2. The energy dependence of proton/pion ratios (left panel), and a comparison of the baryon/meson ratios at top SPS energy to a Blast-Wave parametrization (right panel).

$\bar{p}/\pi^-$ (NA49)



p/π^+ (NA49)

NA49 collaboration, nucl-ex/0510054

R_{AA} for pions





22.5 GeV





62.4 GeV

R_{AA} for kaons





22.5 GeV





62.4 GeV


R_{AA} summary (22 GeV)




R_{AA} summary (62 GeV)



$\bar{p}/p$ ratio vs. p_T




22.5 GeV




62.4 GeV

$\pi^-/\pi^+, K^-/K^+$ vs. p_T (22.5 GeV)









π^-/π^+ , K^-/K^+ vs. p_T (62.4 GeV)



p+p reference data (with fit)

23 GeV

63 GeV



Kaons and protons: Fit results of ISR p+p 63 GeV data. B. Alper et al, NPB100(1975)237.
pions: global fits of p+p data, e.g. nucl-ex/0411049, D. d'Enterria.