


# Diffraction measurements at HERA

Henri Kowalski

BNL-Task Force Meeting

29 of April 2010

very similar to the EDS2009 Talk in Geneva



$Q^2$  - virtuality of the incoming photon

$W$  - CMS energy of the incoming photon-proton system

$M_X$  - invariant mass of all particles seen in the central detector


$t$  - momentum transfer to the diffractively scattered proton

$$\beta = Q^2 / (Q^2 + M^2)$$

$$\times_{IP} = (Q^2 + M^2) / (W^2 + M^2)^2$$

In the first month of HERA data taking the analysis trigger killed the diffractive signal with 100% efficiency


In ZEUS the analysis trigger required energy depositions in forward AND backward calorimeters (to measure vertex timing)



Interesting diffraction  
can be seen with  
small lumi,  $\sim 10 \mu\text{b}^{-1}$

→ Watch  
the trigger and  
the cuts

# Rapidity Gap Selection



Select diffractive events by requirement:


No energy deposition in some area of the detector

-  $\eta_{\text{max}}$  cut

no energy means no cluster with  $> 400 \text{ MeV}$

note: noise  $O(100) \text{ MeV}$  per cell

**ZEUS Collaboration; M.Derrick et al.**  
**Observation of Events with a Large Rapidity Gap in Deep Inelastic Scattering at HERA**  
**DESY 93-093 (July 1993)**  
**Physics Letters B 315 (1993) 481-493**




Shape of MC ?  
Shifts of MC ?

**First diffractive signal seen in DIS**

# $D^*$ (2010) Signal Plots

## ZEUS



→ Watch  
the Monte Carlos

ZEUS Collaboration; J.Breitweg et al.

Measurement of the Diffractive Cross Section in Deep Inelastic Scattering using ZEUS 1994 Data  
DESY 98-084 (July 1998)

*The European Physical Journal* **1994** (1999) 43-66

as a function of  $W$  and  $M_X$




Fig. 1. Reaction  $\gamma^* p \rightarrow X + \text{anything}$ , where  $X$  is the system observed in the detector. *Top:* Distributions of  $M_X$ , the corrected mass of the system  $X$ . The distributions are not corrected for acceptance effects. The shaded histograms show the distributions of events with  $\eta_{\text{max}} < 1.5$ . *Bottom:* Same distributions as above presented in terms of  $\ln M_X^2$ . The straight lines give the nondiffractive contributions as obtained from the fits. The upper curves show the fit results for the sum of the diffractive and nondiffractive contributions





Figure 8: Distributions of  $\ln M_X^2 + \ln(s/W^2)$  for the  $W$  intervals 90 - 110 GeV (dotted), 134 - 164 GeV (dashed), 200 - 245 GeV (solid) ( $\ln W^2 = 9.0 - 9.4, 9.8 - 10.2, 10.6 - 11.0$ ) at a)  $Q^2 = 14 \text{ GeV}^2$  and b)  $31 \text{ GeV}^2$ . Here  $M_X$  is the corrected mass; the distributions are the measured ones, not corrected for acceptance effects. For each  $Q^2$  the three distributions were normalized to the same number of events.






- $W$  - the CMS energy
- $\Delta Y = \ln(W/M_x)^2$  length of the rap-gap in the event
- $\Delta Y = \ln(W/M_x)^2 - \eta_0$  length of the rap-gap seen in the detector

### Non-Diffractive Event



### Diffractive Event



non-diff events are characterized by uniform, uncorrelated particle emission along the whole rapidity axis => probability to see a gap  $\Delta Y$  is

$$\sim \exp(-\lambda \Delta Y) \quad \text{--- Poisson } P(0, \Delta Y)$$

$\lambda$  - Gap Suppression Coefficient

(average multiplicity per unit of  $Y$ )


Examples of probabilities to see a gap  $\Delta Y$  in an non-diff event -  $\exp(-1.7 \Delta Y)$

- $P(1) = 18\%$        $P(2) = 3.3\%$
- $P(3) = 0.6\%$

# Non-Diff Color Dipole MC

$$\Delta Y = \ln(W/M_X)^2$$

$$\Delta Y = \ln(1/x_{IP}) \quad ?$$



In MC  $\lambda$  independent of  $Q^2$  and  $W^2$

$\lambda \sim 2$  in MC

$\lambda \sim 1.7$  in data

→ Watch  
the Monte Carlos

Probability to see a gap  $\Delta Y$  in a non-diff event -  $\exp(-\lambda\Delta Y)$

Physical interpretation of the Gap Suppression Coef.  $\lambda \sim 1.7$

Feynman (~1970):  $\lambda$  depends on the quantum numbers carried by the gap

Photon – Hadron  
Interactions,  
lecture 52

$\lambda = 2$  for the exchange of pion q.n.


$= 1$  for the exchange of rho q.n.

$= 0$  for the exchange of pomeron q.n.



In the Longitudinal Phase Space Model

$\lambda$  – particle multiplicity per unit of rapidity  
cluster

# $M_X$ Method




# Rapidity Gap Selection $\longleftrightarrow$ $M_X$ Method



# Effect of FPC on ZEUS Diffractive Measurement

## FPC was added in 1998




# ZEUS

FPC added

$M_X$ -00

Larger  $W$  and  $M_X$  range  
but  
more proton dissociation




$$\ln M_X^2 < \ln W^2 - \eta_0 - 2$$

$$\ln 46^2 = 7.7 \quad M_X \sim 6 \text{ GeV}$$

$$\ln 222^2 = 10.8 \quad M_X \sim 27 \text{ GeV}$$



# Reggeon Contribution



Reggeon MC  
reweighted to  
reproduce LPS data


Figure 6: Distributions of  $\ln M_X^2$  ( $M_X$  in units of GeV) at the detector level for different  $(W, Q^2)$  bins. The points with error bars show the data. The hatched histograms show the contributions predicted by the exchange of the  $\rho$ -Reggeon trajectory. The dash-dotted lines show the results for the non-diffractive contribution from fitting the sum of the diffractive and non-diffractive contributions in the  $\ln M_X^2$  range delimited by the two vertical dashed lines.




# LRG vs $M_x$

$M_x$  data ( $M_N < 2.3$  GeV) normalised to LRG ( $M_N = m_p$ ): factor  $0.83 \pm 0.04$  determined via a global fit **estimates residual p-diss. background in  $M_x$  sample**

## ZEUS




## ZEUS



→ Overall agreement satisfactory


→ Different  $X_{IP}$  dependence ascribed to IR suppressed in  $M_x$  data

# ZEUS



# ZEUS LRG vs H1 LRG

## ZEUS



**ZEUS corrected to  $M_N < 1.6 \text{ GeV}$  with PYTHIA**

- Remaining normalisation difference of 13% (global fit) covered by uncertainty on p-diss. correction (8%) and relative normalisation uncertainty (7%)
- Shape agreement ok except low  $Q^2$

# Conclusions I

**ZEUS detector covers ~ 6.5 units of rapidity by high quality calorimetry**

**Rapidity Gap Selection &  $M_X$  Method used for Inclusive Diffractive Measurements**

**H1 detectors covers ~ 4.5 units by high quality calorimeter +**

**~ 3-4 units by particle detectors**

**Rapidity Gap Selection used only for Inclusive Diffractive Measurements**

**The agreement between H1 and ZEUS incl. diffractive measurements is fairly good**

**although worst than for  $F_2$ . Personal judgment: Main difficulty is due to the diffractive proton dissociation**

**Measurement of  $F_2^D$  is as fundamental as of  $F_2$ . Combined effort using all methods (including forwards protons) necessary.**

**Lesson for LHC: Extend good calorimeter coverage, build as many forwards detectors as possible**




## Leading Proton Spectrometer

Detector operation using Roman Pots


6 Ro-Pots equipped with micro-strip silicon detectors

pitch 115 micron

3 different strip orientations



## ZEUS 1994



Diffractive analysis using LPS detector allows :

Clean selection of the single diffraction processes (no proton dissociation)

Measurement of  $t$  in diffractive reactions


Good reconstruction of kinematical variables when combined with the central detector

Problem - limited statistics

# t dependence

LPS data


## ZEUS



Fit to  $e^{-b|t|} \rightarrow b = 7.0 \pm 0.4 \text{ GeV}^{-2}$


used in DPDF fits  
see talk by W. Slominski

## ZEUS



Lack of  $Q^2$  dependence and  $b$  much larger than in vector meson production  
 $\rightarrow$  features of a soft process

## ZEUS LPS vs H1 FPS



The cleanest possible comparison in principle...

...but large normalisation uncertainties  
(LPS: +11-7%, FPS: +/-10%)

New comparison plot available with HERA II FPS data!  
see talk by M. Kapishin

→ ZEUS and H1 proton-tagged data agree within normalisation uncertainties

# Conclusions

**Diffractive measurements at HERA achieved an impressive agreement between the different methods**


**Surprise of HERA:**

**Diffractive processes are an important part of short distance physics**


- ↳ implication on understanding of the QCD evolution**
- ↳ implication on understanding of confinement and nuclear structure**



### Pion from $\Delta$ decay at 250 GeV





# Pion from $\Delta$ decay at 100 GeV




# Hard Diffraction - the HERA surprise

Non-Diffractive Event



$$\tau_{\text{res}} \approx \frac{1}{\Delta E} \approx \frac{1}{m_p x} \approx 10 - 1000 \text{ fm}$$

Diffractive Event  
 expected before HERA  
 <0.01%, seen over 10%  
 at  $Q^2=10 \text{ GeV}^2$



Diffraction at HERA is so large because it is a shadow of DIS (i.e. inelastic processes) → **dipole picture**

$$\sigma_{\text{tot}}^{\gamma p} = \frac{1}{W^2} \text{Im} A_{\text{dip}}(W^2, t=0)$$