

LATEST NEWS ON DVCS FROM HERMES



*Caroline Riedl
EIC Meeting at BNL
August 25, 2009*




Latest News on DVCS from HERMES

- * Appetizer: HERA and HERMES
- * Motivation: Generalized Parton Distributions, orbital angular momentum and 3-dimensional nucleon structure
- * Measurements of azimuthal asymmetries in DVCS
 - * Beam helicity and charge asymmetries on hydrogen and nuclear targets
 - * Transverse target spin asymmetry on hydrogen
- * A Recoil Detector for HERMES

HERA @ DESY in Hamburg, Germany


- * Polarized electron/positron beam of 27.6 GeV and 40mA
- * Beam polarization 30...65%, 2 beam helicities
- * Proton beam of 920 GeV / 90 mA



C. Riedl (DESY), EIC Meeting at BNL, August 25, 2009


HERA @ DESY in Hamburg, Germany

- * Polarized electron/positron beam of 27.6 GeV and 40mA
- * Beam polarization 30...65%, 2 beam helicities
- * Proton beam of 920 GeV / 90 mA




C. Riedl (DESY), EIC Meeting at BNL, August 25, 2009


HERMES: HERa MEasurement of Spin




HERMES: HERa MEasurement of Spin



HERMES: HERa MEasurement of Spin




HERMES: HERa MEasurement of Spin



$\vec{H}, \vec{H}, \vec{D}$

H, D
 He, N, Ne, Kr, Xe

HERMES: HERa MEasurement of Spin



→
H, H, D

H, D
He, N, Ne, Kr, Xe

Tracking
momentum resolution:

$\leq 2\%$


angular resolution:

0.3...0.6 mrad

4

C. Riedl (DESY), EIC Meeting at BNL, August 25, 2009

HERMES: HERa MEasurement of Spin




→_H ↑_H →_D

H, D
He, N, Ne, Kr, Xe

Tracking
momentum resolution:
≤ 2%
angular resolution:
0.3...0.6 mrad

Particle Identification
electron ID: 98-99%
hadron contamination <1%
RICH: 2...15 GeV

HERMES: HERa MEasurement of Spin



→
H, H, D

H, D
He, N, Ne, Kr, Xe

Tracking
momentum resolution:
≤ 2%
angular resolution:
0.3...0.6 mrad

Particle IDentification
electron ID: 98-99%
hadron contamination <1%
RICH: 2...15 GeV

Deep Inelastic Scattering in ep collisions

Inclusive kinematics

Virtual photon virtuality:


$$Q^2 \equiv -q^2 := (k - k')^2$$
$$\stackrel{\text{lab}}{\approx} 4EE' \sin^2(\theta/2)$$

Bjorken scaling variable:

$$x_B := \frac{Q^2}{2Pq} \stackrel{\text{lab}}{=} \frac{Q^2}{2M(E - E')}$$

Invariant mass squared of X:

$$W^2 := (P + q)^2$$
$$\stackrel{\text{lab}}{=} M^2 + 2M(E - E') - Q^2$$



Deep Inelastic Scattering in ep collisions

Inclusive kinematics

Virtual photon virtuality:

$$Q^2 \equiv -q^2 := (k - k')^2$$

$$\stackrel{\text{lab}}{\approx} 4EE' \sin^2(\theta/2)$$


Björken scaling variable:

$$x_B := \frac{Q^2}{2Pq} \stackrel{\text{lab}}{=} \frac{Q^2}{2M(E - E')}$$

Invariant mass squared of X:

$$W^2 := (P + q)^2$$

$$\stackrel{\text{lab}}{=} M^2 + 2M(E - E') - Q^2$$



$$\sigma_{\text{DIS}} \sim \sum_X \left| \text{Diagram} \right|^2$$

Deep Inelastic Scattering in ep collisions

Inclusive kinematics

Virtual photon virtuality:

$$Q^2 \equiv -q^2 := (k - k')^2$$

$$\stackrel{\text{lab}}{\approx} 4EE' \sin^2(\theta/2)$$


Björken scaling variable:

$$x_B := \frac{Q^2}{2Pq} \stackrel{\text{lab}}{=} \frac{Q^2}{2M(E - E')}$$

Invariant mass squared of X:

$$W^2 := (P + q)^2$$

$$\stackrel{\text{lab}}{=} M^2 + 2M(E - E') - Q^2$$



$$\sigma_{\text{DIS}} \sim \sum_X \left| \text{Diagram} \right|^2$$

$$\left. \frac{d^2\sigma}{dE' d\Omega} \right|_{1\gamma^*} = \frac{\alpha^2}{2MQ^4} \frac{E'}{E} L_{\mu\nu} W^{\mu\nu}$$

Deep Inelastic Scattering in ep collisions

Inclusive kinematics

Virtual photon virtuality:

$$Q^2 \equiv -q^2 := (k - k')^2$$

$$\stackrel{\text{lab}}{\approx} 4EE' \sin^2(\theta/2)$$


Bjorken scaling variable:

$$x_B := \frac{Q^2}{2Pq} \stackrel{\text{lab}}{=} \frac{Q^2}{2M(E - E')}$$

Invariant mass squared of X:

$$W^2 := (P + q)^2$$

$$\stackrel{\text{lab}}{=} M^2 + 2M(E - E') - Q^2$$



$$\sigma_{\text{DIS}} \sim \sum_X \left| \text{Diagram} \right|^2$$

$$\left. \frac{d^2\sigma}{dE' d\Omega} \right|_{1\gamma^*} = \frac{\alpha^2}{2MQ^4} \frac{E'}{E} L_{\mu\nu} W^{\mu\nu}$$

Semi-inclusive kinematics
involve kinematics of produced hadrons

Exclusive kinematics
complete spectrum of X known

The Composition of the Nucleon's Spin

$$\frac{1}{2}\hbar = J_{\text{quarks}} + J_{\text{gluons}}$$



The Composition of the Nucleon's Spin

$$\frac{1}{2}\hbar = J_{\text{quarks}} + J_{\text{gluons}}$$

$$\frac{1}{2}\Delta\Sigma + L_q$$



The Composition of the Nucleon's Spin

$$\frac{1}{2}\hbar = J_{\text{quarks}} + J_{\text{gluons}}$$

(?)

$$\frac{1}{2}\Delta\Sigma + L_q$$



The Composition of the Nucleon's Spin

$$\frac{1}{2}\hbar = J_{\text{quarks}} + J_{\text{gluons}}$$

(?)

$$\frac{1}{2}\Delta\Sigma + L_q$$



* Spin of quarks: $\Delta\Sigma \approx 1/3$ from DIS and SIDIS

Hermes Phys. Rev. **D75** (2007) 012007: $\Delta\Sigma = 0.330 \pm 0.011(\text{theo}) \pm 0.025(\text{exp}) \pm 0.028(\text{evol})$

The Composition of the Nucleon's Spin

$$\frac{1}{2}\hbar = J_{\text{quarks}} + J_{\text{gluons}}$$

(?)

$$\frac{1}{2}\Delta\Sigma + L_q$$



- * Spin of quarks: $\Delta\Sigma \approx 1/3$ from DIS and SIDIS

Hermes Phys. Rev. **D75** (2007) 012007: $\Delta\Sigma = 0.330 \pm 0.011(\text{theo}) \pm 0.025(\text{exp}) \pm 0.028(\text{evol})$

- * Orbital angular momentum of quarks: L_q ?

$$J_q = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H^q(x, \xi, t) + E^q(x, \xi, t)]$$

Ji relation

Ji, PRL **78** (1997) 610

The Composition of the Nucleon's Spin

$$\frac{1}{2}\hbar = J_{\text{quarks}} + J_{\text{gluons}}$$

(?)

$$\frac{1}{2}\Delta\Sigma + L_q$$



- * Spin of quarks: $\Delta\Sigma \approx 1/3$ from DIS and SIDIS

Hermes Phys. Rev. **D75** (2007) 012007: $\Delta\Sigma = 0.330 \pm 0.011(\text{theo}) \pm 0.025(\text{exp}) \pm 0.028(\text{evol})$

- * Orbital angular momentum of quarks: L_q ?


$$J_q = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H^q(x, \xi, t) + E^q(x, \xi, t)]$$

Ji relation

Ji, PRL **78** (1997) 610

Generalized Parton Distributions (GPDs)

Generalized Parton Distributions




- * Access through hard exclusive reactions
 - * t : squared momentum transfer to target
 - * x : average longitudinal momentum fraction of quark before/after
 - * ξ : $1/2$ difference
- * GPDs $F^q(x, \xi, t)$ parameterize the non-perturbative nucleon structure
 - contain info on parton-parton correlations

* 4 GPDs that conserve quark chirality (Spin-1/2 target, leading-twist)

nucleon helicity ↓	quark helicity independent	quark helicity dependent
	photon: $J^P=1^-$ (DVCS)	
conserved	H	H-tilde
flipped	E	E-tilde
	$J^P=1^-$ mesons	$J^P=0^-$ mesons

GPDs: A unifying picture of nucleon structure





$$H^q(x, 0, 0) = q(x)$$

forward limit $\xi=0, t=0$

$$\int_{-1}^1 dx H^q(x, \xi, t) = F_1^q(t)$$

moments of GPDs

GPDs: A unifying picture of nucleon structure





$$H^q(x, 0, 0) = q(x)$$

forward limit $\xi=0, t=0$

$$\int_{-1}^1 dx H^q(x, \xi, t) = F_1^q(t)$$

moments of GPDs

GPDs: A unifying picture of nucleon structure





$$H^q(x, 0, 0) = q(x)$$

forward limit $\xi=0, t=0$

$$\int_{-1}^1 dx H^q(x, \xi, t) = F_1^q(t)$$

moments of GPDs

GPDs: A unifying picture of nucleon structure



$$H^q(x, 0, 0) = q(x)$$

forward limit $\xi=0, t=0$

$$\int_{-1}^1 dx H^q(x, \xi, t) = F_1^q(t)$$


moments of GPDs

GPDs: A unifying picture of nucleon structure



Parton Distribution
Functions:
longitudinal momentum



GPDs
 $F^q(x, \xi, t)$



Form Factors:
transverse position



$$H^q(x, 0, 0) = q(x)$$

forward limit $\xi=0, t=0$

$$\int_{-1}^1 dx H^q(x, \xi, t) = F_1^q(t)$$


moments of GPDs

GPDs: A unifying picture of nucleon structure




Parton Distribution
Functions:
longitudinal momentum



GPDs
 $F^q(x, \xi, t)$



Form Factors:
transverse position



$$H^q(x, 0, 0) = q(x)$$

forward limit $\xi=0, t=0$

$$\int_{-1}^1 dx H^q(x, \xi, t) = F_1^q(t)$$


moments of GPDs

GPDs: A unifying picture of nucleon structure




Parton Distribution Functions:
longitudinal momentum



GPDs
 $F^q(x, \xi, t)$



Form Factors:
transverse position



$$H^q(x, 0, 0) = q(x)$$

forward limit $\xi=0, t=0$

$$\int_{-1}^1 dx H^q(x, \xi, t) = F_1^q(t)$$

moments of GPDs


“Nucleon Tomography”

GPDs: A unifying picture of nucleon structure



Parton Distribution
Functions:
longitudinal momentum



GPDs
 $F^q(x, \xi, t)$



Form Factors:
transverse position



$$H^q(x, 0, 0) = q(x)$$

forward limit $\xi=0, t=0$

$$\int_{-1}^1 dx H^q(x, \xi, t) = F_1^q(t)$$

moments of GPDs


“Nucleon Tomography”

GPDs: A unifying picture of nucleon structure




Parton Distribution Functions:
longitudinal momentum



GPDs
 $F^q(x, \xi, t)$



Form Factors:
transverse position



$$H^q(x, 0, 0) = q(x)$$

forward limit $\xi=0, t=0$

$$\int_{-1}^1 dx H^q(x, \xi, t) = F_1^q(t)$$

moments of GPDs

“Nucleon Tomography”

Deeply Virtual Compton Scattering (DVCS)

Deeply Virtual Compton Scattering (DVCS)

Reminder: inclusive DIS

$$\sigma_{\text{DIS}} \sim \sum_X \left| \begin{array}{c} \text{wavy line} \\ \text{circle} \\ \text{line} \end{array} \right|^2$$

$\sim \text{Im} \left(\begin{array}{c} \text{wavy line } q \\ \text{circle} \\ \text{wavy line } q \end{array} \right)$

↑
optical theorem


forward
Compton scattering
amplitude

The image contains two Feynman diagrams. The top diagram shows a wavy line (photon) entering a circle (target) from the left, and a line (electron) exiting to the right. A vertical line labeled 'X' is attached to the right side of the circle. The bottom diagram shows a wavy line labeled 'q' entering a circle from the left, and another wavy line labeled 'q' exiting to the right. A blue speech bubble points to this diagram with the text 'forward Compton scattering amplitude'.


Deeply Virtual Compton Scattering (DVCS)

DVCS = hard electroproduction of a real photon


Reminder: inclusive DIS

$$\sigma_{\text{DIS}} \sim \sum_X \left| \text{Diagram} \right|^2$$


optical theorem \uparrow $\sim \text{Im} \left(\text{Diagram} \right)$




forward Compton scattering amplitude



Skewing
due to
mass difference
of photons


the 2 partons have
different longitudinal
momenta!
Skewness $\xi \neq 0$
Leading order: $\xi \approx x_B / (2 - x_B)$




Deeply Virtual Compton Scattering (DVCS)

DVCS = hard electroproduction of a real photon


Reminder: inclusive DIS

$$\sigma_{\text{DIS}} \sim \sum_X \left| \text{Diagram} \right|^2$$


optical theorem \uparrow $\sim \text{Im} \left(\text{Diagram} \right)$




forward Compton scattering amplitude



Skewing
due to
mass difference
of photons

the 2 partons have
different longitudinal
momenta!
Skewness $\xi \neq 0$
Leading order: $\xi \approx x_B / (2 - x_B)$

Forward limit:
 $\xi=0, t=0$



The $\gamma^*N \rightarrow \gamma N$ Cross Section

$$\sigma_{\gamma^* \gamma N} = \left| \text{DVCS} + \text{Bethe-Heitler (BH)} \right|^2$$

DVCS
Bethe-Heitler (BH)

Bjorken limit:
 large Q^2 (small distances)
 large energy of γ^* (small times)
 small t , fixed x_B

$$= |\tau_{\text{DVCS}}|^2 + |\tau_{\text{BH}}|^2 + \tau_{\text{DVCS}}\tau_{\text{BH}}^* + \tau_{\text{DVCS}}^*\tau_{\text{BH}}$$


DVCS-BH
 interference term \mathcal{I}

Contribution at colliders.
 Fixed target:
 $|\tau_{\text{DVCS}}|^2 \ll |\tau_{\text{BH}}|^2$

exactly calculable in QED
 given the nucleon elastic form
 factors F_1 and F_2

Azimuthal Dependences in $\gamma^*N \rightarrow \gamma N$

- Unpolarized target
- Lepton beam with charge C_B and polarization P_B



Fourier expansion in azimuthal angle ϕ


$$|\mathcal{T}_{\text{BH}}|^2 = \frac{K_{\text{BH}}}{\mathcal{P}_1(\phi)\mathcal{P}_2(\phi)} \sum_{n=0}^2 c_n^{\text{BH}} \cos(n\phi)$$

$$|\mathcal{T}_{\text{DVCS}}|^2 = K_{\text{DVCS}} \left[\sum_{n=0}^2 c_n^{\text{DVCS}} \cos(n\phi) + P_B \sum_{n=1}^1 s_n^{\text{DVCS}} \sin(n\phi) \right]$$

$$\mathcal{I} = \frac{C_B K_{\mathcal{I}}}{\mathcal{P}_1(\phi)\mathcal{P}_2(\phi)} \left[\sum_{n=0}^3 c_n^{\mathcal{I}} \cos(n\phi) + P_B \sum_{n=1}^2 s_n^{\mathcal{I}} \sin(n\phi) \right]$$

Azimuthal Dependences in $\gamma^*N \rightarrow \gamma N$

- Unpolarized target
- Lepton beam with charge C_B and polarization P_B



Fourier expansion in azimuthal angle ϕ

$$|\mathcal{T}_{\text{BH}}|^2 = \frac{K_{\text{BH}}}{\mathcal{P}_1(\phi)\mathcal{P}_2(\phi)} \sum_{n=0}^2 c_n^{\text{BH}} \cos(n\phi)$$


$$|\mathcal{T}_{\text{DVCS}}|^2 = K_{\text{DVCS}} \left[\sum_{n=0}^2 c_n^{\text{DVCS}} \cos(n\phi) + P_B \sum_{n=1}^1 s_n^{\text{DVCS}} \sin(n\phi) \right]$$

$$\mathcal{I} = \frac{C_B K_{\mathcal{I}}}{\mathcal{P}_1(\phi)\mathcal{P}_2(\phi)} \left[\sum_{n=0}^3 c_n^{\mathcal{I}} \cos(n\phi) + P_B \sum_{n=1}^2 s_n^{\mathcal{I}} \sin(n\phi) \right]$$

Bethe-Heitler propagators $\mathcal{P}(\phi)$

Azimuthal Dependences in $\gamma^*N \rightarrow \gamma N$

- Unpolarized target
- Lepton beam with charge C_B and polarization P_B



Fourier expansion in azimuthal angle ϕ

$$|\mathcal{T}_{\text{BH}}|^2 = \frac{K_{\text{BH}}}{\mathcal{P}_1(\phi)\mathcal{P}_2(\phi)} \sum_{n=0}^2 c_n^{\text{BH}} \cos(n\phi)$$

$$|\mathcal{T}_{\text{DVCS}}|^2 = K_{\text{DVCS}} \left[\sum_{n=0}^2 c_n^{\text{DVCS}} \cos(n\phi) + P_B \sum_{n=1}^1 s_n^{\text{DVCS}} \sin(n\phi) \right]$$

$$\mathcal{I} = \frac{C_B K_{\mathcal{I}}}{\mathcal{P}_1(\phi)\mathcal{P}_2(\phi)} \left[\sum_{n=0}^3 c_n^{\mathcal{I}} \cos(n\phi) + P_B \sum_{n=1}^2 s_n^{\mathcal{I}} \sin(n\phi) \right]$$

Bethe-Heitler propagators $\mathcal{P}(\phi)$

Wanted:
Fourier coefficients
 s_n and c_n
of BH, DVCS, and \mathcal{I} terms


Measured Azimuthal Asymmetries in DVCS

Born cross-section:

$$\sigma(\phi; P_B, C_B) = \sigma_{UU}(\phi) \cdot [1 + P_B \mathcal{A}_{LU}^{DVCS}(\phi) + C_B P_B \mathcal{A}_{LU}^I(\phi) + C_B \mathcal{A}_C(\phi)]$$

Beam helicity asymmetries

Beam charge asymmetry



Old approach at HERMES
and CLAS: single charge BSA

BSA:
projects out imaginary
part of τ_{DVCS}

BCA:
projects out real part
of τ_{DVCS}

$$\mathcal{A}_{LU}(\phi) \equiv \frac{d\sigma^{\rightarrow} - d\sigma^{\leftarrow}}{d\sigma^{\rightarrow} + d\sigma^{\leftarrow}}$$


no separate access
to s_1^{\uparrow} and s_1^{DVCS}

$$\mathcal{A}_C(\phi) \equiv \frac{d\sigma^+ - d\sigma^-}{d\sigma^+ + d\sigma^-}$$

Measured Azimuthal Asymmetries in DVCS

Born cross-section:

$$\sigma(\phi; P_B, C_B) = \sigma_{UU}(\phi) \cdot [1 + P_B \mathcal{A}_{LU}^{\text{DVCS}}(\phi) + C_B P_B \mathcal{A}_{LU}^{\text{I}}(\phi) + C_B \mathcal{A}_C(\phi)]$$



Beam helicity asymmetries

Beam charge asymmetry

Old approach at HERMES and CLAS: single charge BSA

BSA:
projects out imaginary part of τ_{DVCS}

BCA:
projects out real part of τ_{DVCS}

$$\mathcal{A}_{LU}(\phi) \equiv \frac{d\sigma^{\rightarrow} - d\sigma^{\leftarrow}}{d\sigma^{\rightarrow} + d\sigma^{\leftarrow}}$$

$$\mathcal{A}_C(\phi) \equiv \frac{d\sigma^{+} - d\sigma^{-}}{d\sigma^{+} + d\sigma^{-}}$$

no separate access to s_1^{\uparrow} and s_1^{DVCS}

New approach at HERMES:
 s_1^{\uparrow} and s_1^{DVCS} can be disentangled

Charge difference BSA:

Charge average BSA:

$$\mathcal{A}_{LU}^{\text{I}}(\phi) \equiv \frac{(d\sigma^{+\rightarrow} - d\sigma^{+\leftarrow}) - (d\sigma^{-\rightarrow} - d\sigma^{-\leftarrow})}{(d\sigma^{+\rightarrow} + d\sigma^{+\leftarrow}) + (d\sigma^{-\rightarrow} + d\sigma^{-\leftarrow})}$$

$$\mathcal{A}_{LU}^{\text{DVCS}}(\phi) \equiv \frac{(d\sigma^{+\rightarrow} - d\sigma^{+\leftarrow}) + (d\sigma^{-\rightarrow} - d\sigma^{-\leftarrow})}{(d\sigma^{+\rightarrow} + d\sigma^{+\leftarrow}) + (d\sigma^{-\rightarrow} + d\sigma^{-\leftarrow})}$$

Relation: asymmetries \leftrightarrow Fourier coefficients

Beam helicity asymmetries:

$$A_{LU}^I(\phi) = \frac{1}{D(\phi)} \cdot \frac{x_B}{Q^2} [s_1^I \sin(\phi) + s_2^I \sin(2\phi)]$$

$$A_{LU}^{DVCS}(\phi) = \frac{1}{D(\phi)} \cdot \frac{x_B^2 t \mathcal{P}_1(\phi) \mathcal{P}_2(\phi)}{Q^2} s_1^{DVCS} \sin(\phi)$$

Beam charge asymmetry:

$$A_C(\phi) = -\frac{1}{D(\phi)} \cdot \frac{x_B}{y} [c_0^I + c_1^I \cos(\phi) + c_2^I \cos(2\phi) + c_3^I \cos(3\phi)]$$

Relation: asymmetries \leftrightarrow Fourier coefficients

Beam helicity asymmetries:

$$A_{LU}^I(\phi) = \frac{1}{D(\phi)} \cdot \frac{x_B}{Q^2} [s_1^I \sin(\phi) + s_2^I \sin(2\phi)]$$

$$A_{LU}^{DVCS}(\phi) = \frac{1}{D(\phi)} \cdot \frac{x_B^2 t \mathcal{P}_1(\phi) \mathcal{P}_2(\phi)}{Q^2} s_1^{DVCS} \sin(\phi)$$

Beam charge asymmetry:

$$A_C(\phi) = -\frac{1}{D(\phi)} \cdot \frac{x_B}{y} [c_0^I + c_1^I \cos(\phi) + c_2^I \cos(2\phi) + c_3^I \cos(3\phi)]$$

Additional ϕ dependence
in denominator through
BH propagators

$$D(\phi) = \frac{\sum_{n=0}^2 c_n^{BH} \cos(n\phi)}{(1 + \epsilon^2)^2} + \frac{x_B^2 t \mathcal{P}_1(\phi) \mathcal{P}_2(\phi)}{Q^2} \sum_{n=0}^2 c_n^{DVCS} \cos(n\phi)$$

From Azimuthal Asymmetries to GPDs

- * To obtain Fourier coefficients = asymmetry amplitudes:
 - Combine data with different beam charges and helicities and **fit all amplitudes simultaneously**

- * Compton Form Factors (CFFs) $\mathcal{F}(\xi, t) = \sum_q \int_{-1}^1 dx C_q^\mp(\xi, x) F^q(x, \xi, t)$
 - * Define linear combination of CFFs: $\mathcal{C}_{\text{unp}}^{\mathcal{I}} = F_1 \mathcal{H} + \xi(F_1 + F_2) \tilde{\mathcal{H}} - \frac{t}{4M^2} F_2 \mathcal{E}$
 - * $F_1(t), F_2(t)$: Dirac, Pauli nucleonic form factors

twist-2 GPD

- * Leading twist level (twist-2):

$$c_1^{\mathcal{I}} \propto \frac{\sqrt{-t}}{Q} \Re [\mathcal{C}_{\text{unp}}^{\mathcal{I}}] \propto -\frac{Q}{\sqrt{-t}} c_0^{\mathcal{I}}$$


BCA
constant term

$$s_1^{\mathcal{I}} \propto \frac{\sqrt{-t}}{Q} \Im [\mathcal{C}_{\text{unp}}^{\mathcal{I}}]$$

BSA

Holographic Principle / Femtophotography

- Wanted: 3-dim spatial picture
 - $(FT)^{-1}$ of diffraction pattern, given amplitude $\tau = Ae^{i\varphi}$
 - Need both magnitude A & phase φ
 - Usually $|\tau|^2$ is measured, phase is lost (e.g. PDFs)





- Holography technique:
 - Known BH process as reference amplitude that magnifies DVCS effect
 - Measure phase of DVCS through its interference with BH

DVCS at HERMES 1996-2005 (w/o Recoil)

Detected particles:
electron and photon

Missing mass technique for
 $ep \rightarrow eX\gamma$
 $M_X^2 = (p+q-p_Y)^2$



- e⁺ data
- e⁻ data
- MC sum
- - - elastic BH
- associated BH
- semi-inclusive

$X=p$

resonant excitation: $X=\Delta^+$

$X=\pi^0+\dots$

$\rho\pi^0$

$n\pi^+$



hydrogen target:
25k events
(400 pb⁻¹)

unpolarized
deuterium:
15k events
(300 pb⁻¹)

DVCS at HERMES 1996-2005 (w/o Recoil)

Detected particles:
electron and photon

Missing mass technique for
 $ep \rightarrow eX\gamma$
 $M_X^2 = (p+q-p_Y)^2$



- e⁺ data
- e⁻ data
- MC sum
- ⋯ elastic BH
- associated BH
- ⋯ semi-inclusive

$X=p$

resonant excitation: $X=\Delta^+$

$X=\pi^0+\dots$

$\rho\pi^0$

$n\pi^+$



hydrogen target:
25k events
(400 pb⁻¹)

unpolarized
deuterium:
15k events
(300 pb⁻¹)

DVCS at HERMES 1996-2005 (w/o Recoil)

Detected particles:
electron and photon

Missing mass technique for
 $ep \rightarrow eX\gamma$
 $M_X^2 = (p+q-p_Y)^2$



$X=p$

resonant
excitation: $X=\Delta^+$

$X=\pi^0+\dots$

$\rho\pi^0$

$n\pi^+$

hydrogen target:
25k events
(400 pb^{-1})



unpolarized
deuterium:
15k events
(300 pb^{-1})

C. Riedl (DESY), EIC Meeting at BNL, August 25, 2009

DVCS at HERMES 1996-2005 (w/o Recoil)

Detected particles:
electron and photon

Missing mass technique for
 $ep \rightarrow eX\gamma$
 $M_X^2 = (p+q-p_Y)^2$



- e⁺ data
- e⁻ data
- MC sum
- ⋯ elastic BH
- associated BH
- ⋯ semi-inclusive

$X=p$

resonant excitation: $X=\Delta^+$

$X=\pi^0+\dots$


$\rho\pi^0$

$n\pi^+$

hydrogen target:
25k events
(400 pb⁻¹)

unpolarized
deuterium:
15k events
(300 pb⁻¹)

HERMES: BSA from \mathcal{I} on hydrogen




$$\propto \Im [F_1 \mathcal{H}]$$

← Higher twist (twist-3)

← Fraction of resonant excitation

(*) Phys.Rev. **D60** (1999) and Prog.Nucl.Phys. **47** (2001) 401 17

HERMES: BSA from \mathcal{I} on hydrogen



D-term to restore
 polynomiality


$$\propto \Im [F_1 \mathcal{H}]$$

← Higher twist (twist-3)

← Fraction of resonant
 excitation

(*) Phys.Rev. **D60** (1999) and Prog.Nucl.Phys. **47** (2001) 401 17

HERMES: BSA from \mathcal{I} on hydrogen



D-term to restore polynomiality

Bands by variation of skewness parameters


$$\propto \Im [F_1 \mathcal{H}]$$

← Higher twist (twist-3)

← Fraction of resonant excitation

(*) Phys.Rev. **D60** (1999) and Prog.Nucl.Phys. **47** (2001) 401 17

HERMES: BSA from $|\tau_{DVCS}|^2$ on hydrogen



$$\propto \left[\mathcal{H}\mathcal{H}^* + \tilde{\mathcal{H}}\tilde{\mathcal{H}}^* \right]$$

bilinear combination of CFFs


← Higher twist (twist-3)

GPD model: Vanderhaeghen, Guichon, Guidal (“VGG”)


Phys.Rev. **D60** (1999) and Prog.Nucl.Phys. **47** (2001) 401

Based on double distribution ansatz: “ $DF(x,\xi) \otimes \text{profile}(t)$ ”.

Twist-3 included



HERMES: BCA on hydrogen



constant term:


$$\propto -A_C^{\cos \phi}$$

$$\propto \Re [F_1 \mathcal{H}]$$

← Higher twist (twist-3)

← Gluon leading twist

HERMES: BCA on hydrogen



constant term:

$$\propto -A_C^{\cos \phi}$$

$$\propto \Re [F_1 \mathcal{H}]$$


← Higher twist (twist-3)

← Gluon leading twist

VGG with
 D-term
 disfavored


DVCS on Nuclear Targets

- * How does the nuclear environment modify parton-parton correlations?
- * How do nucleon properties change in the nuclear medium?




DVCS on Nuclear Targets

- * How does the nuclear environment modify parton-parton correlations?
- * How do nucleon properties change in the nuclear medium?




- * DVCS in coherent region: new insights into 'generalized EMC effect'?



- Nuclear GPDs \neq GPDs of free nucleon
- Enhancement of effect when leaving forward limit?
 - caused by transverse motion of partons in nuclei?
 - important role of mesonic degrees of freedom?
 - manifest in strong increase of real part of τ_{DVCS} with atomic mass number A ?


DVCS on Nuclear Targets

- * How does the nuclear environment modify parton-parton correlations?
- * How do nucleon properties change in the nuclear medium?



(Bethe-Heitler)

- * DVCS in coherent region: new insights into 'generalized EMC effect'?




- Nuclear GPDs \neq GPDs of free nucleon
- Enhancement of effect when leaving forward limit?
 - caused by transverse motion of partons in nuclei?
 - important role of mesonic degrees of freedom?
 - manifest in strong increase of real part of τ_{DVCS} with atomic mass number A ?

**HERMES
measurements
on nuclear
targets**

Target	spin	L (pb^{-1})
H	1/2	227
He	0	32
N	1	51
Ne	0	86
Kr	0	77
Xe	0, 1/2, 3/2	47


DVCS on Nuclear Targets

- * How does the nuclear environment modify parton-parton correlations?
- * How do nucleon properties change in the nuclear medium?



(Bethe-Heitler)

- * DVCS in coherent region: new insights into 'generalized EMC effect'?






- Nuclear GPDs \neq GPDs of free nucleon
- Enhancement of effect when leaving forward limit?
 - caused by transverse motion of partons in nuclei?
 - important role of mesonic degrees of freedom?
 - manifest in strong increase of real part of τ_{DVCS} with atomic mass number A ?

HERMES measurements on nuclear targets


Target	spin	L (pb^{-1})
H	1/2	227
He	0	32
N	1	51
Ne	0	86
Kr	0	77
Xe	0, 1/2, 3/2	47

+ deuterium, spin-1, 300 pb^{-1}



HERMES: BCAs on hydrogen and deuterium




HERMES: BCAs on hydrogen and deuterium



low t: coherent





HERMES: BCAs on hydrogen and deuterium




low t: coherent

high t: incoherent




DVCS at HERMES: Nuclear mass dependence



Select for each target two samples (t-cutoffs):

- **coherent enriched**
($\approx 65\%$ coherent fraction)
- **incoherent enriched**
($\approx 60\%$ incoherent fraction)


DVCS at HERMES: Nuclear mass dependence




Select for each target
two samples (t-cutoffs):

- **coherent enriched**
($\approx 65\%$ coherent fraction)
- **incoherent enriched**
($\approx 60\%$ incoherent fraction)


BCA vs. A



BSA ratio: nuclear / free proton vs. A




DVCS at HERMES: Nuclear mass dependence




- Select for each target two samples (t-cutoffs):
- **coherent enriched**
($\approx 65\%$ coherent fraction)
 - **incoherent enriched**
($\approx 60\%$ incoherent fraction)

No nuclear mass dependence of BCA and BSA observed within uncertainties
 \Rightarrow no enhancement of τ_{DVCS}

BCA vs. A




BSA ratio: nuclear / free proton vs. A



Exclusivity at HERMES in a Nutshell

GPD access at HERMES:




nucleon helicity	quark helicity independent	quark helicity dependent
	photon: $J^P=1^-$ (DVCS)	
conserved	H : A_C, A_{LU}, A_{UT}	H-tilde : $A_{UL}, [A_{UT}]$
flipped	E : A_{UT}	E-tilde : $[A_{UT}]$
	$J^P=1^-$ mesons	$J^P=0^-$ mesons

$$J_q = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H^q(x, \xi, t) + E^q(x, \xi, t)]$$

Exclusivity at HERMES in a Nutshell

GPD access at HERMES:




nucleon helicity	quark helicity independent	quark helicity dependent
	photon: $J^P=1^-$ (DVCS)	
conserved	H : A_C, A_{LU}, A_{UT}	H-tilde : $A_{UL}, [A_{UT}]$
flipped	E : A_{UT}	E-tilde : $[A_{UT}]$
	$J^P=1^-$ mesons	$J^P=0^-$ mesons

$$J_q = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H^q(x, \xi, t) + E^q(x, \xi, t)]$$

Exclusivity at HERMES in a Nutshell

GPD access at HERMES:




nucleon helicity	quark helicity independent	quark helicity dependent
	photon: $J^P=1^-$ (DVCS)	
conserved	H : A_C, A_{LU}, A_{UT}	H-tilde : $A_{UL}, [A_{UT}]$
flipped	E : A_{UT}	E-tilde : $[A_{UT}]$
	$J^P=1^-$ mesons	$J^P=0^-$ mesons

$$J_q = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H^q(x, \xi, t) + E^q(x, \xi, t)]$$

Exclusivity at HERMES in a Nutshell

GPD access at HERMES:



nucleon helicity	quark helicity independent	quark helicity dependent
	photon: $J^P=1^-$ (DVCS)	
conserved	H : A_C, A_{LU}, A_{UT}	H-tilde : $A_{UL}, [A_{UT}]$
flipped	E : A_{UT}	E-tilde : $[A_{UT}]$
	$J^P=1^-$ mesons	$J^P=0^-$ mesons


vector mesons
 ρ^0, ω, ϕ

pseudoscalar mesons
 π^+, η

$$J_q = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H^q(x, \xi, t) + E^q(x, \xi, t)]$$

Exclusivity at HERMES in a Nutshell

GPD access at HERMES:



nucleon helicity	quark helicity independent	quark helicity dependent
	photon: $J^P=1^-$ (DVCS)	
conserved	H : A_C, A_{LU}, A_{UT}	H-tilde : $A_{UL}, [A_{UT}]$
flipped	E : A_{UT}	E-tilde : $[A_{UT}]$
	$J^P=1^-$ mesons	$J^P=0^-$ mesons


vector mesons
 ρ^0, ω, ϕ

pseudoscalar mesons
 π^+, η

$$J_q = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H^q(x, \xi, t) + E^q(x, \xi, t)]$$

Exclusivity at HERMES in a Nutshell

GPD access at HERMES:



nucleon helicity	quark helicity independent	quark helicity dependent
	photon: $J^P=1^-$ (DVCS)	
conserved	H : A_C, A_{LU}, A_{UT}	H-tilde : $A_{UL}, [A_{UT}]$
flipped	E : A_{UT}	E-tilde : $[A_{UT}]$
	$J^P=1^-$ mesons	$J^P=0^-$ mesons


vector mesons
 ρ^0, ω, ϕ

pseudoscalar mesons
 π^+, η

$$J_q = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H^q(x, \xi, t) + E^q(x, \xi, t)]$$

DVCS Transverse Target Spin Asymmetry $\mathcal{A}_{UT}(\phi, \phi_s)$

- * \mathcal{A}_{UT} : the only DVCS asymmetry on the proton for which **GPD E is not suppressed**
(Hall-A: BSA on neutron)




- * **HERMES**: transversely polarized hydrogen, 170 pb^{-1} , 2 beam charges
 - Separation of DVCS and interference terms possible

$$A_{UT}^{\mathcal{I}}(\phi, \phi_s) \propto [d\sigma^+(\phi, \phi_s) - d\sigma^-(\phi, \phi_s)] - [d\sigma^+(\phi, \phi_s + \pi) - d\sigma^-(\phi, \phi_s + \pi)]$$

$$A_{UT}^{\mathcal{I}}(\phi, \phi_s) \propto \text{Im}(F_2 \mathcal{H} - F_1 \mathcal{E}) \sin(\phi - \phi_s) \cos \phi + \text{Im}(F_2 \tilde{\mathcal{H}} - (F_1 + \xi F_2) \tilde{\mathcal{E}}) \cos(\phi - \phi_s) \sin \phi$$

DVCS Transverse Target Spin Asymmetry $\mathcal{A}_{UT}(\phi, \phi_s)$

- * \mathcal{A}_{UT} : the only DVCS asymmetry on the proton for which **GPD E is not suppressed**
(Hall-A: BSA on neutron)



- * **HERMES**: transversely polarized hydrogen, 170 pb^{-1} , 2 beam charges
 - Separation of DVCS and interference terms possible


$$\mathcal{A}_{UT}^{\text{DVCS}}(\phi, \phi_s)$$

also sensitive to GPD E
(bilinear combination)

$$\mathcal{A}_{UT}^{\mathcal{I}}(\phi, \phi_s) \propto [d\sigma^+(\phi, \phi_s) - d\sigma^-(\phi, \phi_s)] - [d\sigma^+(\phi, \phi_s + \pi) - d\sigma^-(\phi, \phi_s + \pi)]$$


$$\begin{aligned} \mathcal{A}_{UT}^{\mathcal{I}}(\phi, \phi_s) &\propto \text{Im}(F_2 \mathcal{H} - F_1 \mathcal{E}) \sin(\phi - \phi_s) \cos \phi \\ &+ \text{Im}(F_2 \tilde{\mathcal{H}} - (F_1 + \xi F_2) \tilde{\mathcal{E}}) \cos(\phi - \phi_s) \sin \phi \end{aligned}$$

HERMES DVCS \mathcal{A}_{UT} Amplitudes




Model: VGG with variation of J_u , while $J_d=0$

HERMES DVCS \mathcal{A}_{UT} Amplitudes



Model: VGG with variation of J_u , while $J_d=0$



HERMES 2006-2007: Recoil Detector



^1H (^2H): factor of 1.6 (0.5)
more than 1996-2005

- Purpose:**
- ★ To tag exclusive events
 - Identify recoiling target proton
 - Identify particles from background processes

The Silicon Strip Detector (SSD)



$E_{\text{kin}} \approx 8 \text{ MeV}$
for protons

- Purpose:

- ▶ Track reconstruction
- ▶ Momenta: $> 125 \text{ MeV}/c$
- ▶ PID for low and medium momenta

- 2 layers of 16 double-sided sensors

- ▶ $(10 \text{ cm} \times 10 \text{ cm})$ active area
- ▶ $300 \mu\text{m}$ thickness


- Inside accelerator vacuum, 5 cm close to electron beam

to reach as low t as possible

The Scintillating Fiber Tracker (SFT)



- Purpose:
 - ▶ Track reconstruction
 - ▶ Momenta: 250-1400 MeV/c (protons)
 - ▶ PID for medium and high momenta
- 2 Barrels with each 4 layers of scintillating fibers
- Per Barrel:
 - ▶ 2 parallel layers
 - ▶ 2 stereo-layers
 - ▶ Stereo angle: 10°



The Photon Detector (PD)




- Purpose:
 - ▶ Detection of photons from resonance decay $\Delta^+ \rightarrow p\pi^0$
 - ▶ PID for $p > 600 \text{ MeV}/c$
- 3 layers of tungsten/scintillator sandwich
 - ▶ 1 layer parallel to beam axis
 - ▶ 2 layers under $+45^\circ/-45^\circ$ angles




Tracking with the Recoil Detector

Energy Deposit in the SSD




Transverse View of Detector




- **Low-energy protons:** momentum $\propto (\sum_i \Delta E_i)^{-1}$
- **Medium-energy protons:** momentum $\propto (\frac{dE}{dx})^{-1}$ (Bethe-Bloch)
- **Higher-energy particles (protons/pions):** momentum $\propto eB\rho$

Tracking with the Recoil Detector

Energy Deposit in the SSD




Transverse View of Detector




- **Low-energy protons:** momentum $\propto (\sum_i \Delta E_i)^{-1}$
- **Medium-energy protons:** momentum $\propto (\frac{dE}{dx})^{-1}$ (Bethe-Bloch)
- **Higher-energy particles (protons/pions):** momentum $\propto eB\rho$

Tracking with the Recoil Detector

Energy Deposit in the SSD




Transverse View of Detector




- **Low-energy protons:** momentum $\propto (\sum_i \Delta E_i)^{-1}$
- **Medium-energy protons:** momentum $\propto (\frac{dE}{dx})^{-1}$ (Bethe-Bloch)
- **Higher-energy particles (protons/pions):** momentum $\propto eB\rho$

Tracking with the Recoil Detector

Energy Deposit in the SSD




Transverse View of Detector




- **Low-energy protons:** momentum $\propto (\sum_i \Delta E_i)^{-1}$
- **Medium-energy protons:** momentum $\propto (\frac{dE}{dx})^{-1}$ (Bethe-Bloch)
- **Higher-energy particles (protons/pions):** momentum $\propto eB\rho$

Reconstructed Momenta and Angles



- Recoiling target protons
 - ▶ Large θ -angles $\lesssim 90^\circ$
 - ▶ Small momenta < 1 GeV/c
- Azimuthal ϕ coverage: 76%




- ΔE accounted for in track fitting
 $\Rightarrow \Delta p/p$ improvement




Proton / Pion Separation with the Recoil

- $p < 600 \text{ MeV}/c$: SSD + SFT (6 layers)
- $p > 600 \text{ MeV}/c$: include PD
- Log-likelihood formalism:


$$\text{PID} \equiv \log \frac{\mathcal{P}(\Delta E | \text{proton}, p)}{\mathcal{P}(\Delta E | \text{pion}, p)}$$



$p < 450 \text{ MeV}/c$, PIDcut=0:
pion contamination $\approx 0.1\%$
proton efficiency $> 99\%$





DVCS and the Recoil



- ▶ Missing ϕ : $\Delta\phi = \phi_{\text{meas}} - \phi_{\text{calc}}$
- ▶ Missing p : $\Delta p = p_{\text{meas}} - p_{\text{calc}}$

Missing Mass ($\approx M_p^2$):
 $M_X^2 = (p + p_{\gamma^*} - p_{\gamma})^2$




Separation of Resonant States in DVCS

DVCS / Bethe Heitler

- Elastic:
 - ▶ $ep \rightarrow ep\gamma$
- Resonant ('associated'):
 - ▶ $ep \rightarrow e\Delta^+\gamma$

$$\Delta^+ \rightarrow \begin{cases} n\pi^+, & 1/3 \\ p\pi^0, & 2/3 \end{cases}$$
 - ▶ 12% of signal
- Presence of $\pi^0 \Rightarrow$ proton fails coplanarity cut
 - ▶ Select elastic:
 - ★ $|\Delta\phi| < 0.1$ rad
 - ★ $|p_T^{\text{calc}}| / |p_T^{\text{meas}}| = 0.5 \div 1.5$
 - ▶ Select resonant:
 - ★ $|\Delta\phi| > 0.35$ rad




Separation of Resonant States in DVCS


DVCS / Bethe Heitler

- Elastic:
 - ▶ $ep \rightarrow ep\gamma$
- Resonant ('associated'):
 - ▶ $ep \rightarrow e\Delta^+\gamma$



$$\Delta^+ \rightarrow \begin{cases} n\pi^+, & 1/3 \\ p\pi^0, & 2/3 \end{cases}$$
 - ▶ 12% of signal
- Presence of $\pi^0 \Rightarrow$ proton fails coplanarity cut
 - ▶ Select elastic:
 - ★ $|\Delta\phi| < 0.1$ rad
 - ★ $|\rho_T^{\text{calc}} / \rho_T^{\text{meas}}| = 0.5 \div 1.5$
 - ▶ Select resonant:
 - ★ $|\Delta\phi| > 0.35$ rad



Exclusive Mesons and the Recoil



DVCS Beam Helicity Asymmetry with Recoil





Pre-preliminary,
private analysis

DVCS Beam Helicity Asymmetry with Recoil



**Pre-preliminary,
private analysis**

DVCS Beam Helicity Asymmetry with Recoil



Pre-preliminary,
private analysis

DVCS Beam Helicity Asymmetry with Recoil





Pre-preliminary,
private analysis

Elastic fraction: >95%



Summary

- * Generalized Parton Distributions
 - * Give glimpse of three-dimensional structure of nucleons
 - * Allow to access total angular momentum carried by quarks
- * Deeply Virtual Compton Scattering:
the golden channel to study GPDs
 - * Measurements of cross sections and asymmetries as input to GPD constraints and fits
 - * Measurement of various azimuthal asymmetries at HERMES and other fixed target experiments
- * HERMES high lumi run 2006/2007 with Recoil detector
 - * Exclusive event tagging
 - * First time: separation of elastic and resonant BSA in DVCS


Extraction of GPD $H(x, \xi=x, t)$ from DVCS data



Extraction of GPD $H(x, \xi=x, t)$ from DVCS data





Kinematics	H1/ZEUS	HERMES	JLab
x_B	$10^{-2} \dots 10^{-4}$	0.03...0.35	0.1...0.5
Q^2 [GeV ²]	1...100	1...10	1...5
W [GeV]	30...170	3...6	>2
$ t $ [GeV ²]	0.08...1	0.01...0.7	0.1...0.6



HERA: DVCS Cross Section Measurements

W and Q² dependence




- * Detected particles
 - * H1: electron and photon
 - * ZEUS: electron, photon subsample: proton tagged
- * BH contribution (MC) subtracted from 'DVCS-enriched' sample
- * Steep W^δ dependence
 - * $\delta \approx 0.7$, independent of Q^2
 - * DVCS is hard process
 - * high $W = \text{low } x_B$
 - * gluons resolved
- * Decrease of σ_{DVCS} with Q^2
 - * Q^{2n} dependence, $n \approx 1.5$
 - * slower than for exclusive vector mesons

HERA: Transverse Extension of Partons

DVCS cross section differential in t

- * Extract $d\sigma/dt$ in bins of Q^2 and W
- * Ansatz for quark GPDs: $\exp(-b|t|)$
- * t slope \Rightarrow average impact parameter
 - * Description of transverse extension of parton
 - * Size of proton core (w/o soft periphery)


$$\sqrt{\langle r_T^2 \rangle} = (0.65 \pm 0.02) \text{ fm} \quad \text{at } x_B = 10^{-3}$$



Hall A: DVCS Cross Section Measurements

- * E00-110 in Hall A at JLab
5.75 GeV e- beam on H₂ target
76% beam polarization
- * Detected: e-, γ , proton
- * Valence quark region


Cross section vs. azimuthal angle $\angle(\gamma^*, \gamma)$



Hall A: DVCS Cross Section Measurements

- * E00-110 in Hall A at JLab
5.75 GeV e- beam on H₂ target
76% beam polarization
- * Detected: e-, γ , proton
- * Valence quark region

Cross section vs. azimuthal angle $\angle(\gamma^*, \gamma)$



Helicity-dependent

Helicity-independent

$\text{Im}(\mathcal{I})$

$\sin\phi$ behavior

GPDs @ $x=\xi$

$\text{Re}(\mathcal{I})$


main contributions:
BH + twist-2 interference part
negligible: twist-3 and $|\text{DVCS}|^2$

integral of GPDs over x

Hall A: DVCS Cross Section Measurements

- * E00-110 in Hall A at JLab
5.75 GeV e- beam on H₂ target
76% beam polarization
- * Detected: e-, γ , proton
- * Valence quark region

Cross section vs. azimuthal angle $\angle(\gamma^*, \gamma)$



Helicity-dependent

Helicity-independent

$\text{Im}(\mathcal{I})$

$\sin\phi$ behavior

GPDs @ $x=\xi$

$\text{Re}(\mathcal{I})$

main contributions:
BH + twist-2 interference part
negligible: twist-3 and $|\text{DVCS}|^2$


integral of GPDs over x

Computed BH contribution
< helicity-independent cross section
Important to keep in mind for denominator of
beam helicity asymmetry = $d^4\Sigma/d^4\sigma$

Hall A: DVCS Cross Section Measurements

- * E00-110 in Hall A at JLab
5.75 GeV e- beam on H₂ target
76% beam polarization
- * Detected: e-, γ , proton
- * Valence quark region

Cross section vs. azimuthal angle $\angle(\gamma^*, \gamma)$



Helicity-dependent

Helicity-independent

$\mathcal{I}m(\mathcal{I})$

$\sin\phi$ behavior

GPDs @ $x=\xi$

$\mathcal{R}e(\mathcal{I})$

main contributions:
BH + twist-2 interference part
negligible: twist-3 and $|DVCS|^2$


integral of GPDs over x

No Q^2 dependence of $\mathcal{I}m(\mathcal{I})$ observed

- Indication of factorization
- GPDs accessible at moderate Q^2


Computed BH contribution
< helicity-independent cross section
Important to keep in mind for denominator of
beam helicity asymmetry = $d^4\Sigma/d^4\sigma$

HERA/H1: Beam Charge Asymmetry



- * First measurement at a collider (low $x_B=10^{-4} \dots 10^{-2}$)
 - * $6.5 < Q^2 < 80 \text{ GeV}^2$, $30 < W < 140 \text{ GeV}$, $|t| < 1 \text{ GeV}^2$
- * Positive $\cos\phi$ amplitude $0.16 \pm 0.04 \pm 0.06$
 - $\Rightarrow \text{Re}(\tau_{\text{DVCS}}) > 0$
- * Ratio $\rho = \text{Re}(\tau_{\text{DVCS}}) / \text{Im}(\tau_{\text{DVCS}})$
 - * $\rho = 0.20 \pm 0.05(\text{stat}) \pm 0.08(\text{sys})$
 - * In good agreement with calculation from dispersion relation

Nuclear DVCS at HERMES




Hydrogen, Krypton, Xenon:
2 beam charges



BCA

BSA sensitive to
squared DVCS term

Nuclear DVCS at HERMES








Nuclear DVCS at HERMES




BSA for Hydrogen, Helium, Nitrogen, Neon, Krypton, Xenon

C. Riedl (DESY), EIC Meeting at BNL, August 25, 2009

Corrections and systematic uncertainties


- (, ) Shift of exclusive peak between e^- and e^+ data (small)
- (, ) Semi-inclusive and exclusive background
⇒ Fractions from Monte Carlo
- () Acceptance, bin-width, smearing and detector misalignment (main contribution)
⇒ Estimated from Monte Carlo simulation employing range of available models
⇒ Model dependence
- The contributions from the resonance region, e.g.



stays part of the signal, in average 12%!

The underlying **“associated” asymmetry is unknown!**

Acceptance, Bin-width, Smearing and Misalignment



The difference between “model-generated” and in the HERMES acceptance reconstructed MC amplitudes is taken as systematic uncertainty

Some Literature

* Selected GPD review articles:

- * A. Radyushkin, Phys. Lett. B380, 417 (1996); Phys. Rev. D56, 5524 (1997)
- * A.V. Belitsky, D. Mueller and A. Kirchner, *Nucl. Phys. B* **629** (2002), p. 323
- * M. Diehl, Phys. Rept. 388, 41 (2003).

* Hermes DVCS data:

- * Transverse target spin asymmetry: HERMES Collaboration, A. Airapetian et al., JHEP 0806, 066 (2008).
- * Beam helicity and beam charge asymmetries on hydrogen, deuterium and nuclear targets:
3 publications in fall 2009
- * Longitudinal target spin asymmetries on hydrogen and deuterium to come