Event anisotropy measurement in 200GeV Au+Au collisions at RHIC-PHENIX experiment

ShinIchi Esumi,[a] for the PHENIX collaboration

[a]Inst. of Physics, Univ. of Tsukuba, Tenno-dai 1-1-1, Tsukuba, Ibaraki 305, Japan

Event anisotropy is expected to have sensitivity to the early stage of ultra-relativistic heavy-ion collisions at RHIC. The possible formation of a quark gluon plasma (QGP) could affect how the initial anisotropy in the space coordinate is transferred into the momentum space for the final state. The anisotropy parameter (v_2) is an amplitude of the 2nd harmonic parameter of the azimuthal distribution with respect to the reaction plane. We present here v_2 of identified (π^+, π^-, K^+, K^-, p and $\overline{p}$) and inclusive charged particles measured in the PHENIX central arm detector ($|\eta| < 0.35$) with respect to the reaction plane defined at $|\eta| = 3 \sim 4$ in 200 GeV Au+Au collisions.

1. Introduction and Experimental Setup

The PHENIX experiment at RHIC is designed to measure a variety of observables in ultra-relativistic heavy-ion collisions. The primary goal is to detect signals of the QGP. The v_2 measurements [1,2] at RHIC energies have been performed in several experiments for inclusive charged particles and for identified particles. Extending the v_2 measurements at higher p_T region, especially with identified particles, is important to understand the origin of the v_2 component: hydro-dynamic flow of compressed matter, the production of many mini-jets, etc. It has also been observed that v_2 saturates at $p_T \sim 2$ GeV/c. The cause of this is not yet known, but it is worth noting that at this momentum the particle composition is very different than at low momentum in that the proton yield is comparable to the pion yield [3].

PHENIX has excellent particle identification at mid-rapidity. In this analysis, the time-of-flight (TOF) detector is used to identify final state hadrons up to 4 GeV/c in p_T together with the central arm tracking system consisting of drift- and pad-chambers for momentum reconstruction. The beam counters (BBC) provide collision time and z-vertex position information. The two beam counters are located at $|z|$=1.5m from the collision point, surrounding the beam pipe with 64 photo-multiplier tubes, covering $|\eta| = 3 \sim 4$. The large charged multiplicity at $|\eta| = 3 \sim 4$ and non-zero signal of event anisotropy in this η range enables us to measure the event reaction plane using the BBC with full azimuthal angle coverage. The standard method [4,5] for calculating and correcting (flattening) the reaction plane is applied here. Figure 1 shows the average cosine of the difference between two reaction planes defined by two beam counters. The left panel shows a negative correlation, which indicates a back-to-back type directed (1st) moment between the two reaction planes in the forward and backward rapidities. The right panel shows



Figure 1. Correlation of reaction planes between two beam counters for the 1st (left) and the 2nd (right) moment, respectively. The solid symbol is for the average cosine which corresponds to the reaction plane resolution, the open symbol is for the average sine, which has to be zero because of the left-right symmetry.

the elliptic (2nd) moment, which has a positive correlation with a larger magnitude than that for the directed moment. Note that the reaction plane from the elliptic moment has the same orientation for forward and backward rapidities. Also note that the elliptic moment resolution is better than that for the directed moment. In order to obtain an improved reaction plane resolution, a combined reaction plane is defined by averaging the reaction plane angles obtained from each BBC using the elliptic moment. The estimated resolution of the combined reaction plane, $< cos\ 2(\Phi_{measured} - \Phi_{truth}) >$, is an average of 0.3 with a maximum of about 0.4.

2. Results and Interpretation

Charged particles are measured in the central arm spectrometers ($|\eta| < 0.35$). In the present analysis, the p_T range of the particle identification is $0.2 < p_T < 3$ GeV/c for pions, $0.3 < p_T < 2$ GeV/c for kaons, and $0.5 < p_T < 4$ GeV/c for protons. The centrality of the collision is defined by the correlation between the total number of particles measured in the BBC and the total energy measured in zero degree calorimeter.

Figure 2 shows the centrality dependence of v_2 for inclusive charged particles measured at mid-rapidity ($|\eta| < 0.35$) with respect to three different reaction planes: two determined using each beam counter separately, and one determined from the combination of the two beam counters. Here R.P. stands for the reaction plane within the η range shown in the figure. The three results agree well, indicating that the resolution for the different reaction planes is well understood. The v_2 parameter decreases for both peripheral and central collisions with a maximum at about 50% of the geometric cross section. The reduction of v_2 for peripheral events might indicate that non-flow contributions are less significant in this analysis because of the large rapidity gap between the reaction plane and the central tracking arm acceptance of about 3 units. The data are consistent with a $< sin\ 2(\phi - \Phi) >$



Figure 2. Centrality dependence of v_2 for inclusive charged particle measured at mid-rapidity ($|\eta| < 0.35$) and p_T integrated between 0.2 and 10 GeV/c with respect to the reaction plane defined by the two beam counters separately or combined. Here R.P. stands for the reaction plane within the specified η range.



Figure 3. Transverse momentum dependence of v_2 for inclusive charged particles at mid-rapidity ($|\eta| < 0.35$) with respect to the combined reaction plane ($|\eta| = 3 \sim 4$) for minimum-bias collisions. Included are comparisons to the model calculations, where the details are described in the text below and in refs [6,7].

distribution (represented by the open symbols), which is expected to be zero. Figure 3 shows the transverse momentum dependence of v_2 for inclusive charged particles with respect to the combined reaction plane for minimum-bias events. The data above a p_T of 2 GeV/c clearly show a deviation from the linear behavior seen at smaller p_T. The solid line is from a hydro-dynamic calculation [6]. The three dashed lines are from a hydro + pQCD calculation with three different gluon densities (from top to bottom, dN^g/dy = 1000, 500, and 200) [7].

Figure 4, the transverse momentum dependence of v_2 for identified particles is shown. The solid circles show p and $\overline{p}$, the open triangles show K^+ and K^-, and the open squares show π^+ and π^-. The left panel shows negatively charged particles, while the right panel shows positively charged particles. The solid lines represent hydro-dynamic calculation [6] including a first-order phase transition with a freeze out temperature of 120 MeV. The data show that at lower p_T (< 2 GeV/c) the lighter mass particles have a larger v_2 at a given p_T, which is reproduced by the model calculations. However, the data and the calculation deviate at higher p_T. Lighter particles seem to deviate from the hydro-dynamic behavior at lower p_T than for the heavier particles. Consequently, the mass dependence of v_2 for p_T above 2 GeV/c is reversed.



Figure 4. Transverse momentum dependence of v_2 for identified particles, π^-, K^-, $\overline{p}$ (left) and π^+, K^+, p (right). The solid lines represent a hydro-dynamic calculation [6] including a first-order phase transition with a freeze out temperature of 120 MeV for π, K and p from upper to lower curves, respectively.

3. Summary

The value of v_2 for identified and inclusive charged particles are measured at mid-rapidity with respect to the reaction plane defined in the forward and backward rapidity regions in 200 GeV Au+Au collisions in the PHENIX experiment at RHIC. It was found that the v_2 value of inclusive charged particle increases from central to mid-central collisions reaching a maximum at about 50% of the geometric cross section and then decreases again for more peripheral collisions. As a function of transverse momentum in minimum-bias collisions, the v_2 parameter increases linearly with p_T up to $p_T \simeq 2$ GeV/c and then saturates for inclusive charged particles. The v_2 parameter of identified particles (π^+, π^-, K^+, K^-, p and $\overline{p}$) follow a hydro-dynamic behavior up to 2 GeV/c in p_T, where the lighter mass particles have larger v_2 at a given p_T. However there is an indication that this trend is reversed at around $p_T \simeq 2$ GeV/c, where p and $\overline{p}$ have larger v_2 than π and K.

REFERENCES

1. K.H. Ackermann et al., Phys. Rev. Lett. **86**, 402, (2001), Phys. Rev. Lett. **87**, 182301, (2001), J.Phys. **G28**, 2089, (2002).
2. K. Adcox et al., nucl-ex/0204005.
3. K. Adcox et al., Phys. Rev. Lett. **88**, 242301, (2002).
4. J.Barrette et al., Phys. Rev. Lett. **73**, 2532, (1994), Phys. Rev. **C55**, 1420, (1997), Phys. Rev. **C56**, 3254, (1997),
5. A.Poskanzer and S.Voloshin, Phys. Rev. **C58**, 1671, (1998)
6. P.Huovinen, P.F.Kolb, U.W.Heinz, P.V.Ruuskanen and S.A.Voloshin, Phys. Lett. B503, 58 (2001)
7. M.Gyulassy, I.Vitev and X.N.Wang, Phys. Rev. Lett. 86, 2537, (2001)