Particle Correlation Measurements from

Stephen C. Johnson – Lawrence Livermore National Lab
for the PHENIX Collaboration

HBT results presented here:
submitted to PRL: 16 Jan `02
nucl-ex/0201008

The Physics of Hanbury-Brown Twiss



- Consider a source S(r) of identical bosons (γ or π) whose wave functions can be described as plane waves.
- Assume:
  - Production amplitudes independent of momentum
  - Mutually incoherent

Amplitude for this diagram:

$$A = \frac{1}{\sqrt{2}}\left[e^{ip_1(r_1-x)}e^{ip_2(r_2-y)} + e^{ip_1(r_1-y)}e^{ip_2(r_2-x)}\right]$$

$$A^2 = 1 + e^{i(p_1-p_2)(y-x)}$$

$$\approx 1 + \cos(\Delta p \Delta x)$$

Corresponding normalized probability:

$$P = \int dxdy|A|^2 S(x)S(y) = 1 + \left|\tilde{S}(q)\right|^2 \equiv C_2(q)$$





Some details:

In principle:

$$C_2 = \frac{\langle n \rangle^2}{\langle n(n-1) \rangle} \frac{dn/dp_1 dp_2}{\left(dn/dp_1\right)\left(dn/dp_2\right)}$$

In practice:

$$C_2 = \frac{A(q)}{B(q)}$$

Pairs from same event

Pairs from 'mixed' events



$$E\frac{dN}{d^3p} = \int d^4x S(x,p)$$

$$q = p_1 - p_2$$

$$k = p_1 + p_2/2$$

$$C_2(q,k) = 1 + \frac{\left|\int d^4x S(x,k) e^{iq\cdot x}\right|^2}{\int d^4x S\left(x, k + \frac{1}{2}q\right) \int d^4y S\left(y, k - \frac{1}{2}q\right)}$$

- One dimension works fine when measuring stars:
  - Static
  - Isotropic emission (no position momentum correlations)
- Heavy Ion Collisions are anything but
  - Consider a more complicated source
- S() – effective single particle Wigner phase space density
  - Often replaced by a classical phase-space density in practical calculations
- Note: due to mass shell constraints, the function is non-invertible (model assumptions).

Kinematic Variables

- Currently the field tends to consider projections of the momentum difference into:
  - q_{long}, q_{side}, q_{out}
- q_{long} ↔ R_{long} ~longitudinal extent
- q_{side}/q_{out} ↔ R_{side}/R_{out} ~transverse extent



$$q = p_1 - p_2$$
$$k = p_1 + p_2/2$$

$$C_2(q,k) = 1 + \frac{\left|\int d^4x S(x,k) e^{iq\cdot x}\right|^2}{\int d^4x S\left(x, k + \tfrac{1}{2}q\right) \int d^4y S\left(y, k - \tfrac{1}{2}q\right)}$$

HBT and the QGP

- "Naïve" picture:
  - $R_{out}^2 = R_{side}^2 + (\beta_{pair}\tau)^2$



- Concrete predictions are few:
  - Pratt PRD 1314 (`86): fireball and EOS Ł τ ~ 90 fm/c
  - Bertsch NPA 173 (89) QGP + cascade Ł τ ~ 12 fm/c
  - Hydro calculation of Rischke & Gyulassy expects R_{out}/R_{side} ~ 2->4 @ k_t = 350 MeV.
  - Result robust to T_{freeze}, d_Q/d_H, 1st order vs. rapid cross-over.
- Response: can hadronic rescattering mask this prediction?

D.H. Rischke, M. Gyulassy/Nuclear Physics A 608 (1996) 479–512



Fig. 17. The same as in Fig. 12, but for the ratio R_{out}/R_{side}.



One step closer



FIG. 3. R_{out}/R_{side} for RHIC initial conditions, as a function of K_T at freeze-out (symbols) and at hadronization (lines).

- Soff, Bass, Dumitru (PRL 86)
  - Couple microscopic transport to hydro with phase transition
  - Still expect $R_{out}/R_{side}>1$ ☹ measurements at high k_t are very interesting.
- Note:
  - Hydro: $R_o/R_s(200)<R_o/R_s(160)$
    - Longer time at phase transition
  - Transport: $R_o/R_s(200)>R_o/R_s(160)$
    - Longer time rescattering

PHENIX – Year 1 Configuration

- Both arms provide hadron PID (contrary to popular belief)
- East:
  - DC + TOF (~100 ps)
  - π/K separation to 2 GeV/c
- West:
  - DC + PbSc (~600 ps)
  - π/K separation to 1 GeV/c
- B-field + geometry limits lower k_t bound to 200 MeV.

$$\frac{\partial p}{p} = 0.6\% \oplus 3.6\% \, p$$

I use 'electron' acceptance for this hadron



Centrality definition and sample



- This data sample uses the 30% most central collisions
  - $<cent_{pairs}>$ = 10%
  - 493k events
- After all analysis cuts:
  - 3.1M π^+ pairs
  - 3.3M π^- pairs



Pair acceptance and corrections



- π definition ℄ <1.5σ from π peak && >2.5σ from K peak
- Require pairs from mixed events to have reconstructed vertices within 1cm
  - Acceptance varies as a function of vertex position
- Remove pairs within 2cm in drift chamber
  - Ghosting
- Remove tracks with EMC clusters within 12cm of each other in both real and mixed sample
  - Shower + tower size in EMC
- Correct for two track inefficiencies at low relative φ in the drift chamber.
- Full Coulomb Correction modified for momentum smearing
  - Partial correction changes radii results marginally
- Residual correlations in event mixed background ℄ <2% error
- Momentum smearing correction to correlation function

Systematic Errors:

R_{long} & R_{side}: 8%

R_{out}: 4.5%

Results:

Assume T = 125 MeV, β_f = .69/η_f=.85
From fits to singles spectra in centrality region 5-15%
(J. Burward-Hoy)

- Theoretical hydro-inspired fits.
  - $R_{side}^2(m_T) = \frac{R_{geom}^2}{1+\beta_f^2\left(\frac{m_T}{T}\right)} \Rightarrow R_{geom} = 6.7 \pm 0.2\, fm$

$m_t^2 = k_t^2 + m_\pi^2$

  - Chapman, Nix, Heinz PRC 52, 2694
  - $R_{side}^2(m_T) = \frac{R_{geom}^2}{1+\eta_f^2\left(\frac{1}{2}+\frac{m_T}{T}\right)} \Rightarrow R_{geom} = 8.1 \pm 0.3\, fm$
  - Wiedemann, Scotto, Heinz PRC 53, 918
- However, hydro calculations predict R_o than data, R_s, smaller.
- Much larger than comparable 1D RMS Au radius of 3.07fm
- k_t dependence suggest larger β_f/T (η_f/T) than fits to singles



Systematic Errors:
R_{long} & R_{side}: 8%
R_{out}: 4.5%

Compared to STAR



- Well described by hydro model if consider datasets separately.
  - But they indicate a much higher flow/temp ratio when taken together
  - Need to be careful about systematics between different measures.
  - Both experiments should be able to sort this out in the next dataset.
- Inconsistent with models of QGP that include an hadronic rescattering phase.

Energy dependence …

M. Lisa *et al.*, PRL **84**, 2798 (2000)
R. Soltz *et al.*, to be sub PRC
C. Adler *et al.*, PRL **87**, 082301
I.G. Bearden *et al.*, EJP **C18**, 317 (2000)

- R_{out} and R_{side} are energy independent within error bars.
- Smooth energy dependence in R_{long}
- No immediate indication of very different physics
- Fit R_{long} to: $\frac{A}{\sqrt{m_T}}$
- AGS: A = 2.19 +/- .05
- SPS: A = 2.90 +/- .10
- RHIC: A = 3.32 +/- .03



$A = \tau_0 T$ in 1st order T/m_T calculation

τ_0 = average freeze-out proper time

Even an experimentalist can't predict this

- "A prognostication"
  - Local PHENIX HBT expert keeps his predictions on the web.
- Zajc at Nucleus-Nucleus 97 showed this slide
- Extrapolation to RHIC multiplicities?
  - $(dN/dy)^{1/3}$ => R ~ 9 fm
  - (dN/dy) => R ~20 fm
- Neither.
- What is happening here?
- **



** To be fair: Bill's radii predictions are way off. His guess at the charged particle multiplicity in the extrapolations…bang on

Q & A

- Q: Why are Rout and Rside ~identical over an order of magnitude of beam energy?
  - There is ample evidence at AGS and SPS of dependence of radii on # target+projectile nucleons
    - Even though the larger nuclei have larger flow the radii follow a simple scaling
    - Why no multiplicity/energy dependence?



  - Q: Flow higher at RHIC leads to smaller radii?
    - HBT results suggest high flow but spectra imply flow comparable to SPS **
    - k_t dependence of radii is similar for different energies (competing mechanisms that create similar β_f/T?)
  - Q: Higher opacity at RHIC energies?
  - Why would opacity be higher at RHIC than at AGS?
    - $\pi n \rightarrow \Delta \rightarrow \pi n$ Ł $\pi\pi \rightarrow \rho \rightarrow \pi\pi$
    - And why would size and opacity effects identically cancel?
    - Red herring? – You can't create opaque source with smaller lifetime.
  - Q: Why is it only changing in the longitudinal direction?
    - Some surprise among theorists that the difference is so small.
    - Is there a quantitative expectation of the R_{long} dependence?

** To my knowledge no one has fit both SPS and RHIC results to same hydro model

A taste of the future

- For year-1 we wrote ~5 million min. bias events.
  - 1/2 million events in this analysis after all cuts
- In the past run we wrote ~90 million min. bias events (+14 million rare event triggers)
- Therefore, for the year-2 pion correlations:
  - easily 10 high statistics bins in k_t from .2 to 1.0 GeV + a few bins from 1.0 to 2.0 GeV
    - Capability to exclude detailed theories
    - Systematic errors start to become the real problem
- Beyond pions:
  - 1D proton and kaon correlations
  - Non-identical correlations (πK, πp, etc.)
  - π^0
    - maybe possible in year 2 … year 3?
    - Probe of very high k_t.
  - anti-neutron correlations
    - Better than proton correlations Ł lack of coulomb.



The scientists that make me look good*

PHENIX



University of São Paulo, São Paulo, Brazil
Academia Sinica, Taipei 11529, China
China Institute of Atomic Energy (CIAE), Beijing, P. R. China
Laboratoire de Physique Corpusculaire (LPC), Universite de Clermont-Ferrand, 63170 Aubiere, Clermont-Ferrand, France
Dapnia, CEA Saclay, Bat. 703, F-91191, Gif-sur-Yvette, France
IPN-Orsay, Universite Paris Sud, CNRS-IN2P3, BP1, F-91406, Orsay, France
LPNHE-Palaiseau, Ecole Polytechnique, CNRS-IN2P3, Route de Saclay, F-91128, Palaiseau, France
SUBATECH, Ecole des Mines at Nantes, F-44307 Nantes, France
University of Muenster, Muenster, Germany
Banaras Hindu University,Banaras, India
Bhabha Atomic Research Centre (BARC),Bombay, India
Weizmann Institute, Rehovot, Israel
Center for Nuclear Study (CNS-Tokyo), University of Tokyo, Tanashi, Tokyo 188, Japan
Hiroshima University, Higashi-Hiroshima 739, Japan
KEK, Institute for High Energy Physics,Tsukuba, Japan
Kyoto University, Kyoto, Japan
Nagasaki Institute of Applied Science, Nagasaki-shi, Nagasaki, Japan
RIKEN, Institute for Physical and Chemical Research, Hirosawa, Wako, Japan
University of Tokyo, Bunkyo-ku, Tokyo 113, Japan
Tokyo Institute of Technology, Ohokayama, Meguro, Tokyo, Japan
University of Tsukuba, Tsukuba, Japan
Waseda University, Tokyo, Japan
Cyclotron Application Laboratory, KAERI, Seoul, South Korea
Kangnung National University, Kangnung 210-702, South Korea
Korea University, Seoul, 136-701, Korea

Myong Ji University, Yongin City 449-728, Korea
System Electronics Laboratory, Seoul National University, Seoul, South Korea
Yonsei University, Seoul 120-749, KOREA
Institute of High Energy Physics (IHEP-Protvino or Serpukhov), Protovino, Russia
Joint Institute for Nuclear Research (JINR-Dubna), Dubna, Russia
Kurchatov Institute, Moscow, Russia
PNPI: St. Petersburg Nuclear Physics Institute, Gatchina, Leningrad, Russia
Lund University, Lund, Sweden
Abilene Christian University, Abilene, Texas, USA
Brookhaven National Laboratory (BNL), Upton, NY 11973
University of California - Riverside (UCR), Riverside, CA 92521, USA
Columbia University, Nevis Laboratories, Irvington, NY 10533, USA
Florida State University (FSU), Tallahassee, FL 32306, USA
Georgia State University (GSU), Atlanta, GA, 30303, USA
Iowa State University (ISU) and Ames Laboratory, Ames, IA 50011, USA
LANL: Los Alamos National Laboratory, Los Alamos, NM 87545, USA
LLNL: Lawrence Livermore National Laboratory, Livermore, CA 94550, USA
University of New Mexico, Albuquerque, New Mexico, USA
New Mexico State University, Las Cruces, New Mexico, USA
Department of Chemistry, State University of New York at Stony Brook (USB), Stony Brook, NY 11794, USA
Department of Physics and Astronomy, State University of New York at Stony Brook (USB), Stony Brook, NY 11794-, USA
Oak Ridge National Laboratory (ORNL), Oak Ridge, TN 37831, USA
University of Tennessee (UT), Knoxville, TN 37996, USA
Vanderbilt University, Nashville, TN 37235, USA

*or at least better than I normally do

Supporting slides:

Fitting the Single Particle Spectra:

Jane Burward-Hoy (LLNL)

$$1/m_t \, dN/dm_t = A \int f(\xi) \, \xi \, d\xi \, m_T \; K_1(m_T/T_{fo} \cosh \rho) \, I_0(p_T/T_{fo} \sinh \rho)$$







Switching frames

- Problem: radii extracted depend on the frame in which you measure
- Historically we have used:
  - Q_{inv} (1D pair rest frame)
  - Collision Center of Mass
    - AGS
  - LCMS (longitudinally co-moving frame)
    - Longitudinally boost invariant sources
    - SPS(?), RHIC
  - Jet frame
    - p-p collisions.
  - PCMS (pair center-of-mass – 3D version of Q_{inv})
  - …
- What is the 'correct' frame?? (= the frame in which the source is not moving)
- Can we bound it experimentally?

Our source??

2π HBT measurement is simultaneous in the PCMS frame

$$R_{OUT}^2 = \gamma_S^2 \{(1 - \beta_P v_S)^2 R^2 + (\beta_P - v_S)^2 \tau^2\}$$



Source



Detector

Pair rest frame != source rest frame
What is the β_{source}?
We don't know, but we can bound it:
>0
<β_{pair}

LCMS vs. PCMS





- LCMS
  - A Lorentz contracted measurement of the frame
- PCMS
  - A measurement of the length of the train in the train frame

unphysical

True frame

unphysical

Fits to entire dataset (π^-)



LCMS

PCMS*

*PCMS = Pair Center of Mass System

PCMS results

- R_{side} and R_{long} unchanged
  - And not plotted
- R_{out}^{PCMS} differs by $\langle\gamma_{pair}\rangle$ from R_{out}^{LCMS}
- Be careful about deducing a lifetime from $R_{out}^2=R_{side}^2+(\beta_{pair}\tau)^2$
  - $\beta_{pair}=0$