



# Reconstruction of $\phi$ Mesons in $K^+K^-$ Channel at $\sqrt{s}=200$ GeV by the PHENIX Experiment at RHIC

**Mohammed Muniruzzaman**

University of California Riverside



# Motivation

- *Sensitive to strangeness enhancement*
- *Lifetime of  $\phi$  ( $\tau_\phi \sim 44\text{fm}/c$ ) makes it a wonderful probe of the deconfined phase*



- ▶ An **enhanced  $\phi$ -meson production** has been suggested as a signature for the formation of a deconfined phase
- ▶ **Medium modifications** of  $\phi$ -meson properties might be related to the expected chiral phase transition
- ▶ The **change in the branching ratio** in leptonic and hadronic channels may point to the chiral symmetry restoration.

**R. Rapp nucl-th/0204003**

# PHENIX in Run-11



- PHENIX Central Arm
  - Capable of Measuring hadrons, electrons and photons

## Tracking:

Drift Chamber

Pad Chamber1


Pad Chamber3

## Kaon Identification

Electromagnetic Calorimeter


Time-of-Flight Detector

*Excellent Particle Identification Capability*



# Run Condition

## RHIC Beam Delivered at PHENIX



- Au-Au at  $E_{cm} = 200$  GeV
  - At  $42 \mu\text{b}^{-1}$
  - Vertex between  $\pm 45$  cm
  - Over 50% of Data accumulated in the last two weeks of Au-Au Run
- p-p at  $E_{cm} = 200$  GeV at  $700 \text{ nb}^{-1}$

**92 million min-bias data recorded for Au-Au Run**


**After Imposing Strict Data Quality Control, 27 Million min-bias data were selected for analysis**

# PHENIX Event Display



*So many  
Tracks !!*


**Need good  
Particle ID**




$\phi \rightarrow K^+K^-$   
analysis is  
possible if we  
can point out  
which tracks  
are kaons

# Particle Identification

Particle Identified through  
High Resolution Time-of-  
Flight Detector



Particle Identified through  
Electromagnetic Calorimeter




# Data Sample Selection

- ◆ 27 Million Minimum Bias Au+Au events
- ◆ Track Selection
  - 3  $\sigma$  track projection matching at PC3
  - 3  $\sigma$  matching on PC3
  - Best quality DC tracks
- ◆ Energy Loss in TOF > 2 MeV
- ◆ Momentum between 200 MeV/c and 1.2 GeV/c in Time of Flight
- ◆ Momentum upto 900 MeV/c in EMCal
- ◆ 3 $\sigma$  cut on the calibrated  $mass^2$  of kaon
- ◆ 5 $\sigma$  above the Pion  $mass^2$  band

Mass squared of Particles are Calculated. Sample is selected on its  $mass^2$  Distribution



3 $\sigma$  around the peak of  $mass^2$



# $\phi$ Invariant Mass

Fit with a Breit-Wigner  
Convolved with a Gaussian

Actual pairs      Mixed background



Mixed  $N_{+-}$  Normalized by  
Actual Pair  $N_{++}$  and  $N_{--}$  as  
 $2\sqrt{(N_{++} \cdot N_{--})}$




# $\phi$ Invariant 2k


Another approach to extract  $\phi$  signal by looking at the difference of 4-momenta of kaons:  $k = \sqrt{(p_{1\mu} - p_{2\mu})^2}$

## Advantages:

- Pushes the  $\phi$  peak away from the kinematic edge
- Puts any correlated pairs into a different region of phase space



Mixed  $N_{+-}$  Normalized by Actual Pair  $N_{++}$  and  $N_{--}$  as  $2\sqrt{(N_{++} \cdot N_{--})}$



# Background above $\phi$

- Investigate background above  $\phi$ ,  $m \sim 1.07$  GeV


- ◆ Make looser PID cut ( $4\sigma$ )
- ◆ Subtraction done by normalizing to region above peak
- ◆ Larger contamination from secondary peak

- $K_S$  from misidentified momentum in the TOF array?

- ◆ Reconstruct mass assuming particles are pions
- ◆ Peak shows up at  $\sim 494$  MeV  $\sim M_K$


- Make a tighter PID cut  $1\sigma$

- ◆ Extra mass peak largely disappears




# Acceptance vs $m_t$

- 3 Centrality Classes  
0%-10%, 10%-40%, 40%-92%
- 8  $M_t$  Bins 1.6-1.8, 1.8-2.0, 2.0-2.2, 2.2-2.4, 2.4-2.6, 2.6-2.8, 2.8-3.0, 3.0-4.0
- Fitting for each centrality and  $M_t$  bins are done by fixing the values of mass and  $\Gamma$  from the fit of min bias, and letting the constant free.



Work in progress toward  $m_t$  distribution





# $\phi$ Invariant Mass in EMCal

## Advantages:

- Large Acceptance
- Kaons accepted upto  $p < 900$  Mev
- Complements the TOF measurements in the low  $m_t$  region

Actual pairs

Mixed background



# Summary

- Using the excellent PID capability of PHENIX Time of Flight Detector  $\phi$  is reconstructed via  $K^+K^-$  channel for the Au-Au Collisions at  $\sqrt{s_{NN}}=200\text{GeV}$ .
- The mass is fitted with a Breit-Wigner Convolved with a Gaussian with a resolution of 1.2 MeV determined from simulation. The preliminary fit values for the mass and  $\Gamma$  agrees well the PDB values. The systematic errors on these numbers are still under investigation.
- With the available statistics 3 bins of centrality and 8 bins in  $m_t$  could be made which enabled us to study the yield of  $\phi$ . This analysis is still in progress.
- It has been demonstrated that  $\phi$  can be reconstructed through Electromagnetic Calorimeter. This will enable us to accumulate more statistics and probe into the low  $m_t$  region.