

Colloquium, GSI, 13 November 2007

Raphaël Granier de Cassagnac  
Laboratoire Leprince-Ringuet  
PHENIX experiment

# **CHARMING CHARMONIA IN THE QUARK GLUON PLASMA**

# THE PROGRAM

---

## ✘ The quark gluon plasma

- + What is it ? Where to find it ? When ?
- + How to recognize it ?

## ✘ Charming charmonia

- + How charm quarks and “charmonia” mysteriously appeared, disappeared, reappeared, by magic...

## ✘ Other charms

- + Two golden signatures of the QGP and how charm quarks surprisingly behave...


What ? Where ? When ? How ?

# THE QUARK GLUON PLASMA

---

# QUARKS, MESONS, BARYONS

- ✗ Protons and neutrons are made of quarks tightly bound together by gluons carriers of the strong interaction
- ✗ 6 quarks discovered
  - + (up, charm, top) charge +2/3
  - + (down, strange, beauty) -1/3
- ✗ Allowed "hadrons" :
  - + Baryons = 3 quarks
    - ✗ p (uud), n (udd), etc.
  - + Mesons = quark + antiquark
    - ✗ pion ( $u\bar{d}$ ), etc.




(gluons not depicted)

# THE STRONG INTERACTION...

- ✘ as seen from (HERA...) data and described by quantum chromodynamics (QCD)
- ✘ ... is strong at low energy, i.e. short distance ( $\approx 1\text{fm}$ )


- No free quark
- Bound hadrons
- Confinement



- ✘ ... but weak at high energy → “asymptotic freedom”

# A STRONG PREDICTION


- ✘ (Lattice) QCD predicts a phase transition from nuclear / hadronic matter to a Quark Gluon Plasma (QGP)
- ✘ Critical parameters:
  - +  $T_c \approx 160 \text{ MeV}$  ( $2 \times 10^{12} \text{ K}$ )  
 $\approx 20\,000 \times T_{\text{sun}}$
  - +  $\epsilon_c \approx 1 \text{ GeV/fm}^3$   
 $\approx 6 \times \text{nuclear density}$



Karsch, hep-lat/0106019  
 Lect. Notes Phys.583 (2002) 209

# A (VERY) NAIVE PICTURE OF THE QGP

- ✗ A bit like melting ice...
- ✗ Above  $T_c$






Sorry,  
powerpoint  
animated...

(gluons and secondaries not depicted)

# WHERE/WHEN CAN WE FIND THE QGP?

1. Early in the universe ( $t < 10\mu\text{s}$ )
  - + But little chance to leave relics
2. Core of a compact star
  - + No smoking gun candidate so far
3. In the lab, colliding heavy ions
  - + Freedom for the quarks...
  - + ... for some  $10^{-23}$  s






$\sqrt{s_{nn}}$   
↓


# HEAVY ION COLLISIONS

F  
I  
X  
E  
D  
  
T  
A  
R  
G  
E  
T  
  
  
  
  
  
  
  
  
  
C  
O  
L  
L  
I  
S  
I  
O  
N  
S

1986 BNL - AGS 4 GeV



1994 CERN - SPS 20 GeV



2000 BNL - RHIC 200 GeV



2008? CERN - LHC 5 TeV



F  
I  
X  
E  
D  
  
T  
A  
R  
G  
E  
T  
  
  
  
  
  
  
  
  
  
C  
O  
L  
L  
I  
S  
I  
O  
N  
S

## EPISODE 2, THE PAST

- ✘ SPS: Super Proton Synchrotron
- ✘ @ CERN (Geneva)
- ✘ Fixed target experiments
  - + (WA98, CERES, NA44, NA49, NA57, NA50, NA60...)
- ✘ Various collisions p+p, p+A, A+A up to Pb+Pb
- ✘  $\sqrt{s_{NN}} \approx 20 \text{ GeV}$




→ Estimated energy density  $\varepsilon \approx 3 \text{ GeV/fm}^3 > \varepsilon_c$

→ In principle, QGP have been produced...

## EPISODE 3, THE PRESENT...

- ✘ RHIC: Relativistic Heavy Ion Collider  
@ Brookhaven National Lab. (New-York)
  - + First collisions in 2000
  - + 2 large (STAR & PHENIX)
  - + 2 smaller (PHOBOS & BRAHMS) experiments
- ✘ Can collide anything from p+p (up to 500 GeV)  
to Au+Au (up to 200 GeV per nucleon pairs)



# HOW TO FIND THE QGP?

- ✗ Theorists predict a QGP signature
- ✗ Experimentalists look at it versus A+A collision growing centrality →
- ✗ Then compare to a reference process
  - + Unaffected by QGP
- ✗ Then compare to p+A
  - + No QGP expected
  - + But “normal” nuclear matter could mimic QGP signatures...

↑ centrality

↓ impact parameter


↓ spectator nucleons

↑ number of participants  $N_{\text{part}}$

↑ number of elementary (n+n) collisions  $N_{\text{coll}}$

↑ particle multiplicity

↑ energy density



# WHICH SIGNATURES?

QGP hints seen at RHIC:

- ✘ Total multiplicity
- ✘ High  $p_T$  suppression
- ✘ Back to back jets
- ✘ Elliptic flow
- ✘ Baryon/meson ratio
- ✘  $J/\psi$  suppression
- ✘ Thermal radiation
- ✘ ...

But others ones were also predicted!

*“There was a general feeling that if the quark-gluon plasma was indeed produced, it would manifest itself in a variety of unknown but dramatic ways, including...”*

H. Satz @ Lattice 2000

hep-ph/0009099

# WHICH SIGNATURES?

July 18 1999

BRITAI



*Ready for blastoff: a Brookhaven engineer puts finishing touches to the ion collider*

## Big Bang machine could destroy Earth

by [Jonathan Leake](#)  
*Science Editor*

A NUCLEAR accelerator designed to replicate the Big Bang is under investigation by international physicists because of fears that it might cause "perturbations of the universe" that could destroy the Earth. One theory even suggests that it could create a black hole.

But others ones were also predicted!

*"There was a general feeling that if the quark-gluon plasma was indeed produced, it would manifest itself in a variety of unknown but dramatic ways, including...*

*...the end of the world"*

H. Satz @ Lattice 2000

hep-ph/0009099

One of the quark gluon plasma signature

# CHARMING CHARMONIA

---

# THE CHARM QUARK

- ✗ 4<sup>th</sup> quark to have been discovered (1974)
- + Was expected from weak interaction
  - ✗ “It appeared like a charm” around 1964
  - ✗ A partner to the strange quark
  - ✗ GIM mechanism (1970)
  - ✗ Strong theoretical need for a fourth quark

Glashow, Iliopoulos, Maiani,  
PRD2 (1970) 1285

Lepton	Neutrino
Electron	$\nu_e$
Muon	$\nu_\mu$

Quark +2/3	Quark -1/3
Up	Down
???	Strange


\* Charm is summoned by theoretical magic



# 1973: NON DISCOVERY

Christenson et al, PRD8 (1973) 2016

- ✘  $p+U \rightarrow \mu^+\mu^-$  (@ 30 AGeV)
- ✘ An excess above continuum
  - + around  $3 \text{ GeV}/c^2$
- ✘ No conclusion
  - + Poor resolution ( $\sim 500 \text{ MeV}/c^2$ )
- ✘ We now know this was
  - + Bound state of  $(c\bar{c}) \rightarrow \mu^+\mu^-$



\* Charm appears without being noticed...



Augustin et al, PRL33 (1974) 1404

Aubert et al, PRL33 (1974) 1406

Ting and Richter, Nobel prize 1976

# 1974: J & $\Psi$ DISCOVERY

- ✘ With much better resolutions, two teams discover at the same time:
  - + “Experimental Observation of a Heavy Particle J”
  - + “Discovery of a Narrow Resonance in  $e^+e^-$  Annihilations”  $\rightarrow \Psi$
- ✘ First particle containing charm
  - +  $(c\bar{c})$  most stable bound state  $J/\psi$
  - + Other “charmonia” soon after:  $\Psi'$ ,  $\chi_c$ ...



✧ Charm is seen simultaneously in two different places...

# THE CHARM QUARK

✗ 4<sup>th</sup> quark to have been discovered (1974)

+ Was expected from weak interaction

\* *It appeared like a (magical) charm*

+ But interesting strong properties

✗ Heavy ( $1.5 \text{ GeV}/c^2$ ) more than a proton

✗ Its dynamics are in principle calculable



✗ A few charm quarks produced (always by  $c\bar{c}$  pairs)

+ A few bound to form a charmonia ( $J/\psi \approx 3.1 \text{ GeV}/c^2$ )

✗ Fast formation time  $\approx 10^{-25} \text{ s}$

✗ “Slow” decay rate  $\approx 10^{-21} \text{ s}$

✗ Excellent QGP ( $\approx 10^{-23} \text{ s}$ ) probe




$J/\psi$

# 1986: J/ψ IN QGP

Matsui & Satz, PLB178 (1986) 416


- ✗ J/ψ should melt in the QGP, just above  $T_c$ 
  - + Due to screening of the  $c\bar{c}$  interaction
  - + “Unambiguous signature of quark gluon plasma”



\* The magical potion should make charmonia disappear...

# THE NAIVE PICTURE OF J/ $\Psi$ SUPPRESSION

- ✗ Creation of  $c\bar{c}$  (always in pairs)
- ✗ Separated by QGP



Sorry,  
powerpoint  
animated...

→ D meson

(gluons and secondaries not depicted)


# 1986-92: 1<sup>ST</sup> J/ $\psi$ SUPPRESSION @ SPS

✗ Quickly, NA38 saw J/ $\psi$  suppression

- + normalized to the  $\mu^+\mu^-$  continuum
- + in S+U versus p+U collisions

✗ But this was an illusion


- + not due to QGP !
- + Nuclear absorption



\* Charmonia disappear by (nuclear) dark magic

# J/ $\Psi$ NORMAL NUCLEAR ABSORPTION

- ✗ @SPS, nuclei take  $5 \times 10^{-24}$  s to cross (1.6 fm/c)
- ✗ They can separate  $c\bar{c}$  (formation time  $\approx 10^{-25}$  s)



# 1995–2000: ANOMALOUS SUPPRESSION

## ✘ $J/\psi$ / Drell-Yan

- + DY =  $q\bar{q} \rightarrow \text{gamma} \rightarrow \mu\mu$
- + Unaffected by QGP

## ✘ As a function of L

- + Nuclear thickness
- + Related to centrality


## ✘ In p+A, S+U collisions:

- +  $\exp(-\sigma_{\text{abs}} \rho_0 L)$
- + Normal nuclear absorption

## ✘ In central Pb+Pb:

- + Anomalous suppression
- + Likely to be due to QGP !

$J/\psi$  / Drell-Yan



\* Charmonia finally vanish in the plasma!



# AN ANNOUNCEMENT


- ✘ Based on this (and a bit more) CERN announced the discovery of QGP on February 10<sup>th</sup> 2000
- ✘ First collisions at RHIC began soon after
- ✘  $J/\psi$  was expected to melt further...



Organisation Européenne pour la Recherche Nucléaire  
European Organization for Nuclear research  
Laboratoire Européen pour la Physique des Particules  
European Laboratory for Particle Physics  
Europäisches Laboratorium für Teilchenphysik  
Laboratorio europeo per la fisica delle particelle

PRESS RELEASE

## New State of Matter created at CERN




At a special seminar on 10 February, spokespersons from the experiments on CERN\* 's Heavy Ion programme presented compelling evidence for the existence of a new state of matter in which quarks, instead of being bound up into more complex particles such as protons and neutrons, are liberated to roam freely.

Theory predicts that this state must have existed at about 10 microseconds after the Big Bang, before the formation of matter as we know it today, but until now it had not been confirmed experimentally. Our understanding of how the universe was created, which was previously unverified theory for any point in time before the formation of ordinary atomic nuclei, about three minutes after the Big Bang, has with these results now been experimentally tested back to a point only a few microseconds after the Big Bang.

# J/ψ @ RHIC, WHAT REFERENCE?

- ✗ No Drell-Yan to refer to
- ✗ We define a nuclear modification factor  $R_{AA}$ 
  - + J/ψ normalized by the number of elementary collisions  $N_{coll}$
- ✗ Limited open charm (Dmesons =  $u\bar{c}...$ ) results:
  - ≈ 25% uncertainty
  - + Do scale with  $N_{coll}$
  - + Same initial state a J/ψ


$$R_{AA} = \frac{dN^{AuAu}}{dN^{pp} \times \langle N_{coll} \rangle}$$



\* Charm quarks have reached immortality

# J/ $\Psi$ @ RHIC AND @ SPS

- ✘ Unexpectedly similar pattern
  - + Energy density is higher @ RHIC
  - + Different normal suppression
  
- ✘ This brings the idea of sequential melting...



\* RHIC alchemists change Pb in Au but do not melt more charmonia


# THE IDEA OF SEQUENTIAL MELTING

✘ Some  $J/\psi$  come from excited  $c\bar{c}$  states:

- +  $\Psi' \rightarrow J/\Psi + \dots$
- +  $X_c \rightarrow J/\Psi + \text{photon}$
- + Ratio not known better than 10%
- +  $J/\Psi \approx 0.7 J/\Psi + 0.2 \Psi' + 0.1 X_c$

✘ They melt at different temperatures

- + Theoretically not under perfect control, but some says:
  - ✘  $T(J/\psi) \approx 2 T_c$
  - ✘  $T(\psi') \approx T(X_c) \approx 1 T_c$




- ✘ Anomalous suppression: compatible with 0.7 @SPS
- ✘ Direct  $J/\psi$  could survive both @SPS and @RHIC

→ A good idea, but there is another result @ RHIC...

# MORE OR LESS J/ $\Psi$ @ RHIC

- ✘ Higher suppression in the forward direction,
- ✘ while the energy density should be lower there...
- ✘ Density-induced enhancement mechanism ?




\* Charmonia may be reappearing by enchantment

# THE IDEA OF RECOMBINATION

- ✘ A lot of  $c\bar{c}$  pairs
  - + (10 to 20 in central collisions @ RHIC)
- ✘ Uncorrelated  $c\bar{c}$  could meet at freeze-out and form new charmonia
- ✘ Enhanced production
  - +  $N(J/\psi) \propto (c\bar{c})^2$
- ✘ Suppression vs. enhancement

→ A recombination prediction (among many others)




Grandchamp et al., PRL 92,(2004) 212301



→ A good idea, but predictions not enough constrained... 30

# THE (LOCAL) RECOMBINATION


- ✘ Statistical hadronization model
- ✘ Forward suppression indeed higher
  - + Yellow : initial experimental uncertainty from  $c\bar{c}$  production



Andronic et al., (GSI) NPA789 (2007) 334  
 PBM and Stachel, PLB 490 (2000) 196

# THE NAIVE PICTURE OF RECOMBINATION

- ✗ Uncorrelated  $c\bar{c}$  could meet in QGP



Sorry,  
powerpoint  
animated...

(gluons and secondaries not depicted)





# NORMAL SUPPRESSION @ RHIC ?

✘ Not well measured

+ Very low p+A statistics


✘ Large uncertainty once extrapolated to A+A



# A LAST J/ψ PICTURE

- ✘ J/ψ in A+A divided by normal suppression extrapolations (poorly known @ RHIC)
  - + Need better constraint on normal suppression
  - + Could account for rapidity dependence ?
- ✘ However J/ψ do melt anomalously...
  - + In particular direct J/ψ

J/ψ survival probability in QGP





# ANOMALOUS CONCLUSION

- ✘  $J/\psi$  melt anomalously @ SPS and RHIC
- ✘ Not understood @ RHIC
  - + Less density = more suppression !
    - ✘ Recombination ?
    - ✘ Normal suppression ? (+ sequential melting ?)
  - + Large uncertainties, but new measurements are expected!
- ✘ However, even if the details are not understood, this is one of the QGP signs @ RHIC !

Charm quarks and the golden signatures

# OTHER CHARMS

---



# GOLDEN SIGNATURES @ RHIC

---


- ✘ Two striking features of RHIC data (at least)
  1. “Jet quenching ”
  2. “Elliptic flow”
- ✘ Measured with numerous light particles
- ✘ Are signatures of the quark gluon plasma
  
- ✘ How heavier (charm) particles behave?

# 1. JET QUENCHING


- ✘ RHIC smoking gun signature !
  - + Two PRL covers
- ✘ Energy loss in the matter, looking at high  $p_T$  ( $>2\text{GeV}$ )
  - + Mostly from jet fragmentation
- ✘ “Jet quenching”



## Au-Au (80-92%)



## d+Au (60-88%)




# MOST PERIPHERAL COLLISIONS...


(slightly old, but pedagogical, data)

PHENIX, PRL 91 (2003) 072303

## Au-Au (50-60%)



## d+Au (40-60%)




# LESS PERIPHERAL COLLISIONS...

(slightly old, but pedagogical, data)


PHENIX, PRL 91 (2003) 072303



## Au-Au (20-30%)



## d+Au (20-40%)




# MORE CENTRAL COLLISIONS...


(slightly old, but pedagogical, data)

PHENIX, PRL 91 (2003) 072303


## Au-Au (0-10%)



## d+Au (0-20%)



**MOST CENTRAL COLLISIONS!**




(slightly old, but pedagogical, data)

PHENIX, PRL 91 (2003) 072303



## 2. ELLIPTIC FLOW “ $V_2$ ”

PHENIX, PRL 91 (2003) 182301  
 Huovinen & al, PLB 503 (2001) 58



✘ Pressure gradient

✘  $V_2 = \langle \cos 2\phi \rangle$



→ Strong collective behavior

# IDEAL HYDRODYNAMICS

## ✗ Ideal hydrodynamics...

- + QGP EoS,
- + Early thermalization
  - ✗ (0.6 fm/c)
- + High density
  - ✗ ( $\approx 30 \text{ GeV/fm}^3$ )

## ✗ Little need for viscosity!

## ... reproduces fairly well

1. Single hadron  $p_T$  spectra
    - ✗ (mass dependence)
    - ✗  $\langle \beta_T \rangle \approx 0.6$
  2. Elliptic flow
- ✗ PQG is not the foreseen ideal partonic gas!


- Quarks are strongly interacting (“sQGP”)
- In a liquid like manner (“perfect fluid”)

# HEAVY QUARKS ?

PHENIX, PRL98, (2007) 172301

- ✘ Electrons from heavy flavour's decay (mostly charm, but some beauty)
- 1. Most of charm (low  $p_T$ ) scale with  $N_{coll}$  but high  $p_T$  are suppressed
  - + As much as light particles
- 2. Charm quark also have large elliptic flow
  - + Thermalization?
- ✘ Two surprises !
  - + Not well understood yet

D, B  $\rightarrow$  e + ...



\* Charm quarks have charming collective behaviours

# CONCLUSIONS

---

- ✘ The quark gluon plasma is surprising
- ✘ Unambiguous signatures become ambiguous
  - +  $J/\psi$  suppression
- ✘ Others become golden signature
  - + Jet quenching
  
- ✘ This, is the charm of this field...

# EPISODE 4: THE FUTURE

1. Back to CERN, always higher in energy

+ (i.e. temperature)

✘ Large Hadron Collider :



+ Pb+Pb @  $\sqrt{s_{nn}} = 5.5$  TeV

+ First collisions (p+p) in 2008?

✘ Recombined  $J/\psi$  could raise with centrality?

✘ Golden sign  $\rightarrow$   
(empirical line)

But see Andronic et al.,  
NPA789 (2007) 334





2. @ GSI, FAIR gets higher baryon densities

$\approx 10$  times nuclear

+ Critical point?

+ Collision by 2015?

# EPISODE 4: THE FUTURE







Back-up slides

**THAT'S ALL FOLKS**

---

# ☺ THE ORIGIN OF (MY) MASS...

~ 98% from QCD + 02% from Higgs !



~ 98% poorly understood + 02% not yet seen...

✘ We are mostly made of confinement...

✘ Thus, let's look at deconfinement...

(ok, this is only ~5% of the universe ☹)


# TOTAL MULTIPLICITY (AND $E_T$ )

- ✗  $dN_{ch}/d\eta|_{\eta=0} \sim 670$
- ✗ (6000 particles total)
- ✗ Less than expected!



- + 1000 from p+p fragmentation
- + Low  $x_{Bj}$  gluon start to overlap, recombine, saturate...
- + (even more at forward rapidity)
- + “Color Glass Condensate”

→ The (initial) matter saturates




PRL 91 (2003) 052303

- ✗  $dE_T/d\eta|_{\eta=0}$  related to energy density
- ✗  $\epsilon > 6 \text{ GeV}/\text{fm}^3 > \epsilon_c !$

# HOW TO ESTIMATE ENERGY DENSITY ?

~ Transverse energy @  $y=0$



Bjorken formula

$$\varepsilon = \frac{1}{\pi R^2 \tau_0} \times \left. \frac{dE_T}{dy} \right|_{y=0}$$


$\tau_0$  formation time  
0,35 à 1 fm/c

R = nuclear radius  
 $1.18 A^{1/3}$  fm

$$\varepsilon > 6 \text{ GeV/fm}^3$$

Bjorken, PRD27 (1983) 140

# DRELL-YAN NCOLL SCALING






# J/ $\Psi$ VERSUS MODELS...



R. Rapp et al., nucl-th/0608033

Yan et al., PRL97 (2006) 232301

## ✗ Recombination...




✗ SPS like...



Capella & Ferreiro,  
hep-ph/0610313

- R. Rapp et al. PRL 92, 212301 (2004)
- R. Thews et al, Eur. Phys. J C43, 97 (2005)
- Yan, Zhuang, Xu, PRL97 (2006) 232301
- Bratkovskaya et al., PRC 69, 054903 (2004)
- A. Andronic et al., NPA789 (2007) 334

# HIGH $P_T$ SUPPRESSION




Photon: PRL94 (2005) 232301

Eta: PRC 75 (2007) 024909

Vitev & Gyulassy PRL82 (2002) 252301


→ The matter is dense !  $>1000$  gluons per  $\Delta y$

# BACK TO BACK JETS






## Peripheral collisions (60-80%)



Take a “trigger” particle ( $p_T > 4\text{GeV}$ ) and look at the others ( $p_T > 2\text{GeV}$ ) azimuth

## Central collisions (0-5%)



In central collision, opposite jets disappear because of jet quenching


# BACK TO BACK JETS

# ANOTHER LOOK TO JET QUENCHING...

# BACK TO BACK (D+AU)

STAR, PRL 91 (2003) 072304

- As always, it is very important to check for d+Au




The matter is opaque!

# THERMAL RADIATION

The matter is hot !

PHENIX, PRL94 (2005) 232301  
 Virtual photon are preliminary  
 + various theory papers



Photon spectrum is fairly reproduced by:

@ high  $p_T$  prompt photon (pQCD)

@ low  $p_T$  thermal photons ( $T \sim 400 - 600$  MeV  $\gg T_c$ )