

# Status of the Quark Gluon Plasma (QGP) search at RHIC

- A PHENIX perspective (\*) -

Nuclear and Particle Physics Colloquium  
LNS, MIT, April 4th, 2005

**David d'Enterria**  
**Nevis Labs, Columbia University, NY**

# Overview/Summary

## Introduction:

- Goal of **high-energy A+A collisions** ?

Study/characterize **Quantum Chromo (many-body) Dynamics**

(By comparing A+A to: **p+A** = “cold QCD medium”, **p+p** = “QCD vacuum” colls.)

**Head-on Au+Au collisions @ RHIC energies ( $\sqrt{s}=200$  GeV) produce a strongly interacting system:**

- with the **highest energy densities ever** achieved at the lab:  $\epsilon > 5 \text{ GeV/fm}^3$
- with a **strong degree of collectivity** at very short time-scales:  $\tau_0 < 1 \text{ fm/c}$
- that behaves like an **nearly ideal (hydrodynamical) fluid**:  $dN/dp_T$  ( $p_T < 2 \text{ GeV/c}$ )
- that reaches **chemical equilibrium at** (or before) **hadronization**:  $T_{\text{chem}} \sim T_{\text{crit}}$
- with the **largest initial gluon densities ever** measured:  $dN^g/dy \sim 1000$
- with **degrees of freedom** consistent with **constituent quarks**

## Summary & open questions


# High-energy heavy-ion physics program (in 4 plots)

$$\mathcal{L} = \frac{1}{4g^2} G_{\mu\nu}^a G_{\mu\nu}^a + \sum_f \bar{\psi}_f (i \gamma^\mu D_\mu + m_f) \psi_f$$

where  $G_{\mu\nu}^a \equiv \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf_{abc} A_\mu^b A_\nu^c$   
and  $D_\mu = \partial_\mu + i t^a A_\mu^a$  ( $\alpha_s = g^2/4\pi$ )



$$\alpha_s(Q^2) \sim 1/\ln(Q^2/\Lambda^2), \Lambda \sim 200 \text{ MeV}$$

1. Learn about 2 basic properties of strong interaction: **(de)confinement**, **chiral symm. breaking (restoration)**




4. Study the regime of **non-linear** (high density) many-body **parton dynamics** at small-x (**CGC**)

2. Study the **phase diagram** of QCD **matter**: esp. produce & study the **QGP**









3. Probe quark-hadron **phase transition** of the **primordial Universe** (few  $\mu\text{sec}$  after the Big Bang)



# The "Little Bang" in the lab.

- High-energy **nucleus-nucleus collisions**: fixed-target reactions ( $\sqrt{s}=20$  GeV, SPS) or colliders ( $\sqrt{s}=200$  GeV, RHIC.  $\sqrt{s}=5.5$  TeV, LHC)
- **QGP** expected to be formed in a **tiny region** ( $\sim 10^{-14}$  m) and to last **very short times** ( $\sim 10^{-23}$  s).
- **Collision dynamics**: Diff. observables sensitive to diff. react. stages





# Relativistic Heavy-Ion Collider (RHIC) @ BNL

## Specifications:

**3.83 km** circumference

**2** independent rings:

- 120 bunches/ring
- 106 ns crossing time

**A + A collisions @  $\sqrt{s}_{NN} = 200$  GeV**

Luminosity:  $2 \cdot 10^{26} \text{ cm}^{-2} \text{ s}^{-1}$  ( $\sim 1.4$  kHz)

**p+p collisions @  $\sqrt{s}_{\text{max}} = 500$  GeV**

**p+A collisions @  $\sqrt{s}_{\text{max}} = 200$  GeV**

## 4 experiments:

**BRAHMS, PHENIX, PHOBOS, STAR**


## Runs 1 - 5 (2000 – 2005):

**Au+Au @ 200, 130, 62.4 GeV**

**p+p @ 200 GeV**




**d+Au @ 200 GeV**

**Cu+Cu @ 200, 62.4 GeV**




# The 4 RHIC experiments

## STAR Detector




## BRAHMS detector

## PHOBOS Detector







# RHIC Au+Au luminosities



- RHIC (Au+Au) is currently running at **~2x design luminosity**

	max energy [GeV/u]	no of bunches	ions/bunch [ $10^9$ ]	$\beta^*$ [m]	emittance [mm mrad]	$\mathcal{L}_{peak}$ [ $10^{26} \text{ cm}^{-2} \text{ s}^{-1}$ ]	$\mathcal{L}_{store,ave}$ [ $10^{26} \text{ cm}^{-2} \text{ s}^{-1}$ ]	$L_{week}$ [ $\mu\text{b}^{-1}$ ]
Run-1 (FY2000)	65	55	0.3	3	15-40	0.3	0.2	4
Run-2 (FY2001/2002)	100	55	0.5	1	15-40	3.7	1.5	24
Run-4 (FY2004)	100	45	1.1	1	15-40	15	4	160
Design	100	55	1.0	2	15-40	9	2	50
Enhanced design	100	112	1.0	1	15-40	30	8	300

# Au+Au collisions @ 200 GeV



~ 700 charged particles per unit rapidity at midrapidity (top 5% central)




## (1) Energy densities at RHIC

- The highest energy densities ever achieved at the lab:  $\epsilon > 5 \text{ GeV/fm}^3$


# Energy density (Au+Au @ 200 GeV, y=0)

• Bjorken estimate:  $\epsilon_{Bj} = \frac{dE_T}{dy} \frac{1}{\tau_0 \pi R^2}$

(longitudinally expanding plasma)



- $dE_T/d\eta$  at mid-rapidity measured by calorimetry (using PHENIX EMCal as *hadronic* calorimeter:  $E_T^{\text{had}} = (1.17 \pm 0.05) E_T^{\text{EMCal}}$ )



$\langle dE_T/d\eta \rangle \sim 650$  GeV (top 5% central)

(~70% larger than at CERN-SPS)


$$\epsilon_{\text{Bjorken}} \sim 5.0 \text{ GeV/fm}^3$$

> QCD critical density ( $\sim 1$  GeV/fm<sup>3</sup>)

PHENIX Collab.  
PRL 87, 052301 (2001)  
nucl-ex/0104015

# 1 fm/c thermalization time ?

- Not unrealistic at RHIC... (for the 1<sup>st</sup> time:  $\tau_{\text{therm}} > \tau_{\text{cross}} = 2R/\gamma \sim 0.15 \text{ fm/c}$ )
- Time evolution of energy density in longitud. expanding system:  $\epsilon \sim 1/\tau$






## (2) Elliptic flow at RHIC

- Strong degree of collectivity at very short time-scales:  $\tau_0 < 1 \text{ fm}/c$


# Elliptic flow

- Initial anisotropy in x-space in non-central collisions (overlap) translates into final **azimuthal asymmetry** in p-space (transverse to react. plane)



$$\frac{dN}{d\phi} \propto 1 + 2v_2 \cos 2(\phi - \Phi_{RP})$$


Elliptic flow =  $v_2$   
 2<sup>nd</sup> Fourier coefficient  
 of  $dN/d\phi$



PHENIX Collab.  
 PRL 89, 212301 (02)  
 nucl-ex/0204005


# Elliptic flow

- Initial anisotropy in x-space in non-central collisions (overlap) translates into final **azimuthal asymmetry** in p-space (transverse to react. plane)



$$\frac{dN}{d\phi} \propto 1 + 2v_2 \cos 2(\phi - \Phi_{RP})$$


Elliptic flow =  $v_2$   
2<sup>nd</sup> Fourier coefficient  
of  $dN/d\phi$



PHENIX Collab.  
PRL 89, 212301 (02)  
nucl-ex/0204005

“1 image = 1000 words” ...

“Elliptic flow” in low -T (strongly coupled) **Li atoms**:




K.M.O'Hara,  
Sci. 289, 2179  
(2002)  
T. Bourdel et al.  
PRL 91, 020402  
(2003)




# Elliptic flow

- Initial anisotropy in x-space in non-central collisions (overlap) translates into final **azimuthal asymmetry** in p-space (transverse to react. plane)



$$\frac{dN}{d\phi} \propto 1 + 2v_2 \cos 2(\phi - \Phi_{RP})$$

Elliptic flow =  $v_2$   
2<sup>nd</sup> Fourier coefficient  
of  $dN/d\phi$




1. Truly **collective** effect (absent in p+p collisions).

2. **Early-state** phenomenon:  
develops only in 1<sup>st</sup> instants of reaction.  
Strongly self-quenches after  $t \sim 1$  fm/c


Time evolution of ellipsoid eccentricity:

$$\varepsilon = \frac{\langle y^2 \rangle - \langle x^2 \rangle}{\langle y^2 \rangle + \langle x^2 \rangle}$$



# Elliptic flow at RHIC



- Large  $v_2$  signal at RHIC:  
Exhausts hydro limit for  $p_T < 1.5$  GeV/c



⇒ Strong (collective) pressure grads.  
⇒ Large & fast parton rescattering:  
early thermalization.

- $\sqrt{s}$ -dependence of  $v_2$ :  
~50% increase from CERN-SPS  
(apparent saturation within 62-200 GeV)

- Mass dependence of  $v_2$   
consistent w/ hydrodynamics too:




### (3) Soft particle production at RHIC

- A system that behaves like an **nearly ideal (hydrodynamical) fluid** and reaches **chemical equilibrium** at (or before) **hadronization**:  $T_{\text{chem}} \sim T_{\text{crit}}$




# Soft particle spectra

- Bulk  $\pi^\pm$ ,  $K^\pm$ ,  $p(\bar{p})$  spectra reproduced by **hydro** w/ **QGP EOS** at  $\tau_0 = 0.6$  fm/c




Strong radial **collective flow** built-up at freeze-out:  $\langle\beta_T\rangle \approx 0.6$



# Soft particle spectra


- Bulk  $\pi^\pm$ ,  $K^\pm$ ,  $p(\text{pbar})$  spectra reproduced by **hydro** w/ **QGP EOS** at  $\tau_0 = 0.6 \text{ fm/c}$

Au+Au central ( $b = 2.6 \text{ fm}$ )




D.d'E. & Peressounko  
nucl-th/0503054

Strong radial **collective flow**  
built-up at freeze-out:  $\langle \beta_T \rangle \approx 0.6$



- **Ratios** of hadron yields consistent w/ system at **chemical equilibrium** at hadronization ( $T_{\text{chem. freeze-out}} \sim T_{\text{crit}}$ ) :



- 1 ratio (e.g.  $p/\bar{p}$ ) determines  $\mu/T$

- 2<sup>nd</sup> ratio (e.g. K/pi) provides  $T_{\mu}$ .

PBM, Redlich, Stachel  
nucl-th/0304013  
Kaneta, Xu  
nucl-th/0405068

- David d'Entieria (Columbia Univ.)





## (4) Hard QCD production at RHIC

- The largest initial gluon densities ever measured:  $dN^g/dy \sim 1000$

# Hard QCD probes (I)

● Hard probes: High- $p_T$ , jets, direct  $\gamma$ , heavy-quarks (D, B), ...

1. **Early production** ( $\tau \sim 1/p_T < 0.1$  fm/c) in parton-parton scatterings with large  $Q^2$ :  
Closest experimental probes to underlying QCD ( $q, g$ ) degrees of freedom.
2. Direct probes of **partonic phase(s)**  $\Rightarrow$  Sensitive to QCD medium properties:



# Hard QCD probes (II)

## 3. Production yields theoretically calculable via perturbative-QCD:

“Factorization theorem”:

$$d\sigma_{AB \rightarrow hX} = A \cdot B \cdot f_{a/A}(x_a, Q^2) \otimes f_{b/B}(x_b, Q^2) \otimes d\sigma_{ab \rightarrow cd} \otimes D_{h/c}(z_c, Q^2)$$

Independent scattering of “free” partons:

$$f_{a/A}(x, Q^2) = A f_{a/p}(x, Q^2)$$

$A+B$  = “simple superposition of  $p+p$  collisions”


$$d\sigma_{AB \rightarrow \text{hard}} = A \cdot B \cdot d\sigma_{pp \rightarrow \text{hard}}$$

At impact parameter  $b$ :


$$dN_{AB \rightarrow \text{hard}}(b) = T_{AB}(b) \cdot d\sigma_{pp \rightarrow \text{hard}}$$

geom. nuclear overlap at  $b$

$T_{AB} \sim \# \text{ NN collisions}$  (“ $N_{\text{coll}}$  scaling”)




## Nuclear Modification Factor:




# High $p_T$ p+p baseline data well described by pQCD

- Good theoretical (NLO pQCD) description:

$$p+p \rightarrow \pi^0 X$$



$$p+p \rightarrow h^\pm X \text{ (non singly diffractive)}$$




- Well calibrated (experimentally & theoretically) p+p references at hand




# “NN scaling” in Au+Au @ 200 GeV: Direct Photons

- Direct photon production in Au+Au (all centralities) consistent w/ p+p incoherent scattering (“NN-scaled” pQCD) predictions:




$$R_{AA}(p_T, y; b) = \frac{\text{“hot/dense QCD medium”}}{\text{“QCD vacuum”}} = \frac{d^2 N_{AA}/dy dp_T}{\langle T_{AA}(b) \rangle \cdot d^2 \sigma_{pp}/dy dp_T},$$



- Direct photon production in Au+Au unmodified by QCD medium.


# Suppressed high $p_T$ hadroproduction in Au+Au @ RHIC !

Au+Au  $\rightarrow \pi^0 X$  (peripheral)




Peripheral data **agree** well with **p+p** (data & pQCD) plus  $N_{\text{coll}}$ -scaling

Au+Au  $\rightarrow \pi^0 X$  (central)




Strong **suppression** in central Au+Au collisions

# Suppressed high $p_T$ hadroproduction @ RHIC




D.d'E., HP'04  
nucl-ex/0504001

$R_{AA} \ll 1$ : well below pQCD (collinear factorization) expectations for hard scattering cross-sections




Discovery of  
 high  $p_T$  suppression  
 (one of most significant  
 results @ RHIC so far)

# Unquenched d+Au production at high $p_T$



PHENIX.  
PRL91, 072303(2003)



- Conclusion: High  $p_T$  suppression in central Au+Au due to final-state effects (absent in “control” d+Au experiment)



# “Jet quenching” predictions

- Multiple final-state non-Abelian (gluon) radiation off the produced hard parton induced by the traversed dense medium.

- Parton energy loss  $\propto$  medium properties:

$$\Delta E_{\text{loss}} \sim \rho_{\text{gluon}} \quad (\text{gluon density})$$

$$\Delta E_{\text{loss}} \sim \Delta L^2 \quad (\text{medium length})$$


- Energy is carried away by gluonsstrahlung inside jet cone:  $dE/dx \sim \alpha_s \langle k_T^2 \rangle$

- Correction for expanding (1-D) plasma :

$$\Delta E_{1-D} = (2\tau_0/R_A) \cdot \Delta E_{\text{static}} \sim 15 \cdot \Delta E_{\text{static}} \quad (\tau_0=0.2 \text{ fm}/c, R_A=6 \text{ fm})$$

- Prediction I: Suppression of high  $p_T$  leading hadrons

- Prediction II: Disappearance of back-to-back (di)jet correlations



# “Jet quenching” model vs. data (I)

- Dense medium properties from pQCD+ final-state parton energy loss models:

- ★ Initial gluon densities:

$$dN_g/dy \sim 1100 \quad [\text{Vitev \& Gyulassy}]$$

- ★ Opacities:

$$\langle n \rangle = L/\lambda \approx 3 - 4 \quad [\text{Levai et al.}]$$

- ★ Transport coefficients:


$$\langle q_0 \rangle \sim 14 \text{ GeV/fm}^2 \quad [\text{BDMPS, F.Arleo}]$$

$$[\text{Salgado-Wiedemann}]$$

- ★ Medium-induced radiative energy losses:

$$dE/dx \approx 0.25 \text{ GeV/fm} \quad (\text{expanding})$$

$$dE/dx|_{\text{eff}} \approx 14 \text{ GeV/fm} \quad (\text{static source}) \quad [\text{X.N.Wang}]$$




- Large opacities imply fast thermalization.
- All these values imply energy densities well above  $\epsilon_{\text{crit QCD}}$  (in thermalized syst.)


# “Jet quenching” model vs. high $p_T$ suppression (II)

## sqrt(s)-dependence:


D.d'E., HP'04  
nucl-ex/0504001




## Non-Abelian energy loss:



## Centrality dependence:





## Reaction-plane (in-medium path length) dependence:



# “Jet quenching”: modified (di)jet structure

- Strongly modified  $dN_{\text{pair}}/d\phi$  high  $p_T$  correlations in central Au+Au:

Au+Au central





- Strongly non-Gaussian away-side (“dip”) peak.

PHENIX Preliminary

- Standard back-to-back di-jet topology:

Au+Au periph






## (5) Hadron production at intermediate $p_T$

- Degrees of freedom consistent with constituent quarks


# Unsuppressed baryon production

- $R_{cp}$  (ratio central/peripheral) at intermediate  $p_T = 2 - 4$  GeV/c:

Baryons:  $p, \bar{p}, \Lambda, \bar{\Lambda}$  **NOT** (or much less) suppressed in central Au+Au.




Mesons:  $\pi^0, K_s^0, \eta$ ,  
equally suppressed.



- Particle composition **inconsistent with** known (universal) **fragmentation functions**.
- **Additional production mechanism** for baryons in the intermediate  $p_T$  range

# Enhanced (anti)proton/pion ratio

- **Central** Au+Au:  $p/\pi \sim 0.8$  (at  $p_T = 2 - 4$  GeV/c) at variance with perturbative production mechanisms (favour lightest mesons).
- **Periph.** Au+Au:  $p/\pi \sim 0.2$  as found in  $p+p$  (ISR,FNAL) &  $e+e^-$  jet fragmentation




PHENIX Collab.  
PRL91:172301(2003)

# Enhanced baryonic elliptic flow

- Different  $v_2$  saturation for mesons and baryons:


$$\begin{aligned} v_2^{\text{meson}} &> v_2^{\text{baryon}} && \text{at low } p_T \\ v_2^{\text{meson}} &\approx v_2^{\text{baryon}} && \text{at } p_T \approx 2 \text{ GeV}/c \\ v_2^{\text{meson}} &< v_2^{\text{baryon}} && \text{at higher } p_T \end{aligned}$$



- Simple  $v_2$  scaling behaviour if  $v_2$  and  $p_T$  are normalized by number of constituent quarks:

$$\begin{aligned} n &= 2 \text{ mesons} \\ n &= 3 \text{ baryons} \end{aligned}$$




(“universal” parent quark flow)






# “Quark recombination” models vs. data

- Anomalous baryon enhancement & quark number scaling of  $v_2$  at  $p_T = 2\text{--}5 \text{ GeV}/c$  explained by “quark recombination” (coalescence) in dense (thermal) medium:



- Rethink hadronization at interm.  $p_T$  at RHIC !  
Phase space filled with partons  
Recombine quarks into hadrons



# Summary

## 1. Energy densities:

- Maximum  $dE_T/d\eta \sim 600$  GeV at midrapidity consistent w/ **initial  $\epsilon > 5$  GeV/fm<sup>3</sup>  $> \epsilon_{\text{crit}}$**

## 2. Elliptic flow:

- **Strong** elliptic flow  $v_2$  consistent w/ **short thermalization times  $\tau_0 \sim 1$  fm/c**

## 3. Soft particle spectra:

- Shapes & yields consistent w/ **hydrodyn.** (thermal+coll. velocity) source emission
- Particles ratios consistent w/ **chemically equilibrated** system before hadronization

## 4. Hard particle spectra:

- **Strong high  $p_T$  suppression** in central A+A (compared to p+p, p+A & pQCD) consistent w/ **final-state partonic energy loss** in dense system:  **$dN^g/dy \sim 1100$**

## 5. Intermediate $p_T$ spectra:

- **Enhanced baryon** yields &  $v_2$  (compared to meson) consistent w/ **quark recombination** mechanisms in a thermal and dense system

**All observations consistent with formation of thermalized dense partonic matter in central Au+Au collisions**

# Outline: What do we need to claim “QGP !” at RHIC ?

Somebody called us “overly conservative” but we (experimentalist @ RHIC) would like at least to know ...



(1) What is the **temperature** of the produced system?

# Outline: What do we need to claim “QGP !” at RHIC ?

Somebody called us “overly conservative” but we (experimentalist @ RHIC) would like at least to know ...

(1) What is the **temperature** of the produced system?

= “do we see **thermal photons** from the radiating plasma” ?




D.d'E. & D.Peressounko  
nucl-th/0503054

# Outline: What do we need to claim “QGP !” at RHIC ?

Somebody called us “overly conservative” but we (experimentalist @ RHIC) would like at least to know ...

(1) What is the **temperature** of the produced system?

= “do we see **thermal photons** from the radiating plasma” ?



Current **experimental upper limits** in “interesting” region  $p_T = 1 - 3 \text{ GeV/c}$  preclude a quantitative answer ...

**Wait for Run-4 data**



# Outline: What do we need to claim “QGP !” at RHIC ?


Somebody called us “overly conservative” but we (experimentalist @ RHIC) would like at least to know ...

- (1) What is the **temperature** of the produced system?
- (2) Is the **system deconfined** ?

# Outline: What do we need to claim “QGP !” at RHIC ?

Somebody called us “overly conservative” but we (experimentalist @ RHIC) would like at least to know ...



- (1) What is the **temperature** of the produced system?
- (2) Is the **system deconfined** ?  
= “do we see the predicted **melting of quarkonia** bound states (seen at CERN-SPS) ?



# Outline: What do we need to claim “QGP !” at RHIC ?

Somebody called us “overly conservative” but we (experimentalist @ RHIC) would like at least to know ...



- (1) What is the **temperature** of the produced system?
- (2) Is the **system deconfined** ?  
= “do we see the predicted **melting of quarkonia** bound states (seen at CERN-SPS) ?



# Outline: What do we need to claim “QGP !” at RHIC ?

Somebody called us “overly conservative” but we (experimentalist @ RHIC) would like at least to know ...

- (1) What is the **temperature** of the produced system?
- (2) Is the **system deconfined** ?  
= “do we see the predicted **melting of quarkonia** bound states (seen at CERN-SPS) ?



Again the answer is ...

**Wait for Run-4 data !**



**Stay tuned !**

**backup slides ...**



# Jet production in hot&dense QCD: modified (di)jet structure (I)


- Discovery of “mono-jet” like topologies (away-side disappearance):



# OK ... but didn't we hear the same at CERN-SPS in 2000 ?




## Quark gluon plasma



Creation of a new state of matter, quark-gluon plasma, which probably existed just after the Big Bang.

# Suppressed high $p_T$ hadroproduction @ RHIC: $h^\pm$ vs $\pi^0$



- Inclusive **charged** hadrons **suppressed** by a factor  $\sim 4 - 5$  at  $p_T > 5$  GeV/c




- Universal** (PID-wise) suppression above  $p_T = 5$  GeV/c

# “Anomalous” particle composition: hadron/meson ratio

- p+p collisions: hadron/meson  $\sim 1.6$





- Au+Au (central): hadron/meson  $\sim 2.5$  at  $p_T = 1 - 4$  GeV/c (inconsistent w/ known fragmentation functions).



- Baryon enhancement limited to  $p_T < 5$  GeV/c ( $h^\pm/\pi \sim 1.6$ , perturb. ratio):  $h^\pm, \pi^0$  equally suppressed

# “ $N_{\text{coll}}$ scaling” in Au+Au @ 200 GeV: Total charm

- Open-charm indirect measurement via semi-leptonic channel:  $D \rightarrow e^\pm + X$
- Single  $e^\pm$  Au+Au spectra & total cross-section consistent w/  $N_{\text{coll}}$ -scaled p+p charm production:




- pQCD parton scattering holds for hard processes in Au+Au (all centralities).




# “ $N_{\text{coll}}$ scaling” in A+A @ 17, 31 GeV: High $p_T$ hadrons

- High  $p_T$   $\pi^0$  production in (0-10%) central A+A at SPS (and  $\alpha+\alpha$  @ ISR) energies consistent w/ “ $N_{\text{coll}}$ -scaling” (or Cronin enhancement):




# “ $N_{\text{coll}}$ scaling” in d+Au @ 200 GeV: High $p_T$ hadrons

d+Au @  $\sqrt{s_{NN}} = 200$  GeV




p+A @  $\sqrt{s_{NN}} = 20 - 40$  GeV




- Enhanced high  $p_T$  production in d+Au ( $R_{dAu} > 1$ ) also found in p+A at lower  $\sqrt{s}$  (“Cronin enhancement”):  $p_T$  broadening due to initial-state soft & semihard scattering.
- Expected pQCD behaviour ( $R_{pA,dA} \sim 1$ ) recovered for  $p_T > 8$  GeV/c

# Energy loss in a dense hadronic medium ?

- Hadronic transport models (HSD, UrQMD) or DPM-based models do not produce enough suppression. Additional pre-hadronic energy loss needed.



Cassing, Gallmeister, Bratkovskaya,  
Greiner, Stoecker, nucl-th/0312049





Capella, Ferreiro, Kaidalov, Sousa  
hep-ph/0403081

# Unsuppressed baryon production: not a mass effect !

- $R_{cp}$  (ratio central/peripheral) at intermediate  $p_T = 2 - 4$  GeV/c:

Baryons:  $p, \bar{p}, \Lambda, \bar{\Lambda}$  **NOT** (or much less) suppressed in central Au+Au.

Heavy  $\phi$  as suppressed as other mesons ( $\pi^0, k_s^0, \eta$ )



- Particle composition **inconsistent with** known (universal) **fragmentation functions**.
- **Additional production mechanism** for baryons in the intermediate  $p_T$  range