

ATLAS Japan Workshop
20 Dec 1997 @ Naruto
University of Education

Basic Properties of the Higgs Boson in the Standard Model (SM)

Kensuke Homma
ICEPP, University of Tokyo

Contents

1. Basics
2. Why we need Higgs in SM
3. Properties of SM Higgs

1. Basics

1-1 Review of Lagrangians

$$L \equiv \text{K.E.} - V$$

Classical

Field theory

$$L(q_i, \dot{q}_i, t) \quad \Leftrightarrow \quad \mathcal{L}(\phi, \frac{\partial \phi}{\partial x_\mu}, x_\mu)$$

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_i} \right) - \frac{\partial L}{\partial q_i} = 0 \quad \Leftrightarrow \quad \frac{\partial}{\partial x_\mu} \left(\frac{\partial \mathcal{L}}{\partial (\partial \phi / \partial x_\mu)} \right) - \frac{\partial \mathcal{L}}{\partial \phi} = 0$$

$$\mathcal{L} = \text{K.E.} + \text{Mass} + \text{Interaction (F-B, B-B)}$$

Fields	K.E. (and B-B)	Mass
Fermion	$i\bar{\psi}\partial_\mu\gamma^\mu\psi$	$-m\bar{\psi}\psi$
Complex Scalar	$(\partial^\mu\phi)^*(\partial_\mu\phi)$ $\phi \equiv \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2)$	$-m^2\phi^*\phi$
Abelian Boson	$-\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$ $F^{\mu\nu} \equiv \partial^\mu B^\nu - \partial^\nu B^\mu$	$+\frac{1}{2}m^2 B^\mu B_\mu$
Non-Abelian Boson	$-\frac{1}{4}W_{\mu\nu a}W_a^{\mu\nu}$ $W_a^{\mu\nu} \equiv \partial^\mu W_a^\nu - \partial^\nu W_a^\mu$ $+gf_{abc}W_b^\mu W_c^\nu$	$+\frac{1}{2}m^2 W_a^\mu W_{a\mu}$

Interaction F-B is described as $Q\bar{\psi}\gamma^\mu\psi A_\mu$

1-2 Local Gauge Invariance

Under the transformation, $\psi'(x) \equiv e^{i\alpha(x)}\psi(x)$

$$\psi'(x) = \psi(x) \text{ holds.}$$

For a massive fermion in a free potential

$$\mathcal{L} = i\bar{\psi}\gamma^\mu\partial_\mu\psi - m\bar{\psi}\psi$$

mass term is invariant

$$\psi \rightarrow e^{i\alpha(x)}\psi(x), \quad \bar{\psi} \rightarrow e^{-i\alpha(x)}\bar{\psi}(x)$$

K.E. term is not invariant

$$\partial_\mu\psi \rightarrow e^{i\alpha(x)}\partial_\mu\psi(x) + ie^{i\alpha(x)}\partial_\mu\alpha(x)\psi(x)$$

Introduce covariant derivative D_μ which transforms in the same way as ψ itself

$$D_\mu \rightarrow e^{i\alpha(x)}D_\mu$$

$$D_\mu \equiv \partial_\mu - ieA_\mu$$

$$A_\mu \rightarrow A_\mu + \frac{1}{e}\partial_\mu\alpha(x)$$

$$\begin{aligned}\mathcal{L} &= i\bar{\psi}\gamma^\mu D_\mu\psi - m\bar{\psi}\psi \\ &= i\bar{\psi}\gamma^\mu\partial_\mu\psi - m\bar{\psi}\psi + e\bar{\psi}\gamma^\mu\psi A_\mu\end{aligned}$$

By requiring the local gauge invariance,
a new vector field A_μ emerges !!!

1-3 Lagrangian of mass-less unphysical fields in SM

Fermions in SM

$$L = \begin{pmatrix} \nu_e \\ e \end{pmatrix}_L \quad \blacksquare \quad e_R \quad \times 3 \text{ generations}$$

$$Q = \begin{pmatrix} u \\ d \end{pmatrix}_L \quad u_R, d_R$$

Charges

$$Q = T_3 + \frac{Y}{2}$$

Q is electric charge

T is isospin

Y is weak-hypercharge

fermion	Q	T	T_3	Y
ν_{eL}	0	1/2	1/2	-1
e_L	-1	1/2	-1/2	-1
u_L	2/3	1/2	1/2	1/3
d_L	-1/3	1/2	-1/2	1/3
$\blacksquare$	$\blacksquare$	$\blacksquare$	$\blacksquare$	$\blacksquare$
e_R	-1	0	0	-2
u_R	2/3	0	0	4/3
d_R	-1/3	0	0	-2/3

$$f_{SM} \equiv \mathbf{L}, e_R, \mathbf{Q}, u_R, d_R$$

$$\mathcal{L}_{Ferm} = \sum_{f_{SM}} i \bar{f} \gamma^\mu D_\mu f$$

$$D_\mu = \partial_\mu - i g_1 \frac{Y}{2} B_\mu - i g_2 \frac{\tau^i}{2} W_\mu^i - i g_3 \frac{\lambda^a}{2} G_\mu^a$$

$$\text{where} \qquad \tau^i W_\mu^i \equiv \tau^1 W_\mu^1 + \tau^2 W_\mu^2 + \tau^3 W_\mu^3$$

$$\lambda^a G_\mu^a \equiv \sum_{a=1}^8 \lambda^a G_\mu^a$$

$$\mathbf{U(1)}$$

$$Y = \text{constant}$$

$$\mathbf{SU(2)}$$

$$\tau^1 = \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array} \right) \quad \tau^2 = \left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array} \right) \quad \tau^3 = \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array} \right)$$

$$\mathbf{SU(3)}$$

$$\lambda^1 = \left(\begin{array}{ccc} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right) \qquad \lambda^2 = \left(\begin{array}{ccc} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{array} \right) \qquad \lambda^3 = \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{array} \right)$$

$$\lambda^4 = \left(\begin{array}{ccc} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{array} \right) \qquad \lambda^5 = \left(\begin{array}{ccc} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{array} \right) \qquad \lambda^6 = \left(\begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{array} \right)$$

$$\lambda^7 = \left(\begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{array} \right) \qquad \lambda^8 = \frac{1}{\sqrt{3}} \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{array} \right)$$

1-4 Lagrangian of mass-less physical fields in SM

$$\begin{aligned}
 \mathcal{L}_{Ferm} = & \sum_{f=\nu_e, e, u, d} g_1 Q_f \bar{f} \gamma^\mu f A_\mu \\
 & + \sum_{f=\nu_e, e, u, d} \frac{g_2}{\cos \theta_w} (T_f^3 - Q_f \sin^2 \theta_w) \bar{f}_L \gamma^\mu f_L Z_\mu \\
 & + \sum_{f=e, u, d} \frac{g_2}{\cos \theta_w} (-Q_f \sin^2 \theta_w) \bar{f}_R \gamma^\mu f_R Z_\mu \\
 & + \frac{g_2}{\sqrt{2}} (\bar{u}_L \gamma^\mu d_L + \bar{\nu}_{eL} \gamma^\mu e_L) W_\mu^+ \\
 & + \frac{g_2}{\sqrt{2}} (\bar{d}_L \gamma^\mu u_L + \bar{e}_L \gamma^\mu \nu_{eL}) W^{\mu-} \\
 & + \sum_{q=u, d} \frac{g_3}{2} \lambda_{\alpha\beta}^a \bar{q}_\alpha \gamma^\mu q_\beta G_\mu^a
 \end{aligned}$$

where

$$\begin{aligned}
 W^+ &= (-W^1 + iW^2)/\sqrt{2} \\
 W^- &= (-W^1 - iW^2)/\sqrt{2} \\
 W^0 &= -W^3
 \end{aligned}$$

$$\begin{pmatrix} A_\mu \\ Z_\mu \end{pmatrix} = \begin{pmatrix} \cos \theta_w & \sin \theta_w \\ -\sin \theta_w & \cos \theta_w \end{pmatrix} \begin{pmatrix} B_\mu \\ W_\mu^0 \end{pmatrix}$$

$$\sin \theta_w = \frac{g_1}{\sqrt{g_1^2 + g_2^2}}, \quad \cos \theta_w = \frac{g_2}{\sqrt{g_1^2 + g_2^2}}, \quad e = \frac{g_1 g_2}{\sqrt{g_1^2 + g_2^2}}, \quad \sin^2 \theta_w = 0.23$$

2. Why we need Higgs in SM

2-1 Where is mass ?

If mass terms were added by hand.....

In U(1) gauge transformation, $A^\mu \rightarrow A^\mu + \frac{1}{e}\partial_\mu\alpha(x)$

the mass term $\frac{1}{2}m^2 A^\mu A_\mu$ is not invariant clearly.

In the case of a SU(2) fermion, current term is

$$\bar{\psi}\gamma^\mu\psi = \bar{\psi}_L\gamma^\mu\psi_L + \bar{\psi}_R\gamma^\mu\psi_R .$$

On the other hand the mass term is

$$m\bar{\psi}\psi = m(\bar{\psi}_R\psi_L + \bar{\psi}_L\psi_R)$$

which contains only doublet but not singlet,

therefore breaks SU(2) invariant Lagrangian.

We need a gauge invariant Lagrangian
which generates masses of fermions
and physical gauge bosons !!!

From hep-ph/9708417 20 Aug. 1997

Higgs Boson: Keyboards



Higgs Boson is a classically trained Jazz pianist with a taste for rock and pop music. His influences are far and wide, ranging from Debussy to Pink Floyd and Dave Grusin to Chick Corea.

Instruments:- Ensoniq VFX SD11 / Roland JV 1080 - 880 - U220 - JD 800 / Korg i3 - Wavestation AD / Alesis D4 / E111 Sampler / Akai S1000 / TG100 / Emu Proteus 1 / Juno 6
Instruments:- Ensoniq VFX SD11 / Roland JV 1080 - 880 - U220 - JD 800 / Korg i3 - Wavestation AD / Alesis D4 / E111 Sampler / Akai S1000 / TG100 / Emu Proteus 1 / Juno 6.

Email Higgs Boson:- hboson@enterprise.net

Back to Muscian list

2-2 Introduce Higgs Lagrangian

$$\mathcal{L}_{Higgs} = (D_\mu \phi)^\dagger (D^\mu \phi) - V(\phi)$$

$$\text{where } V(\phi) \equiv \mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2$$

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \quad \begin{aligned} \phi^+ &= \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2) \\ \phi^0 &= \frac{1}{\sqrt{2}}(\phi_3 + i\phi_4) \end{aligned}$$

$$\phi^\dagger \phi = \frac{1}{2}(\phi_1^2 + \phi_2^2 + \phi_3^2 + \phi_4^2)$$

In $-\mu^2 < 0$, find $\phi^\dagger \phi$ where $V(\phi)$ becomes minimum

$$\frac{\partial V}{\partial (\phi^\dagger \phi)} = \mu^2 + 2\lambda \phi^\dagger \phi = 0, \quad \phi_0^\dagger \phi_0 = -\frac{\mu^2}{2\lambda}$$

Choose ϕ_0 so that it satisfies electric charge conservation $\phi_1 = \phi_2 = \phi_3 = 0, \phi_4 = v$

$$\phi_0 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}, \quad \text{then } \phi_0^\dagger \phi_0 = \frac{v^2}{2} = -\frac{\mu^2}{2\lambda}$$

Expand around ϕ_0 so that Higgs field, $H(x)$ and weak gauge bosons can have correct mass terms

$$\phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H(x) \end{pmatrix}$$

2-3 Generate weak boson mass

Out of $(D_\mu \phi)^\dagger (D^\mu \phi) - V(\phi)$, relevant terms are

$$\phi^\dagger \left(i g_1 \frac{Y}{2} B_\mu + i g_2 \frac{\tau^i}{2} W_\mu^i \right)^\dagger \left(i g_1 \frac{Y}{2} B_\mu + i g_2 \frac{\tau^i}{2} W_\mu^i \right) \phi$$

$$= \left(\frac{1}{2} v g_2 \right)^2 W_\mu^+ W^{\mu-} + \frac{v^2}{8} (g_1^2 + g_2^2) Z_\mu Z^\mu$$

$$+ \frac{g_2^2 v}{2} H W_\mu^+ W^{\mu-} + \frac{1}{4} (g_1^2 + g_2^2) v H Z_\mu Z^\mu$$

$$+ \frac{g_2^2}{4} H H W_\mu^+ W^{\mu-} + \frac{1}{8} (g_1^2 + g_2^2) H H Z_\mu Z^\mu$$

$$= M_W^2 W_\mu^+ W^{\mu-} + M_Z^2 Z_\mu Z^\mu$$

$$+ g_2 M_W H W_\mu^+ W^{\mu-} + \frac{g_2 M_Z^2}{2 M_W} H Z_\mu Z^\mu$$

$$+ \frac{g_2^2}{4} H H W_\mu^+ W^{\mu-} + \frac{g_2^2}{8 \cos^2 \theta_w} H H Z_\mu Z^\mu$$

$$HWW > HZZ \gg HHWW > HHZZ$$

2-4 Generate fermion mass

Mass of lepton

$$\begin{aligned}
 \mathcal{L}_{Lep} &= g_e(\bar{\mathbf{L}}\phi e_R + \bar{e}_R\phi^\dagger \mathbf{L}) \\
 &= \frac{g_e v}{\sqrt{2}}(\bar{e}_L e_R + \bar{e}_R e_L) + \frac{g_e}{\sqrt{2}}(\bar{e}_L e_R + \bar{e}_R e_L)H \\
 &= m_e \bar{e}e + \frac{m_e}{v} \bar{e}eH
 \end{aligned}$$

Mass of quark

u_R must have mass, thus we need another Higgs doublet, ϕ_c which is isospin conjugate of ϕ

$$\phi_c = e^{-i\pi\tau^2/2}\phi^\star = -i\tau^2\phi^\star = \frac{1}{\sqrt{2}} \begin{pmatrix} -\phi^{0\star} \\ \phi^- \end{pmatrix} \rightarrow \frac{1}{\sqrt{2}} \begin{pmatrix} v + H(x) \\ 0 \end{pmatrix}$$

$$\begin{aligned}
 \mathcal{L}_Q &= g_d \bar{\mathbf{Q}}_L \phi d_R + g_u \bar{\mathbf{Q}}_L \phi_c u_R + g_d \bar{d}_R \phi^\dagger \mathbf{Q}_L + g_u \bar{u}_R \phi_c^\dagger \mathbf{Q}_L \\
 &= \frac{g_d}{\sqrt{2}}(v + H)(\bar{d}_L d_R + \bar{d}_R d_L) + \frac{g_u}{\sqrt{2}}(v + H)(\bar{u}_L u_R + \bar{u}_R u_L) \\
 &= \frac{g_d v}{\sqrt{2}} \bar{d}d + \frac{g_d}{\sqrt{2}} \bar{d}dH + \frac{g_u v}{\sqrt{2}} \bar{u}u + \frac{g_u}{\sqrt{2}} \bar{u}uH \\
 &= m_d \bar{d}d + \frac{m_d}{v} \bar{d}dH + m_u \bar{u}u + \frac{m_u}{v} \bar{u}uH
 \end{aligned}$$

2-5 Summary of SM and Higgs

$$\mathcal{L}_{SM} = \mathcal{L}_{Ferm} + \mathcal{L}_{Higgs} + \mathcal{L}_{Lep} + \mathcal{L}_Q - \frac{1}{4}W_{\mu\nu}W^{\mu\nu} - \frac{1}{4}B_{\mu\nu}B^{\mu\nu}$$

Scenario toward Higgs

- To guarantee the gauge invariance, the covariant derivatives must be defined which contains gauge bosons consequently.
- We constructed Lagrangians for the SM fermions requiring the gauge invariance.
- The mass-less fermion Lagrangian can fit the physical particles, if theoretical fields are interpreted correctly.
- However, if one adds masses to physical particles by hand, it breaks the gauge invariance.
- We need a gauge invariant potential so that it can give masses to fermions and gauge bosons by breaking symmetry. That is Higgs!!!.

What SM can not predict

- The number of generations
- Coupling strength
- Masses of fermions
- Mass of Higgs

3. Properties of SM Higgs

3-1 Decay of SM Higgs

Points !!!

$$g_f = \frac{g_2}{2M_W} M_f$$

$$\Gamma_{f\bar{f}} \propto m_f^2 M_H$$

$$\Gamma_{WW \text{ or } ZZ} \propto M_H^3$$

$$\Gamma_{gg} \propto \alpha_s^2 M_H^3$$

For low mass region $M_H < 130$ GeV

$$H \rightarrow b\bar{b}, c\bar{c}, \tau\bar{\tau}$$

$$H \rightarrow gg$$

$$H \rightarrow \gamma\gamma$$

$$H \rightarrow Z\gamma$$



For high mass region $M_H > 130$ GeV

$$H \rightarrow WW$$

$$H \rightarrow ZZ$$

$$H \rightarrow t\bar{t}$$

From hep-ph/9712334 10 Dec. 1997



3-2 Production of SM Higgs

At e^+e^- colliders

$$e^+e^- \rightarrow (Z) \rightarrow Z + H$$

$$e^+e^- \rightarrow \nu\bar{\nu}(WW) \rightarrow \nu\bar{\nu} + H$$

$$e^+e^- \rightarrow e^+e^-(ZZ) \rightarrow e^+e^- + H$$

$$e^+e^- \rightarrow (\gamma, Z) \rightarrow t\bar{t} + H$$

At pp colliders



$$gg \rightarrow H$$

$$WW/ZZ \rightarrow H$$

$$q\bar{q} \rightarrow W/Z + H$$

$$gg, q\bar{q} \rightarrow t\bar{t} + H$$

From hep-ph/9710439 22 Oct. 1997




3-3 Mass bound on SM Higgs

$$V(\phi) = \mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2$$

$$v = \sqrt{-\mu^2/\lambda}, \quad \mu^2 < 0$$

- Upper bound can be constrained by requiring that perturbative theory is valid up to the GUT scale.
- Lower bound can be determined by requiring $\lambda > 0$ up to the GUT scale which corresponds to the requirement of vacuum stability.

From hep-ph/9708416 20 Aug. 1997



If $160 < M_H < 170$ GeV, the perturbative treatment is valid up to GUT scale, $\Lambda \sim 10^{19}$ GeV.