



Hadron distribution inside jets for hadronization and spin dynamics

Zhongbo Kang
UCLA

RBRC Workshop: Synergies of pp and pA
Collisions with an EIC
June 26 - 28, 2017

Jets as a precision probe of QCD



- Extraction of strong coupling constant



- PDFs are constrained by collider jet data, especially gluon distribution functions

Jets as a precision probe of QCD

- Extraction of strong coupling constant



- PDFs are crucial for the distribution of the process

process

W asymmetry
W and Z production (differential)
W+c production
Drell-Yan (DY): high invariant mass
Drell-Yan (DY): low invariant mass
W,Z +jets
Inclusive jet and di-jet production

sensitivity to PDFs

→ quark flavour separation
→ valence quarks
→ strange quark
→ sea quarks, high-x
→ low-x
→ gluon medium-x
→ gluon and $\alpha_s(M_Z)$
→ gluon medium, high-x
→ gluon and $\alpha_s(M_Z)$

Direct photon

ttbar, single top

Hadron distribution inside the jet

- Study a hadron distribution inside a fully reconstructed jet

$$p + p \rightarrow \text{jet} (h) + X$$


$$F(z_h; p_T) = \frac{d\sigma^h}{dp_T d\eta dz_h} \bigg/ \frac{d\sigma}{dp_T d\eta}$$

$$F(z_h, j_\perp; p_T) = \frac{d\sigma^h}{dp_T d\eta dz_h d^2 j_\perp} \bigg/ \frac{d\sigma}{dp_T d\eta}$$

$$z_h = p_T^h / p_T^{\text{jet}}$$

$j_\perp$: hadron transverse momentum with respect to the jet direction


- The 1st observable is like collinear fragmentation function, while the 2nd observable is more like a TMD fragmentation function
- Both RHIC and LHC performed measurements for this observable



Lots of data: inclusive jet cross section


- Single inclusive jet cross sections

$$p + p \rightarrow \text{jet} + X$$






Lots of data: jet fragmentation function

- Hadron distribution inside a jet $p + p \rightarrow \text{jet}(h) + X$




Light charged hadrons






Lots of data: jet fragmentation function

- Hadron distribution inside a jet $p + p \rightarrow \text{jet}(h) + X$



Light charged hadrons



Theoretical tools: QCD factorization


- Single inclusive jet production: $p + p \rightarrow \text{jet} + X$

$$\frac{d\sigma^{pp \rightarrow \text{jet} X}}{dp_T d\eta} = \sum_{a,b,c} f_a \otimes f_b \otimes H_{ab}$$



partonic hard-scattering cross section

$$H_{ab} = \alpha_s^2 \left(H_{ab}^{(0)} + \alpha_s H_{ab}^{(1)} + \alpha_s^2 H_{ab}^{(2)} + \dots \right)$$




- The idea is simple: dynamics which happen in very different scales do not interfere with each other: **Λ_{QCD} vs P_T**

Theoretical tools: QCD factorization


- Single inclusive jet production: $p + p \rightarrow \text{jet} + X$

$$\frac{d\sigma^{pp \rightarrow \text{jet} X}}{dp_T d\eta} = \sum_{a,b,c} f_a \otimes f_b \otimes H_{ab}$$



partonic hard-scattering cross section


- The scal



NLO 1990

Ellis, Kunszt, Soper '90


$\sim \mathcal{O}(\alpha_s^2)$ (1) $\sim \mathcal{O}(\alpha_s^2)$ (2) $+ \dots$



NNLO 2016 ...

Currie, Glover, Pires '16


Most recent jet measurements



Jet radius R is small: 0.2 – 0.4, $[\alpha_s \ln(R)]^n$ resummation?

QCD factorization makes things simple

- When $R \ll 1$, the relevant scales for single jet production
 - Two momenta: (1) hard collision: p_T (2) jet radius can build one: $p_T R$
 - A further factorization




$$\frac{d\sigma^{pp \rightarrow \text{jet } X}}{dp_T d\eta} = \sum_{a,b,c} f_a \otimes f_b \otimes H_{ab \rightarrow c} \otimes J_c(z, \mu \sim p_T R)$$

- Good thing: semi-inclusive jet function $J_{q,g}(z, R)$ are purely perturbative

Kang, Ringer, Vitev, arXiv:1606.06732, Dai, Kim, Leibovich, 1606.07411,
see also, Kaufmann, Mukherjee, Vogelsang, 1506.01415

Semi-inclusive jet functions

- NLO quark jet



$$\begin{aligned}
 J_q(z, \omega_J) = & \delta(1-z) + \frac{\alpha_s}{2\pi} \left(\frac{1}{\epsilon} + L \right) \left[P_{qq}(z) + P_{gq}(z) \right] \\
 & - \frac{\alpha_s}{2\pi} \left\{ C_F \left[2(1+z^2) \left(\frac{\ln(1-z)}{1-z} \right)_+ + (1-z) \right] - \delta(1-z) d_J^{q, \text{alg}} \right. \\
 & \left. + P_{gq}(z) 2 \ln(1-z) + C_F z \right\},
 \end{aligned}$$

- Evolution equation of semi-inclusive jet functions

After renormalization, one will find: semi-inclusive quark/gluon jets follow DGLAP evolution equation, just like hadron fragmentation functions


$$\mu \frac{d}{d\mu} J_i(z, \omega_J, \mu) = \frac{\alpha_s(\mu)}{\pi} \sum_j \int_z^1 \frac{dz'}{z'} P_{ji} \left(\frac{z}{z'}, \mu \right) J_j(z', \omega_J, \mu)$$

Ln(R) resummation



- Natural scale for jet dynamics

$$L = \ln \frac{\mu^2}{\omega_J^2 \tan^2 \frac{\mathcal{R}}{2}} \rightarrow \mu = \omega_J \tan \frac{\mathcal{R}}{2} = p_T \cdot R$$

- Jet radius resummation: $(\alpha_s \ln R)^n$




Effect of $\ln(R)$ resummation



Other properties of the new formalism

- Unified factorization formalism for hadron and jet production




$$\frac{d\sigma^{pp \rightarrow hX}}{dp_T d\eta} = \sum_{a,b,c} f_a \otimes f_b \otimes H_{ab \rightarrow c} \otimes D_c^h$$

$$\frac{d\sigma^{pp \rightarrow \text{jet}X}}{dp_T d\eta} = \sum_{a,b,c} f_a \otimes f_b \otimes H_{ab \rightarrow c} \otimes J_c(\mu \sim p_T R)$$

- Consistent definition of what are called quark/gluon jets: relative ratio
- Even though derived for small R, $R = 0.7$, the difference between small R approximation and full result is less than 5%

Jet fragmentation function

- First produce a jet, and then further look for a hadron inside the jet



$$F(z_h, p_T) = \frac{d\sigma^h}{dy dp_T dz_h} \bigg/ \frac{d\sigma}{dy dp_T}$$

$$z_h = p_T^h / p_T$$

$$z = p_T / p_T^c$$

Kang, Ringer, Vitev, arXiv:1606.07063, JHEP

- Just like the single inclusive jet production, we have
 - Semi-inclusive fragmenting jet function

$$\frac{d\sigma}{dy dp_T dz_h} \propto \sum_{a,b,c} \int \frac{dz}{z^2} \mathcal{G}_c^h(z, z_h, \mu) \int \frac{dx'}{x'} f_{a/p}(x') \int \frac{dx}{x} f_{b/A}(x) \hat{\sigma}_{ab \rightarrow c}$$

A blue arrow points from the text "Semi-inclusive fragmenting jet function" to the term $\mathcal{G}_c^h(z, z_h, \mu)$ in the equation.

Two DGLAPs


- Again DGLAP evolution: evolution is for variable z

$$\mu \frac{d}{d\mu} \mathcal{G}_i^h(z, z_h, \mu) = \frac{\alpha_s(\mu)}{\pi} \sum_j \int_z^1 \frac{dz'}{z'} P_{ji} \left(\frac{z}{z'} \right) \mathcal{G}_j^h(z', z_h, \mu)$$

- Relation to standard FFs: relevant to variable z_h

$$\mathcal{G}_i^h(z, z_h, \mu) = \sum_j \int_{z_h}^1 \frac{dz'_h}{z'_h} \mathcal{J}_{ij}(z, z'_h, \mu) D_j^h \left(\frac{z_h}{z'_h}, \mu \right)$$

Kang, Ringer, Vitev, arXiv:1606.07063





Resum $\ln(R)$

Evolve standard FFs from 1 GeV to $p_T \times R$

Some interesting phenomenology


- Works pretty well in comparison with experimental data



Kang, Ringer, Vitev, arXiv:1606.07063, JHEP

Further improve light-hadron FFs


- arXiv:1706.02859



- An ideal observable to constrain collinear FFs, at both small and large z region, *especially for gluon FFs*

D meson distribution inside jets

- Jet FFs for D meson: use FFs fitted from e+e- data




Kneesch, Kniehl, Kramer, Schienbein, 08

Using ZM-VFNS scheme:
Chien, Kang, Ringer, Vitev, Xing,
1512.06851, JHEP 16

D meson distribution inside jets

- Jet FFs for D meson: use FFs fitted from e+e- data



Kneesch, Kniehl, Kramer, Schienbein, 08

Using ZM-VFNS scheme:
Chien, Kang, Ringer, Vitev, Xing,
1512.06851, JHEP 16

$$- - - D_g^D(z, \mu) \rightarrow 2D_g^D(z, \mu)$$

New fit of D-meson FFs:
Anderle, Ringer, Stratmann, Vitev, in progress

A new global analysis of FFs


- New fit of D-meson FFs

New fit of D-meson FFs:
Anderle, Ringer, Stratmann, Vitev, in progress

Confirms


JHEP 1605 (2016) 125

Guess





Anderle, talk at QCD Evolution Workshop 2017

The new fit




With the e^+e^- and pp inclusive cross section

- The new fit does not spoil the comparison with inclusive data






With the e^+e^- and pp inclusive cross section

- The new fit does not spoil the comparison with inclusive data



Hadronization modification in p+A/e+A?


- For jet fragmentation functions, by construction/definition, the effects from the modification of PDFs in the target nucleus do NOT matter
 - Mostly cancel out in the ratio
- Thus any modification should be coming from the modification of hadronization process, *one of key observables at EIC white paper*



Nuclear modification of fragmentation function

- Earlier evidence for nuclear modification of hadronization process

HERMES, 0704.3270



$$R_A^H(\nu, Q^2, z, p_T^2) = \frac{\left(\frac{N^H(\nu, Q^2, z, p_T^2)}{N^e(\nu, Q^2)} \right)_A}{\left(\frac{N^H(\nu, Q^2, z, p_T^2)}{N^e(\nu, Q^2)} \right)_d}$$



- By construction, *the observable is insensitive to nuclear PDFs*

Two models describe the data

- Parton energy loss in cold nuclear matter leads to an effective change of the fragmentation function



$$\tilde{D}_{a \rightarrow h}(z) \approx \frac{1}{1 - \Delta z} D_{a \rightarrow h} \left(\frac{z}{1 - \Delta z} \right)$$

Chang, Deng, Wang, PRC 14
Deng, Wang, 10




Fragmentation function in nuclei

- Phenomenological model of Sassot, Stratmann, Zurita: global fitting of SIDIS as well as d+Au RHIC data




Sassot, Stratmann, Zurita, 0912.1311



P+Pb collisions at the LHC

- arXiv:1706.02859 – usually at smaller jet p_T , these effects should be more dramatic, *RHIC as an ideal place* – both light and heavy meson




What about TMD FFs?

- TMD hadron distribution inside the jet

$$F(z_h, j_\perp; p_T) = \frac{d\sigma^h}{dp_T d\eta dz_h d^2 j_\perp} \bigg/ \frac{d\sigma}{dp_T d\eta}$$

$$z_h = p_T^h / p_T^{\text{jet}}$$

$j_\perp$: hadron transverse momentum with respect to the jet direction



- Factorization formalism

Kang, Liu, Ringer, Xing, 1705.08443

$$\frac{d\sigma}{dp_T d\eta dz_h d^2 j_\perp} \propto \sum_{a,b,c} f_a \otimes f_b \otimes H_{ab \rightarrow c} \otimes \mathcal{G}_c^h(z, z_h, \omega_J R, j_\perp, \mu)$$

- Re-factorization of semi-inclusive fragmenting jet function

$$\begin{aligned} \mathcal{G}_c^h(z, z_h, \omega_J R, \mathbf{j}_\perp, \mu) = & \mathcal{H}_{c \rightarrow i}(z, \omega_J R, \mu) \int d^2 \mathbf{k}_\perp d^2 \boldsymbol{\lambda}_\perp \delta^2(z_h \boldsymbol{\lambda}_\perp + \mathbf{k}_\perp - \mathbf{j}_\perp) \\ & \times D_{h/i}(z_h, \mathbf{k}_\perp, \mu, \nu) S_i(\boldsymbol{\lambda}_\perp, \mu, \nu R) \end{aligned}$$

TMD factorization for DY: $p + p \rightarrow [\gamma^* \rightarrow \ell^+ \ell^-] + X$

- Factorized form and mimic “parton model”

$$\frac{d\sigma}{dQ^2 dy d^2q_\perp} \propto \int d^2k_{1\perp} d^2k_{2\perp} d^2\lambda_\perp H(Q) f(x_1, k_{1\perp}) f(x_2, k_{2\perp}) S(\lambda_\perp) \delta^2(k_{1\perp} + k_{2\perp} + \lambda_\perp - q_\perp)$$

$$= \int \frac{d^2b}{(2\pi)^2} e^{iq_\perp \cdot b} H(Q) f(x_1, b) f(x_2, b) S(b)$$




$$F(x, b) = f(x, b) \sqrt{S(b)}$$

$$= \int \frac{d^2b}{(2\pi)^2} e^{iq_\perp \cdot b} H(Q) F(x_1, b) F(x_2, b)$$

- Rapidity divergences cancel between

$$F(x, b) = f(x, b) \sqrt{S(b)}$$




What's different for hadron in the jet?

- Soft radiation has to happen inside the jet
 - For single inclusive jet production, first we produce a high-pt jet
 - This process only involves hard-collinear factorization, and such a process is not sensitive to any soft radiation
 - This is the usual standard “collinear factorization”

$$\int_0^\infty \frac{dy}{y} \Rightarrow \int_0^{\tan^2 \frac{R}{2}} \frac{dy}{y}$$

$$y \sim \frac{\ell^+}{\ell^-}$$




- Once such a high-pt jet is produced, we further observe a hadron inside the jet
 - At this step, we measure the relative transverse momentum of hadron w.r.t the jet. For such a step, soft radiation matters
 - However, only those soft radiation that happens inside the jet matters
 - Restricts soft radiation to be within the jet: cuts half of the rapidity divergence
- Rapidity divergence cancel between restricted “soft factor” and TMD FFs
 - At least up to this order, the combined evolution is the same as the usual TMD evolution in SIDIS, DY, e+e-; justify the use of same TMD evolution here

$$\sqrt{S(b)} D_c^h(z_h, b)_{e^+e^-} \Rightarrow S(b, R) D_c^h(z_h, b)_{pp}$$

TMD + DGLAP evolution



- Evolution structure



- TMD FFs thus are related to the usual TMD FFs in SIDIS at scale $p_T \times R$
- Thus hadron TMD distribution inside the jet could be used to test the universality of TMD FFs from SIDIS, e^+e^- processes


Hadron TMD distribution inside jets

- Unpolarized p+p collisions: very sensitive to gluon TMDs
- If we want to be able to compare gluon TMDs in p+p and e+p, then p+p measurements are essentially necessary





Collins azimuthal asymmetry

- Transversely polarized p+p collisions



Kang, Prokudin, Ringer, Yuan, 1706.xxxx, to appear

- Test universality of Collins function between e+p, e+e, and p+p
- Test TMD evolution



Summary

- jet cross section and jet substructure for inclusive jet production follow a two-step factorization
 - First step: parton-to-jet production
 - Second step: jet internal substructure
- The hard function associated with the 1st step is the same as that for single inclusive hadron production: DGLAP evolution
- For jet substructure, one could then concentrate on the 2nd step
- Collinear and TMD distribution of hadron in a jet are great processes to probe collinear and/or TMD FFs
 - Great opportunities to study spin-independent and spin-dependent hadronization/fragmentation effects in both vacuum and cold nuclear medium
 - New places to test TMD evolution effects

Thank you!