


Probing Sea Quark TMD with Drell-Yan at SpinQuest/E0139 at Fermilab

Ming Liu


Los Alamos National Laboratory
For the SpinQuest/E1039 Collaboration

QCD Evolution 2019, May 13-17
Argonne National Laboratory




Outline

- SpinQuest/E1039 experiment at Fermilab
 - E1039 polarized NH_3/ND_3 targets
 - Follow up of SeaQuest/E906 unpolarized target program
- Novel physics of sea quarks at $x = 0.1 - 0.4$
 - Flavor asymmetry
 - Sivers & OAM
- Other opportunities
 - Parasitic E1067 dark photon search, 2016 - 2021+
 - Future E1027 polarized beam possibility, 2021+
 - TMD physics complementary to the future EIC program




A series of follow up experiments:
SeaQuest, SpinQuest, DarkQuest ...




Fermilab High Intensity Frontier

Fixed Target
Beam lines




Tevatron
800 GeV

Main
Injector
120 GeV



SpinQuest Dimuon Spectrometer

Year 2015



120 GeV protons from the Main Injector

- 4s beam spill very 60 sec
- 19ns RF, ~10s K protons per RF bucket
- 5×10^{12} Proton On Target (POT) per spill
- Total integrated POT for E1039 (2-year): 1.4×10^{18} POT

E906 unpolarized targets: 2012-2017

- ^1H , ^2D , ^{12}C , ^{56}Fe , ^{184}W


E1039 polarized targets: 2019 – 2021+

- Polarized protons (NH_3)
- Polarized neutrons (ND_3)


E1027 polarized beam

5/13/19

SeaQuest Dimuon Tracking Top View




Drell-Yan @SeaQuest – a Sea Quark Laboratory




$$\frac{d^2\sigma}{dx_t dx_b} = \frac{4\pi\alpha^2}{9x_1 x_2 s} \sum e^2 [q_b(x_b) \bar{q}_t(x_t) + \bar{q}_b(x_b) \cancel{q(x_t)}]$$

$$\approx \frac{4\pi\alpha^2}{9x_1 x_2 s} \sum e^2 [q_b(x_b) \bar{q}_t(x_t)]$$

Fixed target kinematics favors sea-quarks from target – **a sea quark lab!**




Dimuon Mass from SeaQuest/E906



E906 Unpolarized Physics Program

- Thin targets: $\sim 10\%$ interaction length
 - Liquid H/D
 - Solid C, Fe, W
- Physics
 - Sea quark flavor asymmetry, \bar{d}/u
 - Quark energy loss in p+A collisions, dE/dx
 - TMD and more ...
- Experimental runs – 6 years
 - 2012 – commissioning
 - 2017 – completed




Flavor Asymmetry of Sea Quarks at Intermediate x

Proton vs “Neutron” targets:

$$\left. \frac{\sigma^{pd \rightarrow \mu^+ \mu^-}}{\sigma^{pp \rightarrow \mu^+ \mu^-}} \right|_{x_b \gg x_t} \approx \frac{1}{2} \left[1 + \frac{\bar{d}(x_t)}{\bar{u}(x_t)} \right]$$

This could lead to a very interesting physics ...


J. C. Peng et al (FNAL E866/NuSea), PRD 58, 092004 (1998)




Sea Quark Flavor Asymmetry and OAM

Pion cloud model


- Sea-quark flavor asymmetry
- Sea-quark orbital angular motion
- Expect large Siverts function at $x = 0.1 - 0.4$




Pion cloud and Drell-Yan process



$$|p\rangle = a_1 |p_0\rangle + a_2 |p_0 + \pi^0\rangle + a_3 |n + \pi^+\rangle + \dots$$







Nucleon 3-D Structure and Sivers Function

Sivers function $f_{1T}^\perp(x, k_T)$

$$f_{1T}^\perp = \begin{array}{c} \uparrow \\ \bigcirc \end{array} - \begin{array}{c} \bigcirc \\ \downarrow \end{array}$$




Sivers Functions from Global Fits

- Sea Quark Sivers poorly constrained, SIDIS not sensitive to sea quarks at large x

1612.06413, M. Anselmino et al

Sea quarks



RHIC pp500GeV: $W^{+/-} A_N$

$$A_N(W^+) \sim \left(\Delta^N f_{u/p^\uparrow} \otimes f_{\bar{d}/p} + \Delta^N f_{\bar{d}/p^\uparrow} \otimes f_{u/p} \right)$$

$$A_N(W^-) \sim \left(\Delta^N f_{\bar{u}/p^\uparrow} \otimes f_{d/p} + \Delta^N f_{d/p^\uparrow} \otimes f_{\bar{u}/p} \right)$$


RHIC data:

- A mix of valence and sea quark Sivers
- Quark flavor identified
- High Q^2
- Statistically limited, $\sim O(10\%)$
- **Hints of sizable sea-quark Sivers contributions**


E1039:

- low Q^2
- high statistics, $\sim O(1\%)$


Anselmino et al 2016



F. Tian et al. / Nuclear Physics A 968 (2017) 379–390



(b) $A_N^{W^+}$




(a) $A_N^{W^-}$

Polarized NH_3 Target Developed for DY Sivers

Microwave: Induces electron spin flips


- EIO + Power equip:

Refrigerator:



Target:


material: frozen NH₃/ND₃. Irradiation@ NIST



Superconducting
Coils for Magnet: 5T



Dynamic Nuclear Polarization: Pol. ~90%



With DNP,
Pol. ~ 90%

W/o DNP, at thermal equilibrium:

- $T = 1\text{K}$
- $B = 5\text{T}$

Proton target polarization:

$$P_i = 0.5\%$$

$$P_i = \tanh \left(\frac{g_i \mu_i B}{2k_B T} \right)$$

Projected SpinQuest Target and Beam Performance


$$A_{\text{meas}} = f \cdot P_T \cdot A_{\text{phy}}$$

Target		Beam	
Polarization P	88%	Beam	10^{13} p per spill
Packing fraction	.6	spill	5 sec , one per minute
Dilution Factor f	.176	Luminosity	$4 \cdot 10^{35} / \text{cm}^2 / \text{s}$
Density NH_3	$.82 \text{ g/cm}^3$	E_{Beam}	120 GeV
		Total $\mu^+ \mu^-$ pairs	$4.59 \cdot 10^5$
		Experiment available	.48

Projected Drell-Yan Transverse Single Spin Asymmetry

$$A_N^{DY} \propto \frac{u(x_b) \cdot f_{1T}^{\perp, \bar{u}}(x_t)}{u(x_b) \cdot \bar{u}(x_t)}$$



$$\delta A = \frac{1}{f} \frac{1}{P} \frac{1}{\sqrt{N^+ + N^-}}$$





Range x_B	Mean x_B	Total events	ΔA
0.10–0.14	0.123	159097	0.016
0.14–0.17	0.154	136558	0.017
0.17–0.21	0.188	123566	0.018
0.21–0.50	0.258	119508	0.019

Drell-Yan Sivers Asymmetries w/ QCD Evolution

Kang et al, 1401.5078




Sea-quark
Sivers



E1039 Status & Plan

- DOE approval, March 2018
- E906 decommissioned 6/2018
- Polarized target to be installed by fall of 2019
 - E1039 commissioning starts in late 2019
 - Run for 2+ years, 2019-2021+



Fermilab Long Range Plan

Fermilab Program Planning 5-April-18

LONG-RANGE PLAN

		FY18	FY19	FY20	FY21	FY22	FY23	FY24	FY25	FY26	FY27	FY28	FY29	FY30	
LBNF / PIP II	SANFORD FNAL				DUNE	DUNE	DUNE	DUNE	DUNE	DUNE	DUNE	DUNE	DUNE	DUNE	
						LBNF	LBNF	LBNF	LBNF	LBNF	LBNF	LBNF	LBNF	LBNF	
NuMI	MI	MINERv	MINERv	OPEN	OPEN	OPEN	OPEN	OPEN							
		NOvA	NOvA	NOvA	NOvA	NOvA	NOvA	NOvA							
BNB	B	MicroBooNE	MicroBooNE	MicroBooNE	OPEN	OPEN	OPEN	OPEN			OPEN	OPEN	OPEN	OPEN	
		CARUS	CARUS	CARUS	CARUS	CARUS	CARUS	CARUS			OPEN	OPEN	OPEN	OPEN	
		SBND	SBND	SBND	SBND	SBND	SBND	SBND			OPEN	OPEN	OPEN	OPEN	
Muon Complex		g-2	g-2	g-2											
		Mu2e	Mu2e	Mu2e	Mu2e	Mu2e	Mu2e	Mu2e			Mu2e	Mu2e	Mu2e	OPEN	
SY 120	MT	FTBF	FTBF	FTBF	FTBF	FTBF	FTBF	FTBF			FTBF	FTBF	FTBF	FTBF	
	MC	FTBF	FTBF	FTBF	FTBF	FTBF	FTBF	FTBF			FTBF	FTBF	FTBF	FTBF	
	NM4	OPEN	E1039	E1039	E1039	E1039	OPEN	OPEN			OPEN	OPEN	OPEN	OPEN	
		FY18	FY19	FY20	FY21	FY22	FY23	FY24	FY25	FY26	FY27	FY28	FY29	FY30	

	Construction / commissioning		Run		Subject to PAC review		Shutdown
	Capability ended		Capability unavailable				

- NOTES:
1. Mu2e estimates 4 year running starts mid-FY22 after 18 months commissioning. Assume, with contingency, 5.5 years data taking.
 2. DUNE: 1st 10kT detector module commissioned in FY24. Runs without beam FY25 to mid-FY26.
 3. NOvA runs as long as possible [in the spirit of PAC Nov 2017].
 4. Assume NuMI in nubar mode through FY19 - facilitates 12E20 POT for MINERvA [PAC Nov 2017]. Assumption may need re-evaluation.
 5. Assume g-2 completed before Mu2e commissioning start mid-FY20. Very tight. Needs scrutiny.
 6. Assume E1039 fully approved & commissioned by mid-FY19.
Experiment estimates 2 yrs run. Add 1 yr contingency. [Stage 1 approval PAC June 2013, update July 2017]
 7. FY19 and FY20 MicroBooNE running subject to future PAC review [PAC July 2017].

Opportunity for new programs at SeaQuest after E1039



Physics Beyond E1039 Polarized DY A_N

- Dark sector physics search – DarkQuest

- Parasitic run with E1039: 2019 – 2021+
- A new proposal for a dedicated run after E1039

- Physics with polarized beams – E1027


- Polarize the Main Injector 120GeV beam
- Valence quark Sivers
 - Test QCD dynamics in DY vs DIS
- TMD physics



Dark Photons and Dark Higgs Search at SpinQuest/DarkQuest


Photon portal: “vector”

$$\mathcal{L}_{\text{mix}} = \frac{\epsilon}{2} F_{\mu\nu}^{\text{QED}} F_{\text{Dark}}^{\mu\nu}$$



Higgs portal: “scalar”

$$\mathcal{L}_{\text{mix}} = \mu\phi|H^\dagger H|$$




Advantage of hadronic collisions

Dark Sector Physics Search at DarkQuest

2019 ~ 2021: parasitic run for DM search



2021+: a proposal for a long term DM program after E1039; parasitic TMD physics?



Add tracking detectors
close to "target" to
improve mass resolution

Add EMCal, PID
 $e^{+/-}$, $h^{+/-}$, $\pi^{+/-}$

Dark Photon Search at DarkQuest with Future Projections





Spin physics Program with Polarized Main Injector – E1027

- Access both polarized valence and sea quarks
- Fermilab PAC stage-1 approved
- Complementary to the future EIC TMD Physics

Test QCD processes in DY vs DIS over a broad range of kinematics

$$A_N \equiv \frac{N^\uparrow - N^\downarrow}{N^\uparrow + N^\downarrow} \propto \frac{f_{1T}^{\perp,u}(x_B) \cdot \bar{u}(x_T)}{u(x_B) \cdot \bar{u}(x_T)}$$



TMDs probed via DY at Fermilab

Boer-Mulders functions:

- Unpolarized Drell-Yan: $d\sigma_{DY} \propto h_1^\perp \bar{h}_1^\perp \cos(2\phi)$

E906, E1039, E1027

Sivers functions:

- Single transverse spin asymmetry in polarized Drell-Yan:

$$A_N^{DY} \propto f_{1T}^\perp(x_q) f_{\bar{q}}^\perp(x_{\bar{q}})$$

E1039, E1027

Transversity distributions:

- Double transverse spin asymmetry in polarized Drell-Yan:

$$A_{TT}^{DY} \propto h_1(x_q) h_1(x_{\bar{q}})$$

E1027





- Drell-Yan and SIDIS involve different combinations of TMDs
- Drell-Yan does not require knowledge of the fragmentation functions
- T-odd TMDs are predicted to change sign from DIS to DY

(Boer-Mulders and Sivers functions)

Remains to be tested experimentally! → COMPASS, RHIC, EIC/SpinQuest for sea quarks

Summary and Outlook

Experiments	Run Time	Collision Types	Physics
E906	2012-2017	p + targets (H, D, C, Fe, W)	- dbar/ubar asymmetry - quark dE/dx
E1039	2018 – 2021+	p + pol. targets (NH₃, ND₃)	Sea-quark Sivers, TMDs
E1067(para.)	2017-2021+(para.)	p + any targets	dark photon, dark Higgs, ALP ...
DarkQuest	2021+ (dedicated)		
E1027	202x	Pol. p-beam +	- quark Sivers - TMD, spin



SpinQuest/E1039 Collaboration

*A relatively small collaboration,
great opportunities for new
comers to contribute and lead
major detector and physics
efforts*

Collaboration

INSTITUTION	FULL MEMBERS	AFFILIATE MEMBERS
Abilene Christian University	Donald Isenhower (PI), Michael Daugherty, Shon Watson	Haley Stien, John Marsden, Mitchell Schneller, Nathan Rowlands, Roy Salinas, Rusty Towell, Shannon McNease, Yves Ngenzi, Thomas Fitch
Argonne National Laboratory	Paul Reimer (PI), Donald Geesaman	Kawtar Hafidi, Kevin Bailey, Thomas O'Connor, Zhihong Ye, Benjamin Zeidman
Fermi National Accelerator Laboratory	Richard Tesarek (PI), Carol Johnstone, Charles Brown	Robert Bushek, Dave Christian, Donald Mitchell, David Northacker, Mike Geelhoed, Kathy Graden, Bridget Iverson, Ivan Vitev, Jin-Yuan Wu, Maddie Schoell, Steven Timm, Yanqiu Yin
KEK	Shin'ya Sawada (PI)	Shigeru Ishimoto
Los Alamos National Laboratory	Kun Liu (SP), Mikhail Yurov, Chun-Min Jen, Ming Liu, Xuan Li, Walter Sondheim	Jan Boissevain, Melynda Brooks, Matt Durham, David Kleinjan, Sho Uemura, Cesar Da Silva, Patrick McGaughey, Andi Klein
Mississippi State University	Lamiaa El Fassi (PI)	Dipangkar Dutta
New Mexico State University	Stephen Pate (PI), Vassili Papavassiliou, Haiwang Yu, Forhad Hossain	
RIKEN	Yuji Goto (PI)	
Tokyo Institute of Technology	Kenichi Nakano (PI), Toshi-Aki Shibata	
University of Colorado, Boulder	Edward Kinney (PI)	
University of Illinois, Urbana-Champaign	Jen-Chieh Peng (PI), Yen-Chu Chen	Naomi Makins, Ching Him Leung, Daniel Jumper, Jason Dove, Mingyan Tian, Bryan Dannowitz, Randall McClellan, Shivangi Prasad
University of Michigan	Wolfgang Lorenzon (PI), Minjung Kim, Noah Wuerfel	Daniel Morton, Richard Raymond, Marshall Scott
University of New Hampshire	Karl Slifer (PI), David Ruth	Maurik Holtrop
University of Virginia	Dustin Keller (SP), Joshua Hoskins, Zulkaida Akbar, Carlos Ramirez	Donal Day, Donald Crabb, Jixie Zhang, Oscar Rondon, Liliet Diaz, Arthur Conover, Brandon Kriesten, Simonetta Liuti, Ellen Brown, Blaine Norum, Matthew Roberts
Yamagata University	Yoshiyuki Miyachi (PI), Genki Nukazuka	Takahiro Iwata, Norihiro Doshita

backup

SpinQuest Experimental Hall




Beam

Target area


F-Mag

K-Mag

Muon-ID



New Beam Collimator and Target



Target cross section: 18 x 28 mm²

Beam cross section:

Need be well contained within
4 sigma, required by $dR < 2 \times 10^{-4}$

$\text{sigX} = 18/2/4 = 2.2 \text{ mm}$

$\text{sigY} = 28/2/4 = 3.5 \text{ mm}$


Beam jitter: $dX=dY \sim 1\text{mm}$

1 sig = 0.68269

2 sig = 0.95450

3 sig = 0.99730

4 sig = 0.99994



E906 beam profile:

SigX = 4.0mm


SigY = 3.0mm

$$f(x, \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

Beam collimator




120GeV
beam




Target and Beam Dump Event Separation

target at upstream: $Z = -3.5\text{m}$



Projected Dark Sector Physics Search Sensitivity

Dark photon search



Dark Higgs search

