


# 40<sup>th</sup> Anniversary of QCD: High $p_T$ physics from ICHEP 1972 to ICHEP 1982 with applications to RHI collisions


M. J. Tannenbaum  
Brookhaven National Laboratory  
Upton, NY 11973 USA

Zimanyi School  
and  
Wigner Seminar  
Budapest, Hungary  
December 7, 2012




It all began at the 1968  
ICHEP in Vienna.  
Panofsky reported on  
the first DIS results  
from SLAC which  
Bjorken had clarified  
using scaling arguments

# From Panofsky ICHEP 1968-SLAC ep DIS



The old way, hard to understand  
 $\nu W_2(Q^2, \nu)$  vs energy loss  $\nu$



The new way, Bjorken Scaling  
 $\nu W_2(Q^2, \nu)$  scales vs  $\omega = \nu/Q^2$   
 i.e. collapses onto one curve

# Bjorken Scaling in Deeply Inelastic Scattering and the Parton Model---1968

♥ The discovery that the DIS structure function

$$F_2(Q^2, \nu) = F_2\left(\frac{Q^2}{\nu}\right) \quad (1)$$

“**SCALED**” i.e just depended on the ratio

$$x = \frac{Q^2}{2M\nu} \quad (2)$$

independently of  $Q^2$  ( $\sim 1/r^2$ )



♥ as originally suggested by **Bjorken** *Phys. Rev.* **179**, 1547 (1969)

♥ Led to the concept of a proton composed of point-like **partons**. *Phys. Rev.* **185**, 1975 (1969)

□ The probability for a parton to carry a fraction  $x$  of the proton's momentum is measured by  $F_2(x)$

$$\nu = \frac{Q^2}{2Mx}$$

# Typical High Energy Physics—1960's



$\Omega^-$  (SSS)

Associated Production  
(not the actual discovery)

BNL-Barnes, Samios *et al.*, PRL12, 204 (1964)

# Why were some people studying “high $p_T$ ” physics in the 1960’s?

---


# Why were some people studying “high $p_T$ ” physics in the 1960’s?

---

They were searching for the W boson.

# Why were some people studying “high $p_T$ ” physics in the 1960’s?

- The first opportunity to study weak interactions at high energy was provided by the development of neutrino beams at the new accelerators in the early 1960’s **CERN-SpS , BNL-AGS**.



*In Token of Our  
Appreciation for  
Your Contribution  
to the Neutrino  
Run September 1961-  
June 1962.*


*“The  
Neutrino  
Group”*

*Gordon Danby  
Jean-Marc Guillard  
Dino Goulianos  
Warner Hayes*

*Ben Lederman  
Nori Mitsuyoshi  
Mel Schwartz  
Jack Steinberger*



COLUMBIA (NEWS)



BNL

- However, it was soon recognized that the intermediate (weak) boson  $W^\pm$ , might be more favorably produced in nucleon-nucleon collisions.

# The 'Zichichi signature' for the W boson

Proc. 12<sup>th</sup> ICHEP, Dubna 1964

## MUON-PROTON ELASTIC SCATTERING AT HIGH MOMENTUM TRANSFERS \*

*R. Cool, A. Maschke*

Brookhaven National Laboratory, USA

*L. Lederman, M. Tannenbaum*

Columbia University, USA

*R. Ellsworth, A. Melissinos, J. Tinlot, T. Yamanouchi*

University of Rochester, USA

(Presented by J. TINLOT)

We have studied the elastic scattering of negative muons from liquid hydrogen at momentum transfers of 550 MeV/c to 1050 MeV/c ( $q^2 = 7$  to 26 fermi<sup>-2</sup>), using a detecting array of spark chambers and scintillation counters. The experiment was performed at the AGS accelerator of the Brookhaven National Laboratory, and the runs were divided into three stages, as shown below:

of the proton in an aluminum plate spark chamber. One also measures the directions of the recoil proton and the recoil muon. This is equivalent to measuring three independent angles, from which one can infer for each event the value of  $k$ , and still overdetermine the scattering event by two degrees of freedom. This redundancy is used to select true scattering events from the background of false events, such as random coincidences, inelastic  $\mu - p$

### ЭЛЕКТРОМАГНИТНЫЕ ВЗАИМОДЕЙСТВИЯ

It appears that, within the uncertainties of these preliminary results, the muon scattering cross section for momentum transfers of up to 1 GeV/c is correctly described by the Rosenbluth formula and the  $e-p$  form factors. It is still too early to attempt a more quantitative definition of the possible deviation from the electron predictions.

### ДИСКУССИЯ

A. Z i c h i c h i.

I would like to ask Dr. Tinlot what is the accuracy of the measured cross section at 1 GeV/c momentum transfer.

J. T i n l o t.

The accuracy of the highest point (statistical error only), at 1.05 GeV/c, is 25%; at 950 MeV/c, the error in the point is about 15%.

### DISCUSSION






# Searches for W boson in p-p collisions

- 1965-1969 Beam dump experiments at ANL-ZGS and BNL-AGS looking for “large angle” muons didn’t find any. [ZGS-Lamb, et al PRL **15**, 800 (1965), AGS-Burns, et al, ibid 830, AGS-Wanderer et al, PRL **23**, 729 (1969)]
- How do you know how many W should have been produced?
- Chilton, Saperstein, Shrauner [PR**148**, 1380 (1966)] emphasize the importance of the timelike form factor, which is solved by
- Y. Yamaguchi [Nuovo Cimento **43**, 193 (1966)] Timelike form factor can be found by measuring the number of lepton pairs  $e^+e^-$  or  $\mu^+\mu^-$  “massive virtual photons” of the same invariant mass; BUT the individual leptons from these electromagnetically produced pairs might mask the leptons from the  $W^\pm$ .
- This set off a spate of single and di-lepton experiments, notably the discovery by Lederman et al of “Drell-Yan” production at the BNL-AGS, E70 at FNAL and CCR at the CERN-ISR.


# AGS-1969-71 Discovery of 'Drell-Yan' and ??



$$q + \bar{q} \rightarrow \mu^+ + \mu^-$$

$$p + U \rightarrow \mu^+ \mu^- + X$$

$$\sqrt{s_{NN}} = 7.4 \text{ GeV}$$



VIEW LETTERS

4 JANUARY 1971





FIG. 2. Experimental cross section of Christenson *et al.*, Ref. 8.

long forgotten

Christenson, Lederman...PRL **25**, 1523 (1970)

'Theory' Altarelli, Brant Preparata PRL **26** 42 (1971)


# AGS-1969-71 Discovery of 'Drell-Yan' and ??



$$q + q \rightarrow \mu^+ + \mu^-$$

$$p + U \rightarrow \mu^+ \mu^- + X$$

$$\sqrt{s_{NN}} = 7.4 \text{ GeV}$$



This is why I  
NEVER plot  
theory curves  
on any of my  
data

VIEW LETTERS

4 JANUARY 1971




FIG. 2. Experimental cross section of Christenson *et al.*, Ref. 8.

long forgotten

Christenson, Lederman...PRL **25**, 1523 (1970)

'Theory' Altarelli, Brant Preparata PRL **26** 42 (1971)

# LML very excited in 1970: AGS-diu continuum + Bj scaling $\rightarrow$ W cross section at any $\sqrt{s}$

Proposal to The National Accelerator Laboratory

## E70-(F)NAL

"Study of Lepton Pairs from Proton-Nuclear Interactions;

Search for Intermediate Bosons and Lee-Wick Structure"

W. LEE, L.M. LEDERMAN, J. APPEL, Columbia University,

M. TANNENBAUM, Harvard University, L. READ, J. SCULLI,

T. WHITE, and T. YAMANOUCHI, National Accelerator


Laboratory.

### ABSTRACT

We propose to observe lepton pairs emerging from high energy proton-nuclear collisions. Large effective mass pairs probe the hadronic electromagnetic structure. The continuum mass spectrum will be measured and any resonant structures in the mass range up to  $\sim 28$  GeV will be detected with great sensitivity. The data provides a prediction, via Conserved Vector Current theory, for the production cross section for weak vector bosons and these are also sought in the mass range  $\sim 8$ -25 GeV. We also propose an initial photon-electron beam survey at high transverse momentum which is also a W-search with good sensitivity.

June 17, 1970 + addendum Dec 1970  $\rightarrow$

Correspondent: L. M. Lederman, Columbia University



# LML very excited in 1970: AGS-diμ continuum +Bj scaling → W cross section at any √s

Proposal to The National Accelerator Laboratory

## E70-(F)NAL

"Study of Lepton Pairs from Proton-Nuclear Interactions;

Search for Intermediate Bosons and Lee-Wick Structure"

W. LEE, L.M. LEDERMAN, J. APPEL, Columbia University,

M. TANNENBAUM, Harvard University, L. READ, J. SCULLI,

T. WHITE, and T. YAMANOUCHI, National Accelerator

Laboratory.


### ABSTRACT

We propose to observe lepton pairs emerging from high energy proton-nuclear collisions. Large effective mass pairs probe the hadronic electromagnetic structure. The continuum mass spectrum will be measured and any resonant structures in the mass range up to  $\sim 28$  GeV will be detected with great sensitivity. The data provides a prediction, via Conserved Vector Current theory, for the production cross section for weak vector bosons and these are also sought in the mass range  $\sim 8-25$  GeV. We also propose an initial photon-electron beam survey at high transverse momentum which is also a W-search with good sensitivity.


June 17, 1970

+ addendum Dec 1970 →

Correspondent: L. M. Lederman, Columbia University



# ?? explained by $J/\psi$ in 1974 at AGS + SLAC



VOLUME 33, NUMBER 23

PHYSICAL REVIEW LETTERS

2 DECEMBER 1974

## Experimental Observation of a Heavy Particle $J^\dagger$

J. J. Aubert, U. Becker, P. J. Biggs, J. Burger, M. Chen, G. Everhart, P. Goldhagen, J. Leong, T. McCorriston, T. G. Rhoades, M. Rohde, Samuel C. C. Ting, and Sau Lan Wu  
*Laboratory for Nuclear Science and Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

and

Y. Y. Lee

*Brookhaven National Laboratory, Upton, New York 11973*

(Received 12 November 1974)

We report the observation of a heavy particle  $J$ , with mass  $m = 3.1$  GeV and width approximately zero. The observation was made from the reaction  $p + \text{Be} \rightarrow e^+ + e^- + x$  by measuring the  $e^+e^-$  mass spectrum with a precise pair spectrometer at the Brookhaven National Laboratory's 30-GeV alternating-gradient synchrotron.

This experiment is part of a large program to study the behavior of timelike photons in  $p + p \rightarrow e^+ + e^- + x$  reactions<sup>1</sup> and to search for new particles which decay into  $e^+e^-$  and  $\mu^+\mu^-$  pairs.

daily with a thin Al foil. The beam spot size is  $3 \times 6$  mm<sup>2</sup>, and is monitored with closed-circuit television. Figure 1(a) shows the simplified side view of one arm of the spectrometer. The two

## Discovery of a Narrow Resonance in $e^+e^-$ Annihilation\*

J.-E. Augustin,<sup>†</sup> A. M. Boyarski, M. Breidenbach, F. Bulos, J. T. Dakin, G. J. Feldman, G. E. Fischer, D. Fryberger, G. Hanson, B. Jean-Marie,<sup>†</sup> R. R. Larsen, V. Lüth, H. L. Lynch, D. Lyon, C. C. Morehouse, J. M. Paterson, M. L. Perl, B. Richter, P. Rapidis, R. F. Schwitters, W. M. Tanenbaum, and F. Vannucci<sup>‡</sup>

*Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305*

and

G. S. Abrams, D. Briggs, W. Chinowsky, C. E. Friedberg, G. Goldhaber, R. J. Hollebeek, J. A. Kadyk, B. Lulu, F. Pierre,<sup>§</sup> G. H. Trilling, J. S. Whitaker, J. Wiss, and J. E. Zipse

*Lawrence Berkeley Laboratory and Department of Physics, University of California, Berkeley, California 94720*  
(Received 13 November 1974)

We have observed a very sharp peak in the cross section for  $e^+e^- \rightarrow \text{hadrons}$ ,  $e^+e^-$ , and possibly  $\mu^+\mu^-$  at a center-of-mass energy of  $3.105 \pm 0.003$  GeV. The upper limit to the full width at half-maximum is 1.3 MeV.

# Now, Back To High $p_T$ “Hard-Scattering”

# BBK 1971

S.M.Berman, J.D.Bjorken and J.B.Kogut, Phys. Rev. **D4**, 3388 (1971)

- BBK calculated for p+p collisions, the inclusive reaction

$$A+B \rightarrow C + X \quad \text{when particle } C \text{ has } p_T \gg 1 \text{ GeV}/c$$

- The charged partons of DIS **must scatter electromagnetically** “*which may be viewed as a lower bound on the real cross section at large  $p_T$ .*”





FIG. 1. Kinematics of lepton-nucleon scattering in the parton model.





# BBK 1971

S.M.Berman, J.D.Bjorken and J.B.Kogut, Phys. Rev. **D4**, 3388 (1971)

- BBK calculated for p+p collisions, the inclusive reaction



$$A+B \rightarrow C + X \quad \text{when particle } C \text{ has } p_T \gg 1 \text{ GeV}/c$$

- The charged partons of DIS **must scatter electromagnetically** “*which may be viewed as a lower bound on the real cross section at large  $p_T$ .*”



# CCR discovery 1972

# Bj-prediction 1971



CCR, R. Cool, ICHEP 1972

Bjorken-International Lepton-Photon  
Cornell 1971-see discussion of Bj with  
Feynman about whether partons are  
hard (Bj) or soft (Feynman) pp.296-7

# BBK 1971-continued: the era of SCALING

♡ BBK propose a **General Form** for high  $p_T$  cross sections, for the **EM** scattering, which must exist:

$$E \frac{d^3\sigma}{dp^3} = \frac{4\pi\alpha^2}{p_T^4} \mathcal{F}(x_1 = \frac{-\hat{u}}{\hat{s}}, x_2 = \frac{-\hat{t}}{\hat{s}}) \quad (4)$$

♡ The two factors are a  $1/p_T^4$  term, characteristic of single photon exchange and a form factor  $\mathcal{F}$


♡ Note that  $x_{1,2}$  are not  $x_{BJ}$

□ The point is that  $\mathcal{F}$  **scales**, i.e. is only a function of the ratio of momenta.

♡ Vector ( $J = 1$ ) Gluon exchange gives the same form as Eq. 4 but much larger.

# CCR at the CERN-ISR

## Discovery of high $p_T$ $\pi^0$ production in p-p



F.W. Büsser, *et al.*,  
CERN, Columbia, Rockefeller  
Collaboration  
Phys. Lett. **46B**, 471 (1973)

Bjorken scaling PR179(1969)1547 →  
BermanBjKogut scaling PRD4(71)3388  
→ Blankenbecler, Brodsky, Gunion  
 $x_T = 2p_T/\sqrt{s}$  Scaling PL **42B**, 461 (1972)


$$E \frac{d^3 \sigma}{dp^3} = \frac{1}{p_T^{n_{\text{eff}}}} F\left(\frac{p_T}{\sqrt{s}}\right) = \frac{1}{\sqrt{s}^{n_{\text{eff}}}} G(x_T)$$

$n_{\text{eff}}$  gives the form of the force-law  
between constituents:  $n_{\text{eff}}=4$  for QED

- $e^{-6p_T}$  breaks to a power law at high  $p_T$  with characteristic  $\sqrt{s}$  dependence
- Large rate indicates that partons interact strongly ( $\gg$  EM) with other.
- Data follow  $x_T = 2p_T/\sqrt{s}$  scaling but with  $n_{\text{eff}}=8!$ , not  $n_{\text{eff}}=4$  as expected for QED

# $x_T$ scaling with $n=8$ , not 4

## Inspires Constituent Interchange Model



$$E \frac{d^3\sigma}{dp^3} = \frac{1}{p_T^n} F\left(\frac{p_T}{\sqrt{s}}\right)$$

$$x_T = 2p_T/\sqrt{s}$$

$n=4$  for QED or vector gluon

$n=8$  for quark-meson  
scattering by the exchange  
of a quark

CIM-Blankenbecler, Brodsky, Gunion,  
Phys.Lett.**42B**,461(1972)

# Constituent Interchange Model

Blankenbecler, Brodsky, Gunion, *Inclusive Processes at High Transverse Momentum*,  
Phys. Lett. **42B**, 461(1972)

♥ Inspired by the *dramatic features* of pion inclusive reactions revealed by “the recent measurements at CERN ISR of single-particle inclusive scattering at 90° and large transverse momentum”, Blankenbecler, Brodsky and Gunion propose a new general scaling form:

$$E \frac{d^3\sigma}{dp^3} = \frac{1}{p_T^n} F\left(\frac{p_T}{\sqrt{s}}\right) \quad (5)$$

♥  $n$  gives the form of the force-law between constituents


♥  $n = 4$  for QED or Vector Gluon

♥ Perhaps more importantly, BBG predict  $n=8$  for the case of quark-meson scattering by the exchange of a quark, **C.I.M.**, as apparently observed.

# State of the Art

## Fermilab 1977

D. Antreasyan, J. Cronin, et al., PRL **38**, 112 (1977)



Beautiful  $x_T$  scaling at all 3 fixed target energies with  $n=8$

Totally Misleading--Not CIM or QCD but  $k_T$

# First prediction using 'QCD' 1975

R. F. Cahalan, K. A. Geer, J. Kogut and Leonard Susskind, Phys. Rev. **D11**, 1199 (1975)

*“Asymptotic freedom and the “absence” of vector-gluon exchange in wide-angle hadronic collisions”*



♥ **Abstract:** The naive, pointlike parton model of Berman, Bjorken and Kogut is generalized to scale-invariant and asymptotically free field theories. The asymptotically free field generalization is studied in detail. Although such theories contain vector fields, **single vector-gluon exchange contributes insignificantly to wide-angle hadronic collisions**. This follows from (1) the smallness of the invariant charge at small distances and (2) the *breakdown of naive scaling* in these theories. These effects should explain the apparent absence of vector exchange in inclusive and exclusive hadronic collisions at large momentum transfers observed at Fermilab and at the CERN ISR.

♥ An interesting **Acknowledgement:** ... Two of us (J. K. and L. S. also thank S. Brodsky for *emphasizing to us repeatedly* that the present data on wide-angle hadron scattering *show no evidence for vector exchange*.

♥ Nobody's perfect, they get *one* thing right! They introduce the “effective index”  $n(x_T, \sqrt{s})$  to account for ‘scale breaking’:

$$E \frac{d^3\sigma}{dp^3} = \frac{1}{p_T^{n(x_T, \sqrt{s})}} F\left(\frac{p_T}{\sqrt{s}}\right) = \frac{1}{\sqrt{s}^{n(x_T, \sqrt{s})}} G\left(\frac{p_T}{\sqrt{s}}\right)$$

# CCOR 1978--Discovery of “REALLY high $p_T > 7$ GeV/c” at ISR



CCOR A.L.S. Angelis, et al,  
Phys.Lett. **79B**, 505 (1978)

QCD: Cahalan, Geer, Kogut,  
Susskind, PRD**11**, 1199 (1975)

$$E \frac{d^3\sigma}{dp^3} = \frac{1}{\sqrt{s}^{n_{\text{eff}}(x_T, \sqrt{s})}} G(x_T)$$

$$\left( \frac{\sqrt{s_1}}{\sqrt{s_2}} \right)^{n_{\text{eff}}(x_T, \sqrt{s})} = \frac{E \frac{d^3\sigma}{dp^3}(x_T, \sqrt{s_2})}{E \frac{d^3\sigma}{dp^3}(x_T, \sqrt{s_1})}$$

$$\heartsuit E d^3\sigma/dp^3 \simeq p_T^{-5.1 \pm 0.4} (1 - x_T)^{12.1 \pm 0.6}$$


$$7.5 \leq p_T \leq 14.0 \text{ GeV/c,}$$

$$53.1 \leq \sqrt{s} \leq 62.4 \text{ GeV}$$


(including *all* systematic errors).

$n_{\text{eff}}=5$  ( $=4^{++}$ ) as predicted for QCD

# 1978- $n_{\text{eff}}(x_T, \sqrt{s})$ WORKS $n_{\text{eff}} \rightarrow 5 = 4^{++}$





C.Kourkoulis, et al  
Phys.Lett. **84B**, 279 (1979)




cross sections vary  
by factor of 2

But  $n(x_T, \sqrt{s})$  agrees




A.Adare, et al, PHENIX  
PRD**79** (2009) 012003



# p.s. Fermilab experiment (1980) kills CIM


H. J. Frisch, et al., PRL **44**, 511 (1980)  $\pi^- + p \rightarrow \pi^\pm + X$

If quark-meson scattering by exchange of a quark dominates  
then  $\pi^-$  should dominate  $\pi^+$  at large  $p_T$



Statement in PLB**637**, 58 (2006): "We find that high- $p_T$  hadrons are produced by different mechanisms at fixed-target and collider energies. For pions, higher-twist subprocesses where the pion is produced directly dominate at fixed target energy," is contradicted by this measurement

# CCRS-1974 Discovery of direct $e^\pm \sim 10^{-4} \pi^\pm$ at ISR not due to internal conversion of direct photons



CCRS PLB**53**(1974)212; NPB**113**(1976)189

Data points  $(e^+ + e^-)/2$  lines  $10^{-4} (\pi^+ + \pi^-)/2$

•Farrar and Frautschi PRL**36**(1976)1017 proposed that direct leptons are due to internal conversion of direct photons with  $\gamma/\pi \sim 10\text{-}20\%$  to  $e^+e^-$  ( $d\sigma/dm \sim 1/m$ ) for  $p_T > 1.3$  GeV/c. CCRS looks, finds very few events, sets limits excluding this.

95% confidence level upper limits for a particle of mass  $m$ , or a mass continuum, which decays to  $e^+e^-$  with branching ratio  $B$ , at  $\sqrt{s} = 52.7$  GeV/c

| Mass<br>(GeV/c <sup>2</sup> ) | $B \frac{d\sigma}{dy} (p_T^* > 1.3 \text{ GeV/c})$<br>(cm <sup>2</sup> ) | Fraction of single<br>electron signal |
|-------------------------------|--------------------------------------------------------------------------|---------------------------------------|
| 0.400                         | $5.54 \times 10^{-33}$                                                   | 0.064                                 |
| 0.500                         | $8.37 \times 10^{-33}$                                                   | 0.104                                 |
| 0.600                         | $1.64 \times 10^{-32}$                                                   | 0.178                                 |

p.s. these direct  $e^\pm$  are due to semi-leptonic decay of charm particles not discovered until 1976, 2 year later: Hinchliffe and Llewellyn-Smith NPB**114**(1976)45

# J/Psi and direct $e^\pm$ at the CERN-ISR

First





FIG. 2

Fig. 2. Invariant mass distribution for the observed  $e^+e^-$  pairs. The curves represent the shapes of the acceptance, as a function of the  $e^+e^-$  invariant mass value, for the Arm 1 and Arm 2 triggers, respectively.


Best

A.G. Clark et al. / Electron pair production at the ISR



Not cause of direct  $e^\pm$

F.W. Büsser et al. / Electrons at the ISR



CCRS PLB**56**(1975)482  
2nd J/ $\Psi$  in Europe

CSZ NPB**142**(1978)29  
 $\langle p_T \rangle = 1.10 \pm 0.05$  GeV/c

CCRS NPB**113**(1976)189  
direct  $e^\pm$  not due to J/ $\Psi$

# Diversion: An application leads to a DISCOVERY

## Charm in PHENIX—a signal or a background?



M. J. Tannenbaum

Brookhaven National Laboratory  
Upton, NY 11973-5000 USA

*Received 26 June 1996*

**Abstract.** Charm, as well as Strangeness, plays an important role in searches for the Quark Gluon Plasma.  $J/\Psi$  Suppression and Strangeness Enhancement are two of the earliest proposed QGP signatures. Recent theoretical work on charm in Relativistic Heavy Ion collisions has focussed on dilepton production. However, even before the discovery of the  $J/\Psi$ , evidence of open charm was seen in hadron collisions via the observation of prompt **single** leptons “resulting from the semi-leptonic decays of charm particles.”[1] The ‘copious’ yield of direct (i.e. not from Dalitz decays) single electrons and muons—at a level  $e/\pi \sim 10^{-4}$  for  $p_T \geq 1.3$  GeV/c—observed in the early 1970’s was explained by Hinchliffe and Llewellyn-Smith and Bourquin and Gaillard as evidence of open-charm production. It is likely that  $e/\pi$  at RHIC is large and is a good measure of charm production. Thus, a measurement of single electrons with moderate  $p_T > 1.5$  GeV/c at RHIC should give a clean charm signal in heavy ion collisions, **with no combinatoric background.**

# Quark Matter 2005-Budapest-single $e^\pm$ in AuAu



Single  $e^\pm$  from heavy quark decay suppressed as much as  $\pi^0$  from light quarks; and they flow. Disfavors radiative energy loss

Theory curves

(1-3) from N. Armesto, *et al.*, PRD 71, 054027

(4) from M. Djordjevic, M. Gyulassy, S.Wicks, PRL 94, 112301


**Major Discovery---still not understood in 2012**

# Back to QCD

# Direct photon production-simple theory hard experiment

See the classic paper of Fritzsche and Minkowski, PLB **69** (1977) 316-320

$$A+B \rightarrow \gamma + X$$



Analytical formula for  $\gamma$ -jet cross section for a photon at  $p_T$ ,  $y_c$  (and parton (jet) at  $p_T$ ,  $y_d$ ):

$$\begin{aligned} \frac{d^3\sigma}{dp_T^2 dy_c dy_d} = & x_1 g_A(x_1, Q^2) F_{2B}(x_2, Q^2) \frac{\pi\alpha\alpha_s(Q^2)}{3\hat{s}^2} \left( \frac{1 + \cos\theta^*}{2} + \frac{2}{1 + \cos\theta^*} \right) \\ & + F_{2A}(x_1, Q^2) x_2 g_B(x_2, Q^2) \frac{\pi\alpha\alpha_s(Q^2)}{3\hat{s}^2} \left( \frac{1 - \cos\theta^*}{2} + \frac{2}{1 - \cos\theta^*} \right) \end{aligned}$$

$$\cos\theta^* = \tanh\frac{(y_c - y_d)}{2}$$

$$x_{1,2} = x_T \frac{e^{\pm y_c} + e^{\pm y_d}}{2}$$


$g(x)$  and  $F_2(x)$  are g and q pdf's in nuclei A,B

# The first 'evidence' for direct- $\gamma$ was wrong; but internal conversions provided a stringent limit






- L.~Yuan, E.~Amaldi Rome, BNL, CERN PLB **77** (1978) 240 set a limit on real photons using PbGl:  $\gamma/\pi^0=0.021\pm0.012$   $3.5<p_T<5.0$  GeV/c
- This corrected a **notoriously wrong** result of  $\gamma/\pi^0=0.20\pm0.06 \pm 0.07$  for  $2.8<p_T<3.8$  GeV/c also using PbGl P.Darriulat *et al*, NPB**110** (1976) 365
- The most stringent limit came from (non observation) of low mass e+e- pairs from internal conversion BNL CERN Syracuse Yale, J. Cobb *et al*, PLB**78** (1978) 519  
 $\gamma/\pi^0=0.006\pm0.009$   $2<p_T<3$  GeV/c

# The first 'evidence' for direct- $\gamma$ was wrong; but internal conversions provided a stringent limit

- L.~Yuan, E.~Amaldi Rome, BNL, CERN PLB **77** (1978) 240 set a limit on real photons using PbGl:  $\gamma/\pi^0=0.021\pm0.012$   $3.5<p_T<5.0$  GeV/c
- This corrected a **notoriously wrong** result of  $\gamma/\pi^0=0.20\pm0.06 \pm 0.07$  for  $2.8<p_T<3.8$  GeV/c also using PbGl P.Darriulat *et al*, NPB**110** (1976) 365
- The most stringent limit came from (non observation) of low mass  $e^+e^-$  pairs from internal conversion BNL CERN Syracuse Yale, J. Cobb *et al*, PLB**78** (1978) 519  $\gamma/\pi^0=0.006\pm0.009$   $2<p_T<3$  GeV/c



# Final ISR direct- $\gamma$ production + correlations




No evidence for bremsstrahlung contribution to direct  $\gamma$ --same side correlation is zero--see CMOR NPB327, 541 (1989) for full list of references.



# → QM2005-direct $\gamma$ in AuAu via internal conversion

Kroll Wada PR98(1955) 1355

PHENIX NPA774(2006)403



$$\frac{1}{N_\gamma} \frac{dN_{ee}}{dm_{ee}} = \frac{2\alpha}{3\pi} \frac{1}{m_{ee}} \left(1 - \frac{m_{ee}^2}{M^2}\right)^3 |F(m_{ee}^2)|^2 \sqrt{1 - \frac{4m_e^2}{m_{ee}^2}} \left(1 + \frac{2m_e^2}{m_{ee}^2}\right)$$



Eliminating the  $\pi^0$  background by going to  $0.2 < m_{ee} < 0.3$  GeV enables direct  $\gamma$  signal to be measured for  $1 < p_T < 3$  GeV/c in Au+Au. It is exponential, does that mean it is thermal? Yes: the p-p direct  $\gamma$  turns over as  $p_T \rightarrow 0$  follows the same function  $B(1+p_T^2/b)^{-n}$  used in Drell

Fit to Au+Au is  $[A e^{-p_T/T} + \langle T_{AA} \rangle B_{pp} (1+p_T^2/b_{pp})^{-n_{pp}}]$ . Significance of exponential (thermal) is  $> 3 \sigma$ . Fitted  $T=220\text{MeV}$  (is time average)

# Back to 1978

## Status of ISR single particle measurements

---

♡ Hard-scattering was visible both at ISR and FNAL (Fixed Target) energies by single particle inclusive at large  $p_T \geq 2\text{-}3 \text{ GeV}/c$ .

♡ Scaling and dimensional arguments for plotting data revealed the systematics and underlying physics.

♡ The theorists had the basic underlying physics correct; but many (inconvenient) details remained to be worked out, several by experiment.

♡  $k_T$ , the transverse momentum imbalance of outgoing partons (due to initial state radiation), was discovered by experiment.

**$k_T$  is what made  $n=4^{++} \rightarrow n=8$**

# $k_T$ is not a parameter, it can be measured

- In leading order QCD or the Quark-Parton model, the net transverse momentum  $\langle p_T \rangle_{\text{pair}} = \sqrt{2} \times \langle k_T \rangle$ , of a hard-scattering jet-pair, or a Drell-Yan pair, or a pair of high  $p_T$  photons, or the  $\gamma$ + Jet pair for direct photon production is zero. All the above pairs should be coplanar with the incident beam axis.

- However, early Drell-Yan and inclusive high  $p_T$  particle studies showed that  $k_T$  was measurable and non-zero.

♡ The history of  $k_T$  is worth reviewing as  $k_T$  was predicted to be zero by theorists, but was discovered to be non-zero by experimentalists. The CCHK experiment [M. Della Negra, et al., Nucl. Phys. **B127**, 1 (1977)] discovered that back-to-back jets had considerable out of plane transverse momentum  $p_{\text{out}}$ , and proposed that this was due to transverse momentum of partons inside a proton.

# Feynman Field & Fox to the rescue

♥ This was elaborated by Feynman, Field and Fox, [Nucl. Phys. **B128**, 1, (1977), Phys Rev. **D18**, 3320 (1978)] who introduced the  $k_T$  phenomenology of a parton in a proton, which they discussed in terms of ‘intrinsic transverse momentum’ from confinement which would be constant as a function of  $x$  and  $Q^2$ , and NLO effects due to hard gluon emission which would vary with  $x$  and  $Q^2$ , but they used an constant ‘effective’  $k_T$  to ‘explain’ the available measurements.

♥ A subsequent ISR experiment, CCOR, showed that  $k_T$  for jet-pairs was roughly the same as for Drell-Yan and increased similarly with  $\sqrt{s}$  (and  $p_T$ ) i.e. was not constant.

# $k_T$ and NLO are distinct---e.g. Drell Yan





FIG. 12. Invariant yield of dimuons as a function of the transverse momentum  $p_T$  of the muon pair for 400 GeV incident protons. A.S.Ito, et al, PRD23,604 (1981)




Note Gaussian shape, no power-law tail!


# $\langle p_T \rangle (= \sqrt{2}k_T)$ vs $\sqrt{s}$ in Drell-Yan

A.L.S. Angelis et al. / Massive electron pairs


CMOR, NPB348, 1 (1991)



# Owens and FFF QCD calculations inclusive $\pi^0$



Owens, Kimel PRD18(1978)3313



Feynman, Field, Fox, PRD18(1978)3320

Note that  $k_T$  smearing dramatically improves agreement at lower  $p_T$

# Status of QCD Theory in 1978

- The first modern QCD calculation and prediction for high  $p_T$  single particle inclusive cross sections including non-scaling and initial state radiation was done in 1978 by J. F. Owens, E. Reya, M. Gluck, PRD **18**, 1501 (1978), “*Detailed quantum-chromodynamic predictions for high- $p_T$  processes*,” and J.F. Owens, J. D. Kimel, PRD **18**, 3313 (1978), “*Parton-transverse-momentum effects and the quantum-chromodynamic description of high- $p_T$  processes*”.
- This work was closely followed and corroborated by Feynman, Field, Fox PRD **18**, 3320 (1978), “*Quantum-chromodynamic approach for the large-transverse-momentum production of particles and jets*.”
- Unfortunately jets in  $4\pi$  Calorimeters at ISR energies or lower are invisible below  $\sqrt{s} \approx 10$  GeV, which led to considerable confusion in the period 1980-1982.

# QCD and Jets

are now a cornerstone of the standard model

- Incredibly at the famous Snowmass conference in July 1982, many if not most people in the U.S. were skeptical

e.g. MJT Seminar in 1982

WHY I BELIEVE  
IN JETS

IN SPITE OF  
CALORIMETER EXPERIMENTS


M. J. TANNENBAUM 3/22/82

FOR QCD TESTS MUST  
MINIMIZE THE EFFECT OF JETS


- The International HEP conference in Paris, three weeks later, July 26--31, 1982 changed everything.

# Paris 1982-THE UA2 Jet

From 1980--1982 most high energy physicists doubted jets existed because of the famous NA5  $E_T$  spectrum which showed NO JETS. This one event from UA2 in 1982 changed everybody's opinion.



(a)




(b)

# Why nobody (in the U.S.) believed in jets

- In 1972-73, soon after hard-scattering was discovered in p-p collisions, [Bjorken PRD8 \(1973\) 4098](#) and [Willis \(ISABELLE Physics Prospects-BNL-17522\)](#) proposed  $4\pi$  hadron calorimeters to search for jets from fragmentation of scattered partons with large  $p_T$  realizing that a substantial increase in rate would be expected in measuring the entire jet at a given  $p_T$  rather than just the leading fragment. (Bjorken's parent-child effect)
- It took until 1980 to get a full azimuth  $\Delta\eta \sim \pm 0.88$  ( $\Delta\Theta \sim \pm 45^\circ$ ) calorimeter but meanwhile experiments were done with smaller back-to-back calorimeters each with aperture  $\Delta\Phi \sim \pm 45^\circ$   $\Delta\eta \sim \pm 0.55$  and many new trigger biases were discovered, for instance, jets wider than the calorimeter aperture would deposit less energy than narrow jets of the same  $p_T$  and be suppressed by the steeply falling spectrum  $\Rightarrow$  jet structure is dominated by the calorimeter geometry [e.g. see [M. Dris NIM 158 \(1979\) 89](#)]


# Willis 'impactometer' from Isabelle Study 1972

$4\pi$  hadron calorimeter non-magnetic detector. Sound familiar?



**IMPACTOMETER**  
FOR 200 GEV PROTONS ON P OR E  
MEASURES  $\pi\psi = \sum_i E_i \sin \theta_i \leftarrow E_T$

# (In)famous FNAL E260 found “Jets” (1977)





- In each of 2 back to back calorimeters with  $\Delta\Phi \sim \pm 45^\circ$   
 $\Delta\eta \sim \pm 0.36$  (same as PHENIX)  
the invariant cross section of several particles with a vector sum  $p_T$  is much larger than a single particle of the same  $p_T$ .  
The authors took this as evidence for the exactly back-to-back in azimuth jets of constituent scattering  $\Rightarrow$  **Never let an interested theorist collaborate on an experiment.**

**C.Bromberg et al E260, PRL 38 (1977)1447, NPB134 (1978) 189**

# But, experiments with different apertures got different results


- The first  $4\pi$  experiment was a bubble chamber(!) 110 GeV/c  $K^-$  on p [M. Deutschmann, et al, ABCCLVW collab, NPB**155** (1979)307]



- multiparticle cross section for  $p_T > 1.5 \text{ GeV}/c \gg$  single particle
- Data extrapolate nicely to those of E260 [8] in slope and magnitude.
- But “principal axis” analysis of the data shows “the vast majority of events with large  $p_T$  multiparticle systems DO NOT exhibit jet-like structure.”

# NA5-the coup-de-grâce to jets (1980)

- Full azimuth calorimeter  $-0.88 < \eta^* < 0.67$  ( $\rightarrow$  NA35, NA49)



- plus triggered in two smaller apertures corresponding to E260.

- No jets in full azimuth data

- All data way above QCD predictions-----

- The large  $E_T$  observed is the result of “a large number of particles with a rather small transverse momentum”--the first  $E_T$  measurement in the present terminology.


K. Pretzl, Proc 20th ICHEP (1980)

C. DeMarzo et al NA5, PLB112(1982)173


For more on  $E_T$  see MJT IJMPA 4 (1989)3377

# Back to Paris 1982-THE UA2 Jet

From 1980--1982 most high energy physicists doubted jets existed because of the famous NA5  $E_T$  spectrum which showed NO JETS. This one event from UA2 in 1982 changed everybody's opinion.





(a)



(b)

# UA1-Carlo himself explained $E_T$ (no jet) dist. before seeing UA2 plot. Explanation is correct






UA1 (1982) Paris-withdrawn (C.Rubbia)  $\sqrt{s}=540$  GeV. No Jets because  $E_T$  is like multiplicity ( $n$ ), composed of many soft particles near  $\langle p_T \rangle$  ! CERN-EP-82/122.

OOPS UA2 discovers jets ~5-6 orders of magnitude down in  $E_T$  distribution!


# Also Paris 1982-first measurement of QCD subprocess angular distribution using $\pi^0$ - $\pi^0$ correlations

DATA: CCOR NPB 209, 284 (1982)

Di Pion Angular Distributions *CONSTITUENT COM POLAR ANGLE*  
 $\sqrt{s} = 62.4 \text{ GeV}$



QCD







$$\frac{d^3\sigma}{dx_1 dx_2 d\cos\theta^*} = \frac{1}{s} \sum_{ab} f_a^A(x_1) f_b^B(x_2) \frac{\pi\alpha_s^2(Q^2)}{2x_1 x_2} \Sigma^{ab}(\cos\theta^*)$$

$\Sigma^{ab}(\cos\theta^*)$ , the characteristic subprocess angular distributions  
 and  $\alpha_s(Q^2) = \frac{12\pi}{25 \ln(Q^2/\Lambda^2)}$  are predicted by QCD

# Also Paris 1982-first measurement of QCD subprocess angular distribution using $\pi^0$ - $\pi^0$ correlations

DATA: CCOR NPB 209, 284 (1982)


Di Pion Angular Distributions  $\sqrt{s} = 62.4 \text{ GeV}$  *CONSTITUENT COM POLAR ANGLE*



$$\frac{d^3\sigma}{dx_1 dx_2 d\cos\theta^*} = \frac{1}{s} \sum_{ab} f_a^A(x_1) f_b^B(x_2) \frac{\pi \alpha_s^2(Q^2)}{2x_1 x_2} \Sigma^{ab}(\cos\theta^*)$$

$\Sigma^{ab}(\cos\theta^*)$ , the characteristic subprocess angular distributions  
and  $\alpha_s(Q^2) = \frac{12\pi}{25 \ln(Q^2/\Lambda^2)}$  are predicted by QCD

# LO-QCD in 1 slide



# LO-QCD in 1 slide

## Cross Section in p-p collisions c.m. energy $\sqrt{s}$

The overall p-p reaction cross section  
is the sum over constituent reactions

$$a + b \rightarrow c + d$$

$f_a^A(x_1)$ ,  $f_b^B(x_2)$ , are structure functions, the differential probabilities  
for constituents  $a$  and  $b$  to carry momentum fractions  $x_1$  and  $x_2$   
of their respective protons, e.g.  $u(x_1)$ ,

$$\frac{d^3\sigma}{dx_1 dx_2 d\cos\theta^*} = \frac{1}{s} \sum_{ab} f_a^A(x_1) f_b^B(x_2) \frac{\pi\alpha_s^2(Q^2)}{2x_1 x_2} \Sigma^{ab}(\cos\theta^*)$$

$\Sigma^{ab}(\cos\theta^*)$ , the characteristic subprocess angular distributions  
and  $\alpha_s(Q^2) = \frac{12\pi}{25 \ln(Q^2/\Lambda^2)}$  are predicted by QCD

# $\Sigma^{ab}(\cos\theta^*)$ in LO-QCD

a)  $qq' \rightarrow qq'$   $\frac{4}{9} \frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2}$   
 $\bar{q}q' \rightarrow \bar{q}q'$

b)  $qq \rightarrow qq$   $\frac{4}{9} \left( \frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2} + \frac{\hat{s}^2 + \hat{t}^2}{\hat{u}^2} \right) - \frac{8}{27} \frac{\hat{s}^2}{\hat{u}\hat{t}}$

c)  $\bar{q}q \rightarrow \bar{q}'q'$   $\frac{4}{9} \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2}$


d)  $\bar{q}q \rightarrow \bar{q}q$   $\frac{4}{9} \left( \frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2} + \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2} \right) - \frac{8}{27} \frac{\hat{u}^2}{\hat{s}\hat{t}}$

e)  $\bar{q}q \rightarrow gg$   $\frac{32}{27} \frac{\hat{u}^2 + \hat{t}^2}{\hat{u}\hat{t}} - \frac{8}{3} \frac{\hat{u}^2 + \hat{t}^2}{\hat{s}^2}$



f)  $gg \rightarrow \bar{q}q$   $\frac{1}{6} \frac{\hat{u}^2 + \hat{t}^2}{\hat{u}\hat{t}} - \frac{3}{8} \frac{\hat{u}^2 + \hat{t}^2}{\hat{s}^2}$

g)  $qg \rightarrow qg$   $-\frac{4}{9} \frac{\hat{u}^2 + \hat{s}^2}{\hat{u}\hat{s}} + \frac{\hat{u}^2 + \hat{s}^2}{\hat{t}^2}$

h)  $gg \rightarrow gg$   $\frac{9}{2} \left( 3 - \frac{\hat{u}\hat{t}}{\hat{s}^2} - \frac{\hat{u}\hat{s}}{\hat{t}^2} - \frac{\hat{s}\hat{t}}{\hat{u}^2} \right)$




qq MOLLER






$\bar{q}q$  BHABHA

COMPTON




qg




gg



gg




DRELL-YAN



$\bar{q}q$

QCD is like QED except for the gluon self coupling

# Constituent Kinematics






$$Q^2 = -\hat{t} = \hat{s} \frac{(1 - \cos \theta^*)}{2}$$

$$-\hat{u} = \hat{s} \frac{(1 + \cos \theta^*)}{2}$$


# Back to Paris 1982-first measurement of QCD subprocess angular distribution using $\pi^0$ - $\pi^0$ correlations: need $\alpha_s(Q^2=\hat{t})$

DATA: CCOR NPB 209, 284 (1982)

Di Pion Angular Distributions *CONSTITUTENT*  
 $\sqrt{s} = 62.4 \text{ GeV}$  *CoM POLAR ANGLE*



QCD







$$\frac{d^3\sigma}{dx_1 dx_2 d\cos\theta^*} = \frac{1}{s} \sum_{ab} f_a^A(x_1) f_b^B(x_2) \frac{\pi\alpha_s^2(Q^2)}{2x_1 x_2} \Sigma^{ab}(\cos\theta^*)$$

$\Sigma^{ab}(\cos\theta^*)$ , the characteristic subprocess angular distributions  
 and  $\alpha_s(Q^2) = \frac{12\pi}{25 \ln(Q^2/\Lambda^2)}$  are predicted by QCD

# Back to Paris 1982-first measurement of QCD subprocess angular distribution using $\pi^0$ - $\pi^0$ correlations: need $\alpha_s(Q^2=\hat{t})$

DATA: CCOR NPB 209, 284 (1982)

Di Pion Angular Distributions  $\sqrt{s} = 62.4 \text{ GeV}$  *CONSTITUTENT COM POLAR ANGLE*



$$\frac{d^3\sigma}{dx_1 dx_2 d\cos\theta^*} = \frac{1}{s} \sum_{ab} f_a^A(x_1) f_b^B(x_2) \frac{\pi\alpha_s^2(Q^2)}{2x_1 x_2} \Sigma^{ab}(\cos\theta^*)$$

$\Sigma^{ab}(\cos\theta^*)$ , the characteristic subprocess angular distributions  
and  $\alpha_s(Q^2) = \frac{12\pi}{25 \ln(Q^2/\Lambda^2)}$  are predicted by QCD

# Eventually this was measured with di-jets





Figure 10 (a) Distribution of  $\cos \theta^*$  for hard parton scattering as measured in the UA1 experiment (42). The normalization is defined by setting the value at  $\cos \theta^* = 0$  equal to 1. (b) Distribution of  $\cos \theta^*$  for hard parton scattering as measured in the UA2 experiment (43). All the different QCD processes (except for  $\rightarrow q'\bar{q}'$ ), separately normalized to the data, lie in the area between the two dashed curves. The full line is the overall QCD prediction, normalized to the data.

see L. Di Lella ARNPS 35 (1985) 107--134


# QCD really works: CCOR p-p follows q-q, UA1 p-p follows q-q



plot I made in 1983


Why I believed in Jets:  
At the CERN ISR from  
1975-1982 two-particle  
correlations showed  
unambiguously  
that high  $p_T$  particles  
come from jets

# How everything you want to know about JETS was measured with 2-particle correlations




CCOR, A.L.S. Angelis, et al Phys.Lett. **97B**, 163 (1980) PhysicaScripta **19**, 116 (1979)

$p_{Tt} > 7 \text{ GeV/c}$  vs  $p_T$





Away side  $p_{out} \sim p_T \Delta\phi$  is not constant i.e  $\Delta\phi \neq 1/p_T$ , indicating jets not collinear in azimuth  $\Rightarrow k_T$

# How everything you want to know about JETS was measured with 2-particle correlations




CCOR, A.L.S. Angelis, et al Phys.Lett. **97B**, 163 (1980) PhysicaScripta **19**, 116 (1979)

$p_{Tt} > 7 \text{ GeV/c}$  vs  $p_T$





Away side  $p_{out} \sim p_T \Delta\phi$  is not constant i.e  $\Delta\phi \neq 1/p_T$ , indicating jets not collinear in azimuth  $\Rightarrow k_T$

# How everything you want to know about JETS was measured with 2-particle correlations




CCOR, A.L.S. Angelis, et al Phys.Lett. **97B**, 163 (1980) PhysicaScripta **19**, 116 (1979)

$p_{Tt} > 7 \text{ GeV/c}$  vs  $p_T$





Away side  $p_{out} \sim p_T \Delta\phi$  is not constant i.e  $\Delta\phi \neq 1/p_T$ , indicating jets not collinear in azimuth  $\Rightarrow k_T$

# How everything you want to know about JETS was measured with 2-particle correlations




CCOR, A.L.S. Angelis, et al Phys.Lett. **97B**, 163 (1980) PhysicaScripta **19**, 116 (1979)

$p_{Tt} > 7 \text{ GeV/c}$  vs  $p_T$





Away side  $p_{out} \sim p_T \Delta\phi$  is not constant i.e  $\Delta\phi \neq 1/p_T$ , indicating jets not collinear in azimuth  $\Rightarrow k_T$

# How everything you want to know about JETS was measured with 2-particle correlations



CCOR, A.L.S. Angelis, et al Phys.Lett. **97B**, 163 (1980) PhysicaScripta **19**, 116 (1979)

$p_{Tt} > 7 \text{ GeV/c}$  vs  $p_T$



Away side  $p_{out} \sim p_T \Delta\phi$  is not constant i.e  $\Delta\phi \neq 1/p_T$ , indicating jets not collinear in azimuth  $\Rightarrow k_T$

# Feynman, Field and Fox said that $x_E$ distribution from single particle or Jet measures $D(z)$

38

R.P. Feynman et al. / Large transverse momenta

FFF Nucl.Phys. B128(1977) 1-65




Fig. 23. Comparison of the  $\pi^0$  and jet trigger away-side distribution of charged hadrons in pp collisions at  $W = 19.4$  GeV,  $\theta_1 = 90^\circ$ , and  $p_\perp(\text{trigger}) = 4.0$  GeV/c from the quark-quark scattering model. The upper figure shows the single-particle ( $\pi^0$ ) trigger results plotted versus  $z_p = -p_x(h^\pm)/p_\perp(\pi^0)$  and the jet trigger plotted versus  $z_J = -p_x(h^\pm)/p_\perp(\text{jet})$  (see table 1). In the lower figure, we plot both versus  $z_J$ , where for the jet trigger  $z_J = z_J$  but for the single-particle trigger  $z_J = \langle z_c \rangle z_p$ . The away hadrons are integrated over all rapidity  $Y$  and  $|180^\circ - \phi| \leq 45^\circ$  and the theory is calculated using  $\langle k_\perp \rangle_{h \rightarrow q} = 500$  MeV.  $\bullet$   $h_1 = \pi^0$ ,  $\times$   $h_1 = \text{jet}$ .

“There is a simple relationship between experiments done with single-particle triggers and those performed with jet triggers. The only difference in the opposite side correlation is due to the fact that the ‘quark’, from which a single-particle trigger came, always has a higher  $p_\perp$  than the trigger (by factor  $1/z_{\text{trig}}$ ). The away-side correlations for a single-particle trigger at  $p_\perp$  should be roughly the same as the away side correlations for a jet trigger at  $p_\perp(\text{jet}) = p_\perp(\text{single particle}) / \langle z_{\text{trig}} \rangle$ .”

# As shown at the ISR by Darriulat, et al, and believed by most High Energy Physicists

P. Darriulat, et al, Nucl.Phys. **B107** (1976) 429-456




Figure 21 Jet fragmentation functions measured in different processes:  $\nu$ -p interactions (open triangles, Van der Welde 1979);  $e^+e^-$  annihilations (solid line, Hanson et al 1975); and pp collisions (full circles CS,  $p_T < 6$  GeV/c, open circles CS,  $p_T > 6$  GeV/c, full squares CCOR,  $p_T > 5$  GeV/c, open squares CCOR,  $p_T > 7$  GeV/c).







Figure 19 The slopes  $b$  obtained from exponential fits to the jet fragmentation function in the interval  $0.2 < z < 0.8$  in  $e^+e^-$  annihilation (full circles) and LPTH data of the BS Collaboration (open circles).

Figures from P. Darriulat, ARNPS **30** (1980) 159-210 showing that Jet fragmentation functions in  $\nu p$ ,  $e^+e^-$  and pp (CCOR) are the same with the same dependence of  $b$  (exponential slope) on “ $\hat{s}$ ”

# But first, CCHK discovered $k_T$ by lack of $x_E$ scaling






CCHK  
NPB127(1977) 1-42



Cern-College de France, Heidelberg, Karlsruhe (CCHK) collab. found that  $x_E$  distributions were not universal for  $2 < p_{Tt} < 4$  GeV/c and attributed this to the large out of plane momentum  $p_{out}$  which was not constant with  $x_E$  as in fragmentation but increased with increasing  $x_E$  as if the di-jets were not collinear due to initial state  $k_T$

# CCOR $\langle |p_{out}| \rangle^2$ vs $x_E^2$ -- $k_T$ not constant


$$\langle |p_{out}| \rangle^2 = x_E^2 [2 \langle |k_{Ty}| \rangle^2 + \langle |j_{Ty}| \rangle^2] + \langle |j_{Ty}| \rangle^2$$



CCOR PLB 97 (1980) 163  
 $k_T$  varies with  $p_{Tt}$  and  $\sqrt{s}$   
 Not intrinsic !


# But $j_T$ is constant-independent of $p_{Tt}$ and $\sqrt{s}$

## Characteristic of jet fragmentation



- it took the  $e^+ e^-$  people several more years to get this correct--- because they didn't understand the seagull effect: ( $j_T < p_T$ )

# $x_E$ distribution measures fragmentation fn.



CCOR, Physica Scripta **19**, 116 (1979)

$$x_E \sim Z / \langle Z_{\text{trig}} \rangle$$


$$\langle Z_{\text{trig}} \rangle = 0.85 \text{ measured}$$

$$e^{-5.3x_E} \Rightarrow D_{\pi}^q(z) \sim e^{-6z}$$

- independent of  $p_{Tt}$

See M. Jacob's talk Proc. EPS 1979  
Geneva (CERN). p512

# $x_E$ distribution measures fragmentation fn. not!



CCOR, Physica Scripta **19**, 116 (1979)

$$x_E \sim Z / \langle Z_{\text{trig}} \rangle$$

$$\langle Z_{\text{trig}} \rangle = 0.85 \text{ measured}$$

$$e^{-5.3x_E} \Rightarrow D^q_{\pi}(z) \sim e^{-6z} *$$

- independent of  $p_{Tt}$

See M. Jacob's talk Proc. EPS 1979  
Geneva (CERN). p512

\* but we did learn something new  
on this issue in PHENIX.

# This is the only thing we didn't understand correctly at the ISR. Maybe we could be forgiven because Feynman said it.

---

\* At RHIC we learned that the  $x_E$  distribution from a trigger fragment does not measure the fragmentation function.



For more info, see: M. J. Tannenbaum “Review of hard scattering and jet analysis”, PoS (CFRNC2006) cited by Kronfeld and Quigg in “Resource Letter: QCD” arXiv:1002.5032v2. Even

better, see J. Rak and M. J. Tannenbaum, “High  $p_T$  physics in the Heavy Ion Era”, Cambridge University Press, available March 2013

[http://www.cambridge.org/us/knowledge/isbn/item6947421/High-/?site\\_locale=en\\_US](http://www.cambridge.org/us/knowledge/isbn/item6947421/High-/?site_locale=en_US)

For the past decade these single and two-particle techniques were used exclusively at RHIC for hard-scattering, with outstanding results...

# Jet measurements of QCD in pp collisions are now standard after a $\sim 30$ year learning curve



The measured crosssection is in agreement with NLO pQCD predictions after the necessary nonperturbative parton-to-hadron corrections are taken into account. i.e. Make sure to read the fine print!

A. Abulencia, et al, CDF PRL 96 (2006) 122001- $k_T$  algorithm




At RHIC, inclusive single particles provide a precision pQCD probe, well calibrated in pp, dAu... collisions

# PHENIX excellent in hard-scattering measurements via single-inclusive and two-particle correlations, STAR better with Jets

PHENIX PRL91 (2003) 241803

PHENIX arXiv:1205.5533

PHENIX PRL97 (2006) 252002




direct-single-e from  
c and b quark decays


In p-p collisions, since 1978, NLO pQCD agrees very well with all measurements.

# Of course LHC MUCH Better with Jets

STAR PRL97 (2006) 252001



CMS PRL107 (2011) 132001




Again agree very well with NLO pQCD in p-p collisions. But, I have known that QCD worked for hard scattering since 1978. What I learn from the CMS plot is that partons are pointlike up to  $Q^2 \approx t \approx 2p_T^2 = 2,000,000 \text{ GeV}^2$  i.e.  $r \ll 1.4 \times 10^{-4} \text{ fm}!!$

# Direct $\gamma$ p-p data and pQCD c. 2007

PHENIX direct- $\gamma$  in p-p  
PRL **98** (2007) 012002

PHENIX direct  
photon p-p data  
clarify longstanding  
data/theory puzzle

P. Aurenche et al Phys. Rev.  
D **73**, 094007 (2006)




# Direct $\gamma$ p-p data and pQCD c. 2007

PHENIX direct- $\gamma$  in p-p  
PRL **98** (2007) 012002


PHENIX direct  
photon p-p data  
clarify longstanding  
data/theory puzzle

P. Aurenche et al Phys. Rev.  
D **73**, 094007 (2006)


New PHENIX p-p results this year  
arXiv:1205.5533 are even better!



# $x_T$ scaling: a) Direct- $\gamma$ b) Jets




Direct  $\gamma$   $n \approx 5$  2005



Jets  $n \approx 4.5$


# QCD in Action 2012



$x_T$  scaling with  $n_{\text{eff}}=4$  (parton model) QCD non-scaling is visible

Collection of World's direct- $\gamma$  measurements (p+p/ p+pbar) including PHENIX low  $p_T$  msmt. to be described next.


# QCD in Action 2012



$x_T$  scaling with  $n_{\text{eff}}=4.5$  works for direct- $\gamma$  due to QCD non-scaling

Collection of World's direct- $\gamma$  measurements (p+p/ p+pbar) including PHENIX low  $p_T$  msmt. to be described next.

# The END



Gunther Wolf, Rapporteur, Proc. 1982 ICHEP

# Recap of discoveries and techniques from the CERN ISR in 1972-1982

G. Giacomelli and M. Jacob, Phys. Rept. **55** (1979) 1-132  
M. Jacob and K. Johnsen, CERN Yellow Report 84-13

- The rapidity plateau. (Not discussed in this talk. )
- Hard scattering in p-p collisions via particle production at large  $p_T$  which proved that the partons of DIS strongly interacted with each other.  $x_T$  scaling measurements to find the underlying physics.
- direct lepton ( $e^\pm$ ) production from the decay of (unknown at that time-1974) particles composed of b and c quarks.
- first and only J/Psi cross section measurement for all pair  $p_T \geq 0$  at a hadron collider, until PHENIX at RHIC [[PRL 92 \(2004\) 051802](#)] and CDF [[PRD 71\(2005\) 032001](#) (15 years after their first publication)]
- direct photon production
- Proof using same-side and away side two particle correlations that high  $p_T$  particles in p-p collisions are produced from states with two roughly back-to-back jets which are the result of scattering of constituents of the nucleons as described by **QCD**, which was developed during the course of these measurements.

# Conclusions from ISR

- Hard Scattering in p-p collisions was discovered at the CERN ISR in 1972 by the method of leading particles.
- A very large flux of high  $p_T$  pions was observed with a power-law tail which varied systematically with  $\sqrt{s}$ , the c.m. energy of the collision.
- The huge flux of high  $p_T$  particles proved that the partons of DIS strongly interacted with each other.
- Scaling arguments allowed the form of the force law between ‘partons’ to be determined but there was some early confusion caused by initial transverse momentum  $k_T$  which distorted the spectra.
- Further ISR measurements utilizing inclusive single or pairs of hadrons established that high transverse momentum particles are produced from states with two roughly back-to-back jets which are the result of scattering of constituents of the nucleons as described by Quantum Chromodynamics.