

Lepton Physics at RHIC

M. J. Tannenbaum
BNL

VII Workshop on Non-Perturbative QCD
Villefranche-sur-Mer, La Citadelle
January 2003


Results on physics at RHIC using outgoing leptons will be presented from Au+Au collisions at nucleon-nucleon c.m. energies $\sqrt{s_{NN}} = 130$ GeV and 200 GeV, and from p-p collisions at $\sqrt{s_{NN}} = 200$ GeV. Introduction and motivation will be presented both from the theoretical and experimental perspectives. Topics include open charm production via single $e^\pm$, $J/\Psi \rightarrow e^+ + e^-$, $\mu^+ + \mu^-$, leptonic decay of vector mesons and the dilepton continuum.

7TH QCD WORKSHOP, VILLEFRANCHE-SUR-MER, FRANCE,
JAN 2003

Quantum Chromo Dynamics

QCD

- QCD is the theory of the strong interaction
- The force between a quark and an antiquark is free and coulomb-like at short distances but is string-like at large r



- The coulomb-like part leads to narrow bound states
 - The string-like tension leads to confinement
- Narrow bound states of heavy quarks, c , b , were discovered in 1970's via large branching ratios to di-leptons $\sim 10^{-2}$
 - c.f. $\sim 10^{-4}$ for light quark vector mesons ρ , ω .

Best Lepton Pair data c. 1977





Figure 1: Lederman's spectrum of di-muon invariant mass at mid-rapidity

Recent State of the Art FNAL



Bound States follow the QCD potential Model

VOLUME 34, NUMBER 6

PHYSICAL REVIEW LETTERS

10 FEBRUARY 1975




FIG. 1. The spectrum of charmonium. The vertical scale is schematic; our predictions of masses for the

TABLE I. γ ray widths.^a


Transition	Γ_γ	Γ_γ (keV)
$2^3S \rightarrow 3P_2$	$5I_1\alpha k^3$	120
$\rightarrow 3P_1$	$3I_1\alpha k^3$	70
$\rightarrow 3P_0$	$1I_1\alpha k^3$	25
$3P_2 \rightarrow 1^3S$	$I_2\alpha k^3$	240
$3P_1 \rightarrow 1^3S$	$I_2\alpha k^3$	240
$3P_0 \rightarrow 1^3S$	$I_2\alpha k^3$	240
$1^1P_1 \rightarrow 1^1S$	$I_2\alpha k^3$	240
$3D_1 \rightarrow 3P_2$	$1I_2\alpha k^3$	7
$\rightarrow 3P_1$	$15I_2\alpha k^3$	110
$\rightarrow 3P_0$	$20I_2\alpha k^3$	150
$2^3S \rightarrow 1^1S$	$I_4\alpha k^7$	~ 1

^aIn the second column $1/\alpha = 137$, k is the energy of the transition, and I_n is a radial integral. The last column is based on our wave functions and energy differences, with fine-structure splittings and S-D mixing ignored.

P multiplet lies about 230 MeV below that of the $2S$ levels. This energy difference is not very sensitive to our choice of parameters: It decreases to 160 MeV if α_s and m_c assume the

Charmonium becomes Open Charm

- The force between a quark and an antiquark is free and coulomb-like at short distances but is string-like at large r





- The string-like tension leads to confinement BUT
- At larger r , the string tension limit is exceeded
- the string breaks forming a $q \bar{q}$ pair which binds to $Q\bar{Q}$
- making a $D=Q\bar{q}$ $\bar{D}=\bar{Q}q$ open charm (Beauty) pair.

$$V(r) = -\frac{4}{3}\frac{\alpha_s}{r} + \sigma r$$

- $V(r \rightarrow \infty) = 2M_D$ for $T = 0$

High Temperature $T \rightarrow T_c$


- $\sigma \rightarrow 0$ as $T \rightarrow T_c$, $2M_D \rightarrow 0$
- Excited quarkonia dissociate to open charm, beauty




- ‘Like’ Chiral Symmetry Restoration

Tightest bound quarkonia survive at T_c but

- $\alpha_s(T) \downarrow$ as $T \uparrow$ but $\alpha_s(T_c)$ is finite $\Rightarrow$ Quarkonia can exist above T_c .
- BUT attractive potential is ‘Debye screened’ in QGP $\Rightarrow$ Dissociation of Quarkonia in QGP = Deconfinement!



$\chi_c, \psi', J/\psi$



$\Upsilon'', \chi'_b, \Upsilon', \chi_b, \Upsilon$


[H. Satz, et al., Phys. Lett. B**178**, 416 (1986), Phys. Lett. B**514**, 57 (2001), Phys. Rev. D**64**, 094015 (2001)]

QUARKONIA

The Road To Success in HEP


LETTERS B

5 March 1987



$p_T(\mu) \geq 3 \text{ GeV}/c$, UA1 Phys. Lett. B186, 237 (1987)

The Road To Success in HIP



The Physics of Open Charm (Beauty)

- Produced via $g + g \rightarrow c + \bar{c}$ so measures gluon structure function.
- charm and J/ψ both suppressed $\Rightarrow$ Shadowing in Gluon Structure Fn.
- Large mass scale $\Rightarrow$ sensitive to initial T
- Large mass scale $\Rightarrow$ energy loss in QGP is less than for light quarks $\Rightarrow$ less suppression of c jets than light quark jets due to ‘jet quenching’ in hot (deconfined) medium [‘Dead Cone Effect’, Y. L. Dokshitzer and D. E. Kharzeev, Phys. Lett. B**519**, 199 (2001)]

But a Key reason to study Charm is Experimental

- Large semi-leptonic branching ratio $\sim 7\text{-}17\%$ per lepton (e, μ)
- Large mass $\Rightarrow \sim 1$ GeV leptons ($\gg \langle p_T \rangle$)
- Lepton I.D. in large hadronic and photonic background is an experimental challenge **BUT**
- Charm via single lepton measurement has **NO Combinatoric Background** $\Rightarrow$ same difficulty in p-p and A+A collisions.

Why (Some) Experimentalists Study Leptons at High p_T From Hadron Collisions

They indicate weak ($e^\pm \nu$) or EM decays ($e^+ e^-$)
 $\Rightarrow$ Long Lived Particles $\tau \gg 1 \text{ fm}/c$ ($3.3 \times 10^{-23} \text{ sec}$)

- Originally (1964) inspired by the search for the Intermediate W boson of weak interactions in p-p collisions.

- “Zichichi” Signature [Proc. 12th ICHEP, Dubna, 1964, p35 (footnote)] hadron spectra fall like e^{-6p_T} , decay lepton spectra fall faster and can be calculated, upon which the decay $W \rightarrow e + \nu$ from heavy W produced at rest (mid-rapidity) would give a (jacobian) peak at $p_{T_e} = M_W/2$

That’s Actually How the $W^\pm$ was discovered in 1983!




Figure 2: $W^\pm \rightarrow e^\pm + X$ from UA2 c. 1984

However Great Excitement in 1974

Discovery of prompt electrons in p-p collisions

$e^\pm/\pi^\pm \sim 10^{-4}$, $p_T \geq 1.3$ GeV/c—no peak




Figure 3: Invariant cross section for $(e^+ + e^-)/2$ vs p_T for 5 values of $\sqrt{s}$ at the CERN-ISR compared to fits (solid lines) to corresponding data for $(\pi^+ + \pi^-)/2$ [CCRS, *Phys. Lett. B***53**, 212 (1974), *Nucl. Phys. B***113**, 189 (1976)]

- Discovered before J/ψ
- Not due to J/ψ but $\langle p_T \rangle|_{J/\psi}$ is a key issue.

F.W. Büsser et al. / Electrons at the ISR




Figure 4: CCRS calculation of $e^\pm$ from J/ψ as a function of $\langle p_T \rangle$ of the J/ψ

1976—Prompt Electrons = Open Charm


- Some previous (and even post) spurious results: MJT in 1974 “All those who try to measure direct leptons become the world’s experts in η Dalitz Decay ($\eta \rightarrow \gamma + e^+ + e^-$)”
- First suggestion of direct photons in p-p collisions as the source of prompt leptons [G. R. Farrar and S. C. Frautschi, Phys. Rev. Lett. **36**, 1017 (1976), and others] well before pQCD predicts $g + q \rightarrow \gamma + q$ “Inverse QCD compton effect”
- 1976—Finally explained as evidence for open charm. [I. Hinchliffe and C. H. Llewellyn Smith, Phys. Lett. B**61**, 472 (1976); M. Bourquin and J.-M. Gaillard, Nucl. Phys. B**114**, 334 (1976) (The original “Coctail Model”)]

The ‘modern’ view

Heavy flavor measurement

• ideal but very challenging

- direct reconstruction of heavy flavor decays (e.g. $D^0 \rightarrow K^- \pi^+$)



• alternative but indirect

- heavy flavor semi leptonic decays contribute to single lepton and lepton pair spectra

Experimental Issues

Real Backgrounds, Falling Spectra

Photons from $\pi^0 \rightarrow \gamma + \gamma$ follow the π^0 curve
down by a factor of $2/(n-1)$




Figure 5: PHENIX π^0 spectrum for $\sqrt{s} = 200$ GeV p-p collisions

$$\frac{dn_{\pi^0}}{p_T dp_T} \propto p_T^{-n}$$

where $n \sim 8$, so $\gamma|_{\pi^0}/\pi^0 \sim 1.2 \times 2/7 = 0.34$ and the factor 1.2 includes $\eta \rightarrow \gamma + \gamma$ (estimated).

- This assumes that e.g. one 10 GeV photon and two 5 GeV photons (a 10 GeV π^0) measure at exactly the same energy in the experiment $\Rightarrow$ **nonlinearity** is a big experimental issue for photons.

PHENIX π^0 spectra at $\sqrt{s_{NN}} = 200$ GeV





Figure 6: PHENIX QM02 π^0 spectra in Au+Au peripheral and central collisions

- The inclusive γ spectrum **calculated** from the above π^0 spectrum shown below as $\gamma|_{\pi^0}/\pi^0$.



- Note that $\gamma|_{\pi^0}/\pi^0 \sim \text{constant}$ for $p_T \geq 3$ GeV/c consistent with power law with $n \sim 8$ for π^0 (and assumed for other particles).

PHENIX inclusive γ spectrum





Figure 7: PHENIX QM02 inclusive γ spectrum in Au+Au central collisions at $\sqrt{s_{nn}} = 200$ GeV **statistical errors only**

- Are the γ direct or all from decays? Hard to compare two 6-7 order log plots. Analysis is best in $\gamma/\pi^0_{\text{meas}}/\gamma/\pi^0_{\text{decays}}$



- Background from $n, \bar{n}$? Possible non-linearity? η/π^0 ratio? ...

QM02 inclusive γ results PHENIX





Figure 8: PHENIX QM02 $\gamma/\pi^0|_{meas}/\gamma/\pi^0|_{decays}$ vs. p_T in Au+Au collisions at $\sqrt{s_{nn}} = 200$ GeV **Systematic Errors Included**

STAR

Boltzmann Assumption



π^0 Bose-Einstein Assumption




Figure 9: STAR QM02 $\gamma/\pi^0|_{\pi^0}/\gamma/\pi^0|_{meas}$ **Systematic Errors Included**

- Ratio is upside-down!
- Background from $n, \bar{n}$? Background from η, ω, η' ?
- Shape of π^0 spectrum!?
- Non-linearity not an issue—measured via conversions.

All Inclusive Photons create Background e^+e^- pairs




Figure 10: **CCRS** yield of inclusive electrons vs total external radiation length t/X_0 for (b) selected conversions (extrapolates to zero at the "Dalitz point") and (a) Accepted events with prompt electrons (non-zero intercept at Dalitz point). Yields are relative, normalized to 1 at the standard thickness $t/X_0 = 0.16$

- Probability of internal and external conversion per γ

$$\frac{e^-|_{\gamma}}{\gamma} = \frac{e^+|_{\gamma}}{\gamma} = \frac{\delta_2}{2} + \frac{t}{\frac{9}{7}X_0} \equiv \delta_{eff}$$

where $\delta_2/2 =$ Dalitz (internal conversion) branching ratio per photon = 0.6% π^0 , 0.8% for $\eta \rightarrow \gamma\gamma$.

- Must keep external $t/\frac{9}{7}X_0$ comparable $\sim 0.6\%$ to avoid additional background from external conversions.

- But can add small external converter in test run to determine whether $e^-/\gamma \rightarrow 0$ at "Dalitz Point" $-\frac{9}{7}\delta_2/2 \sim 0.8 - 1.0\%$ in units of radiation lengths, **insensitive** to the η/π^0 ratio. Converter curve measures the photonic background!

Effect of Falling Spectra

• Still assuming the p_T^{-n} power law for the π^0 and thus decay γ Spectra:

$$\frac{e^-}{\pi^0} \bigg|_{\pi^0} (p_T) = \frac{(e^- + e^+)}{2\pi^0} \bigg|_{\pi^0} (p_T) = \delta_{eff} \times \frac{2}{(n-1)^2} > 0.6\%/7^2 = 1.2 \times 10^{-4}$$

• Need $\sim 10^4$ rejection against $\pi^\pm$ just to be able to see the $e^\pm$ background from π^0 Dalitz. $\Rightarrow$ some experimental discussion.

For the record, it's actually easier in Au+Au at RHIC








Figure 11: PHENIX inclusive $\pi^\pm$, π^0 , h^- , $e^\pm$ p_T spectra in minimum bias Au+Au collisions at $\sqrt{s_{nn}} = 130$ GeV

- $e/\pi \sim 1/500 = 2 \times 10^{-3} \gg 1.2 \times 10^{-4}$
- Measured Conversions/Dalitz = 1.8 ± 0.2
- Since π are suppressed at high p_T , suggests that charm is not suppressed ...




Detecting electrons means detecting all particles=PHENIX



- ElectroMagnetic Calorimeter measures Energy of photons and electrons
 - reconstructs π^0 from 2 photons. Measures decent Time of Flight
 - hadrons deposit Minimum Ionization, or higher if they interact
- For electron ID require RICH (cerenkov) and matching energy in EMCal
 - Electron and photon energy can be matched to $< 1\%$ --No nonlinearity problem
- momentum +TOF=charged particle ID
- High Resolution TOF completes the picture giving excellent charged hadron PID



PHENIX--charm via single $e^\pm$ in Au+Au





● main systematic errors (band)

– pion spectra, ratio π/π^0 , ratio conversion/Dalitz (material) □ Need converter run

PHENIX--charm via single $e^\pm$ in Au+Au 130 cont'd

- Min bias and central Au+Au p_T distributions are consistent with pointlike binary scaling from pythia (p-p)
- consistent with ISR and NLO pQCD in measured $d\sigma_e/dy$ and total deduced charm cross section.



To go from p-p to A+A collisions: We know from DIS that Hard Scattering is Point-Like

E. Gabathuler, Total cross-section




Fig. 14. The A dependence of the inelastic muon cross-section as presented by Tannenbaum (see discussion).

AGS $\mu - A$ scattering data, from E. Gabathuler's talk, [Proc. 6th Int. Symposium on Electron and Photon Interactions at High Energies, Bonn (1973)].


♡ DIS is pointlike $A^{1.00}$ even at modest q^2 —no shadowing.

♡ Photoproduction is shadowed— $A^{0.91}$


♡ In the region of hard scattering ($p_T > 2 \text{ GeV}/c$) scaling from p-p to nuclear collisions should be simply proportional to the relative number of point-like encounters, corresponding to A (p+A), $A \times B$ (A+B) for the total rate and to T_{AB} , the overlap integral of the nuclear profile functions, as a function of centrality.

QM02 PHENIX 200 GeV charm via $e^\pm$

PHENIX Detector - Second Year Physics Run







- Some data with 1.7 % X_0 converter added
- converter only affects photonic component



- converter effect is much greater at low p_T , indicating
 - relatively much larger photonic component at low p_T
 - relatively smaller photonic (i.e. larger non-photonic) component at higher p_T (i.e. charm)
- Reduced systematic errors.

Single e p_T distribution vs Centrality


Consistent with binary (pointlike) scaling $\square$ charm is not suppressed



Conclusions from single $e^\pm$ measurement

- $e^\pm$ from charm decays measured by PHENIX in Au+Au collisions at RHIC
 - $e^\pm$ yields and spectral shapes
 - consistent with PYTHIA calculations for pp:
$$\sigma_{cc} = 330 \text{ (650)} \sigma_b \text{ at } 130 \text{ (200) GeV}$$
 - scaled to Au+Au by number of binary collisions
 - NA50: factor ~ 3 charm enhancement inferred at SPS
from $\pi^+ \pi^-$ pair enhancement (will discuss later)
 - no enhancement observed at RHIC in inclusive charm production.
 - RHIC: factor 3-4 suppression of high p_T hadrons relative to binary scaling
 - no suppression observed in $e^\pm$ from charm decays---follows binary scaling
- Seems to IMPLY**
- normal gluon structure function in initial state
□ no saturation, no large shadowing
 - possibly less energy loss of heavy quarks in dense medium:
“dead cone effect”? (Y. Dokshitzer, D. Kharzeev: hep-ph/0106202)

Lepton Pairs---Kinematics, Notation



$$\vec{p}_T = \vec{p}_{T1} + \vec{p}_{T2} \quad \vec{p}_L = \vec{p}_{L1} + \vec{p}_{L2}$$

$$m^2 = m_1^2 + m_2^2 + 2(E_1 E_2 - p_1 p_2 \cos \phi)$$



$$y = \frac{1}{2} \ln[(E + p_L)/m_T]$$

ϕ^* , θ^* , azimuthal and polar angles of the pair in its cm system

- simple if $p_T=0$, more complicated $p_T \neq 0$ collins-soper-sterman

[J. C. Collins, D. E. Soper and G. Sterman, **NPB250**, 199 (1985)]

Acceptance is geometrical but:



Acceptance (geometrical) = prob. to fall on detector for given p_T y m

- an average over $\square^*$, $\square^*$, (sometimes over p_T , y)---can be model dependent (e.g. if averaged $-3 < y < 3$, is y distribution flat?)


Efficiency = detection prob for lepton pair or lepton in acceptance

Survey of Physics Topics using lepton pairs

M. J. Tannenbaum, 7th QCD Workshop, Villefranche-sur-Mer, France, Jan 2003

19

MJT Physics topics using e^+e^- 11/96



Topic	Method	Comment
π^0	external conversions (+Dalitz) $m_{e^+e^-} \leq m_{\pi^0}$	Inclusive γ
η	$140 \leq m_{e^+e^-} \leq 400 \text{ MeV}/c^2$	Dalitz decay
Direct γ	$400 \leq m_{e^+e^-} \leq 600 \text{ MeV}/c^2$	Internal Conversion
ρ^0	$600 \leq m_{e^+e^-} \leq 900 \text{ MeV}/c^2$	Lose money on every sale...
ω^0	$m_{e^+e^-} \simeq 782 \pm 8 \text{ MeV}/c^2$	Should work (?)
ϕ	$m_{e^+e^-} \simeq 1019 \pm 4 \text{ MeV}/c^2$	compare K^+K^-
Charm	$e^\pm$ inclusive $p_T \geq 1.1 \text{ GeV}/c$	No combinatoric bkg.
$c - \bar{c}$	$1.6 \leq m_{e^+e^-} \leq 5 \text{ GeV}/c^2$	Jet Quenching
$c - \bar{c}$	$1.6 \leq m_{e^\pm\mu^\mp} \leq 5 \text{ GeV}/c^2$	Jet Quenching Msmt. Charm bkg for D-Yan
Drell-Yan	$4.0 \leq m_{e^+e^-} \text{ GeV}/c^2$	QCD Sanity check
J/Ψ	$m_{e^+e^-} \simeq 3097 \pm 0.09 \text{ MeV}/c^2$	QGP or Bust Inclusive, Central $(AB)^\alpha$ p_T dependence
Ψ'	$m_{e^+e^-} \simeq 3686 \pm 0.3 \text{ MeV}/c^2$	Rate Limit ?
Υ	$9.6 \leq m_{e^+e^-} \leq 10.6 \text{ GeV}/c^2$	Rate Limit


MJT Physics topics using e^+e^- 11/96

Topic	Method	Comment
π^0	external conversions (+Dalitz) $m_{e^+e^-} \leq m_{\pi^0}$	Inclusive γ
η	$140 \leq m_{e^+e^-} \leq 400 \text{ MeV}/c^2$	Dalitz decay
Direct γ	$400 \leq m_{e^+e^-} \leq 600 \text{ MeV}/c^2$	Internal Conversion
ρ^0	$600 \leq m_{e^+e^-} \leq 900 \text{ MeV}/c^2$	Lose money on every sale...
ω^0	$m_{e^+e^-} \simeq 782 \pm 8 \text{ MeV}/c^2$	Should work (?)
ϕ	$m_{e^+e^-} \simeq 1019 \pm 4 \text{ MeV}/c^2$	compare K^+K^-
Charm	$e^\pm$ inclusive $p_T \geq 1.1 \text{ GeV}/c$	No combinatoric bkg.
$c - \bar{c}$	$1.6 \leq m_{e^+e^-} \leq 5 \text{ GeV}/c^2$	Jet Quenching
$c - \bar{c}$	$1.6 \leq m_{e^\pm\mu^\mp} \leq 5 \text{ GeV}/c^2$	Jet Quenching Msmt. Charm bkg for D-Yan
Drell-Yan	$4.0 \leq m_{e^+e^-} \text{ GeV}/c^2$	QCD Sanity check
J/Ψ	$m_{e^+e^-} \simeq 3097 \pm 0.09 \text{ MeV}/c^2$	QGP or Bust Inclusive, Central $(AB)^\alpha$ p_T dependence
Ψ'	$m_{e^+e^-} \simeq 3686 \pm 0.3 \text{ MeV}/c^2$	Rate Limit ?
Υ	$9.6 \leq m_{e^+e^-} \leq 10.6 \text{ GeV}/c^2$	<i>Rate Limit</i>

Low Mass Pairs---ISR best data 1979

[A. Chilingarov, et al., Nucl. Phys. B151, 29 (1979)]

- ISR 'same side' low mass $e^+ e^-$ pair production with the condition $p_T \geq 1.6 \text{ GeV}/c + 0.5 m(\text{GeV}/c^2)$ and $\cos\theta \approx 1$ $e^+ e^-$ peaks are clear. Solid curve for $m \leq 500 \text{ MeV}/c^2$ is $\pi^0 \pi$ Dalitz decay. Other solid curve for $m \geq 100 \text{ MeV}/c^2$ is $c \bar{c}$ production with $73 \mu\text{b}$ cross section.
- Sum of 2 solid curves (dashed curve) seems to explain excess pairs in the range $400 < m < 600 \text{ MeV}/c^2$ where there is no known source of Dalitz decay.
- Also, excess could be explained by internal conversion of direct photon with $\sigma/\sigma_0 = 3.8\%$ at $p_T = 1.9 \text{ GeV}/c$, but this would scale like $1/m$ to lower masses in disagreement with data



These expt's are tricky:
continuing ``low mass pair anomaly'' results
in the period 1976-91 explained by wrong
assumption for π/π^0 ratio!!!!!!!


- pp measurement, [K. J. Anderson et al, PR37, 799 (1976)]
They noted π dalitz could explain effect if $\pi/(\pi+\pi^0)=60\%$
- Quark Matter conference `highlights' from 1984, H. J. Specht,
until 1991 U. Goerlach, when they use correct π/π^0 !!!!!
- Final publication Helios Collab., Eur. Phys. J. **C13**, 433 (2000)
- However `low mass anomaly' persists in more recent CERES
experiment at SpS fixed target.

CERES--low mass pairs at CERN-SpS

[CERES Collaboration, G. Hering, et al, nucl-ex/0203004]


- Excess in $0.2 < m < 0.6 \text{ GeV}/c^2$
- source of many theoretical fantasies.
- Authors claim it's not charm?
- Huge background subtraction.
- Strange way to quote msmt, not centrality based

?



On to the RHIC data □ □


RHIC data
PHENIX e^+e^- spectrum Au+Au 200 GeV



PHENIX e^+e^- spectrum at RHIC 200 GeV Au+Au


- The huge $e^\pm$ rate from conversions gives a HUGE random $e^+ e^-$ background. In general $N_{e^+} \neq N_{e^-}$
- $(P_{e^+} + P_{e^-})^2 = P_{e^+} P_{e^+} + P_{e^-} P_{e^-} + 2P_{e^+} P_{e^-}$
- Background subtraction by mixed $e^+ e^-$ events with normalization $N_{e^+e^-} = 2\sqrt{N_{e^+e^+} N_{e^-e^-}}$

Real and **Mixed** e^+e^- Distribution




Real - Mixed with systematic errors

Green multiplicative, yellow additive (bkg subtraction)




PHENIX intermediate mass results



- Need less background (i.e. Dalitz rejection) or more data before we can say anything serious



This is the mass range for thermal pairs

[Ralf Rapp, PRC63, 054907 (2001), also nucl-th/0204003]





[More classical reference: K. Kajantie, et al (KKMM) PRD**34**, 2746 (1986)]

Vector Meson Interlude---PHENIX $\phi \rightarrow KK ee$



$K^+ K^-$




$e^+ e^-$

PHENIX--phi comparison

Do observed branching ratios change?


$$\sqrt{s} \quad e^+e^- : \quad \frac{dN}{dy} = 5.4 \pm 2.5(stat)_{\pm 2.8}(sys)$$

$$\sqrt{s} \quad K^+K^- : \quad \frac{dN}{dy} = 2.01 \pm 0.22(stat)_{\pm 0.52}(sys)$$




- Cross sections are consistent using 'vacuum' branching ratios □ No medium effect (yet?)

STAR $\pi^+\pi^-$





- Even $\pi^+\pi^-$ is hard, long wait for e^+e^- but great hadron gas signature
- For the record: STAR $\pi^+\pi^-$ K^+K^- [PRC 65, 04190(R) (2002)]





Drell-Yan production is pointlike

Scales like AB or T_{AB} (ncoll)

Drell-Yan: Experimental K factor




J/ψ and Drell-Yan production-summary



- DY is pointlike
- J/ψ is hadronic
- Complicates suppression issue

NA50 J/ψ Suppression at CERN 1988-2000




- Is it normal $T_{AB}^{0.92}$?
- Plus comovers?
- Discontinuities?
- χ_c χ + χ suppression?
- J/ψ suppression?
- pT dependence is relevant

[NA50 collaboration, M.C. Abreu, et al., PLB **477**, 28 (2000)]

Is this QGP suppression--- p_T dependence?


$J/\psi < p_t^2 >$ vs. E_t : Model Comparison



can a QGP model fit simultaneously $(J/\psi)/(DY)$ and $\langle p_t^2 \rangle$ vs. E_t

J/ψ Production--State of the Art NA50


CERN fixed target Pb+Pb $s_{NN}=17.2$ GeV



J/ψ Production--PHENIX State of the Art


RHIC Au+Au $s_{NN}=200\text{GeV}$ J/ψ → e⁺e⁻

$N = 10.8 \pm 3.2 \text{ (stat)} \pm 3.8 \text{ (sys)}$



Minimum bias

$N = 5.9 \pm 2.4 \text{ (stat)} \pm 0.7 \text{ (sys)}$




Versus centrality


J/ψ Production--PHENIX State of the Art

RHIC Au+Au $s_{NN}=200\text{GeV}$ J/ψ → e⁺e⁻

$N=4.5 \pm 2.1$ (stat) ± 0.5 (sys)




$N=3.5 \pm 1.9$ (stat) ± 0.5 (sys)




J/ψ Production--PHENIX State of the Art

RHIC Au+Au $s_{NN}=200\text{GeV}$ J/ψ e⁺e⁻



J/ψ Production--PHENIX State of the Art



RHIC Au+Au $s_{NN}=200\text{GeV}$ J/ψ → e⁺e⁻



- Inconclusive, need 100-1000 times more data!
- p-p data also sparse, but more conclusive

J/ψ Production--PHENIX RHIC p-p s=200GeV

J/ψ e⁺e⁻ J/ψ μ⁺μ⁻




$$N_{J/\psi} = 24 \pm 6(\text{stat}) \pm 4(\text{sys})$$

$$B d\psi/dy = 52 \pm 13(\text{stat}) \pm 18(\text{sys}) \text{ nb}$$

$$1.2 < y < 1.7 \quad N_{J/\psi} = 26 \pm 6(\text{stat}) \pm 2.6(\text{sys})$$

$$1.7 < y < 2.2 \quad N_{J/\psi} = 10 \pm 4(\text{stat}) \pm 1.0(\text{sys})$$

J/ψ B-d σ /dy and σ (p-p $\sqrt{s}=200$ GeV)




$$B \cdot \sigma(pp \rightarrow J/\psi + X) = 226 \pm 36(stat) \pm 79(sys) \text{ nb}$$

$$\sigma(pp \rightarrow J/\psi + X) = 3.8 \pm 0.6(stat) \pm 1.3(sys) \text{ } \mu\text{b}$$

J/ψ $\sqrt{s}$ dependence in p-p

RHIC/PHENIX measurement is respectable and instructive!





CEM predictions (Phys.Lett.B390:323-328,1997)

J/ψ $\langle p_T \rangle$ (p-p $\sqrt{s}=200$ GeV)


$J/\psi \rightarrow e^+e^-$

$J/\psi \rightarrow \mu^+\mu^-$



RHIC p-p $\sqrt{s}=200$ GeV $\langle p_T \rangle_{J/\psi} \sim 1.6\text{--}2.0$ GeV/c

seems much larger than at CERN $\sqrt{s}=17.2$ GeV



$\langle p_T \rangle$ vs $\sqrt{s}$

Agrees with trend from lower $\sqrt{s}$ [Apanasevich, et al, PRD59, 074007 (1999)]




FIG. 1. $\langle p_T \rangle$ of pairs of muons, photons, and jets produced in hadronic collisions versus $\sqrt{s}$.

Conclusions--leptons/photons at RHIC

- Proof of principle established for J/ψ in Au+Au
 - need a factor 100-1000 (10000) more data
- Significant result for J/ψ cross section in p-p with 60 events
- charm via single $e^\pm$ in AuAu indicates point-like scaling
 - suggestive of difference in light and heavy quark jets
 - first measurement of charm in RHI collisions
- Low mass region, inconclusive, huge background
 - Need lots more data, or, preferably Dalitz rejection
- π , η , ω some early results, still a long way to go in e^+e^-
- Direct photons---systematics still under study, results this yr.

A very successful start of what
we hope should be a long and fruitful program