PHENIX results on small systems collectivity at RHIC

Ron Belmont
University of North Carolina at Greensboro

RHIC & AGS Annual Users' Meeting
Online
22 October 2020

Standard model of heavy ion physics



Based on developments in hydro theory over the last few years, we might replace "thermalization" with "hydrodynamization"

Azimuthal anisotropy measurements





$$\frac{dN}{d\varphi} \propto 1 + \sum_{n=1}^{\infty} 2v_n \cos n\varphi \qquad v_n = \langle \cos n\varphi \rangle \qquad \varepsilon_n = \frac{\sqrt{\langle r^n \cos n\varphi \rangle + \langle r^n \sin n\varphi \rangle}}{\langle r^n \rangle}$$

- Hydrodynamics translates initial shape (including fluctuations) into final state distribution

Azimuthal anisotropy measurements





$$\frac{dN}{d\varphi} \propto 1 + \sum_{n=1}^{\infty} 2v_n \cos n\varphi \qquad v_n = \langle \cos n\varphi \rangle \qquad \varepsilon_n = \frac{\sqrt{\langle r^n \cos n\varphi \rangle + \langle r^n \sin n\varphi \rangle}}{\langle r^n \rangle}$$

- Hydrodynamics translates initial shape (including fluctuations) into final state distribution

Important discovery in 2005

PHOBOS Plenary, Quark Matter 2005 (see also Phys.Rev.C 77, 014906 (2008))



A nucleus isn't just a sphere

Important discovery in 2005

PHOBOS Plenary, Quark Matter 2005 (see also Phys.Rev.C 77, 014906 (2008))



A nucleus isn't just a sphere

Important discovery in 2005

PHOBOS Plenary, Quark Matter 2005 (see also Phys.Rev.C 77, 014906 (2008))



A nucleus isn't just a sphere

Important discovery in 2005

R. Andrade et al, Eur. Phys. J. A 29, 23-26 (2006)

NeXSPheRIO results on elliptic flow at RHIC and connection with thermalization

R.Andrade[1], F.Grassi[1], Y.Hama[1], T.Kodama[2], O.Socolowski Jr.[3], and B.Tavares[2]

[1] Instituto de Física, USP,
C. P. 66318, 05315-970 São Paulo-SP, Brazil

[2] Instituto de Física, UFRJ,
C. P. 68528, 21945-970 Rio de Janeiro-RJ , Brazil

[3] CTA/ITA,
Praça Marechal Eduardo Gomes 50, CEP 12228-900 São José dos Campos-SP, Brazil

Received 1 January 2004



Worth noting that lumpy initial conditions were predicted some time in 2003

Important discovery in 2010

Alver and Roland, Phys. Rev. C 81, 054905 (2010)



Nucleon fluctuations can produce non-zero ε_n for odd n

Symmetry planes ψ_n can be different for different harmonics

$$\varphi = \phi_{lab} - \psi_n$$

Fluctuations in large systems

PHOBOS, Phys. Rev. C 81, 034915 (2010)



Fluctuations should also be translated, so measure $\sigma_{v_2}/\langle v_2\rangle$

$|\eta| < 1$

Generally good agreement with models of initial geometry

Fluctuations in large systems

PHENIX, Phys. Rev. C 99, 024903 (2019)



Fluctuations should also be translated, so measure $\sigma_{v_2}/\langle v_2\rangle$

$1 < |\eta| < 3$

Generally good agreement with models of initial geometry

Fluctuations in large systems

PHENIX, Phys. Rev. C 99, 024903 (2019)



Fluctuations should also be translated, so measure $\sigma_{v_2}/\langle v_2 \rangle$

$1 < |\eta| < 3$

Central: breakdown of small-variance limit (assumed in data and solid line)

Peripheral: non-linearity in hydro response (e.g. J. Noronha-Hostler et al Phys. Rev. C 93, 014909 (2016))

Testing hydro by controlling system geometry

PHYSICAL REVIEW LETTERS

Highlights Recent Accepted Collections Authors Referees Search Press About

Exploiting Intrinsic Triangular Geometry in Relativistic $^{3}\mathrm{He} + \mathrm{Au}$ Collisions to Disentangle Medium Properties

J. L. Nagle, A. Adare, S. Beckman, T. Koblesky, J. Orjuela Koop, D. McGlinchey, P. Romatschke, J. Carlson, J. E. Lynn, and M. McCumber

Phys. Rev. Lett. **113**, 112301 – Published 12 September 2014

- Collective motion translates initial geometry into final state distributions
- To determine whether small systems exhibit collectivity, we can adjust the geometry and compare across systems
- We can also test predictions of hydrodynamics with a QGP phase

Testing hydro by controlling system geometry

nature physics

Letter | Published: 10 December 2018

Creation of quark–gluon plasma droplets with three distinct geometries

PHENIX Collaboration

Nature Physics **15**, 214–220(2019) | Cite this article



Testing hydro by controlling system geometry

Nature Physics 15, 214–220 (2019)





Testing hydro by controlling system geometry

Nature Physics 15, 214–220 (2019)



- v_2 and v_3 ordering matches ε_2 and ε_3 ordering in all three systems
  —Collective motion of system translates the initial geometry into the final state

Testing hydro by controlling system geometry

Nature Physics 15, 214–220 (2019)



- v_2 and v_3 vs p_T predicted or described very well by hydrodynamics in all three systems
  —All predicted (except v_2 in d+Au) in J.L. Nagle et al, PRL 113, 112301 (2014)
  —v_3 in p+Au and d+Au predicted in C. Shen et al, PRC 95, 014906 (2017)

Testing hydro by controlling system geometry

Nature Physics 15, 214–220 (2019)



- Initial state effects alone do not describe the data
  —Phys. Rev. Lett. 123, 039901 (Erratum) (2019)

Testing hydro by controlling system geometry

Nature Physics 15, 214–220 (2019)



- Important to include initial state effects
  —B. Schenke et al, Phys. Lett. B 803, 135322

Comparisons with STAR



Good agreement between STAR and PHENIX for v_2

Comparisons with STAR



Good agreement between STAR and PHENIX for v_2

Large discrepancy between STAR and PHENIX for v_3

Comparisons with STAR

PHENIX takes the issue seriously, so we are doing our due diligence!

The published small systems results use the event plane method, where the resolution nominally follows
$R(\chi) = \frac{\sqrt{\pi}}{2}\chi e^{-\frac{\chi^2}{2}} \left(I_0(\frac{\chi^2}{2}) + I_1(\frac{\chi^2}{2}) \right)$
In small systems we're in the limiting case where $\chi \ll 1$ so $R \propto \chi$ (note that $\chi = v_n\sqrt{N_{ch}}$).

The set of PHENIX event plane resolutions do not follow the expected pattern.

The origin of this effect is the beam and angle offset relative to the detector and an additional offset of the PHENIX central carriage (all of these things vary between operational periods). **The effect is qualitatively reproduced in toy simulation studies that utilize the full analysis procedure.**

The three-subevent 2-particle correlation method uses event mixing, which corrects these effects quite well. **Checks with the 3x2PC method show no such bias as seen in EP method for all systems, and all of these checks agree with published EP results within uncertainties.**

Further checks on going as part of due diligence!

Comparisons with STAR





- Most PHENIX results use backward rapidity detectors in combination with mid-rapidity
- STAR uses mid-rapidity only (TPC, $|\eta| < 1$)

Comparisons with STAR





- Most PHENIX results use backward rapidity detectors in combination with mid-rapidity
- STAR uses mid-rapidity only (TPC, $|\eta| < 1$)
- PHENIX can use a backward-mid-forward combination to better mimic STAR acceptance
  - —Unsubtracted STAR results match PHENIX backward-mid-forward
  - —Non-flow subtracted STAR results match PHENIX backward-backward-mid

Comparisons with STAR





- Most PHENIX results use backward rapidity detectors in combination with mid-rapidity
- STAR uses mid-rapidity only (TPC, $|\eta| < 1$)
- PHENIX can use a backward-mid-forward combination to better mimic STAR acceptance
  —Unsubtracted STAR results match PHENIX backward-mid-forward
  —Non-flow subtracted STAR results match PHENIX backward-backward-mid
- v_3 with both detector combinations coming soon!

Initial eccentricities

Table compiled by J.L. Nagle

System	Nagle Nucleons w/o NBD fluctuations	Welsh Nucleons w/ NBD fluctuations	Welsh Quarks w/ NBD and Gluon fluctuations	IPGlasma w/ Nucleons t=0	IP-Glasma w/ 3 Quarks t=0
ε_2 p+Au	0.23	0.32	0.38	**0.10**	**0.50**
ε_2 d+Au	0.54	0.48	0.51	**0.58**	**0.73**
ε_2 ^{3}He+Au	0.50	0.50	0.52	**0.55**	**0.64**
ε_3 p+Au	0.16	0.24	0.30	**0.09**	**0.32**
ε_3 d+Au	0.18	0.28	0.31	**0.28**	**0.40**
ε_3 ^{3}He+Au	0.28	0.32	0.35	**0.34**	**0.46**

- Nagle et al: https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.113.112301
- Welsh et al: https://journals.aps.org/prc/abstract/10.1103/PhysRevC.94.024919
- IP-Glasma run by S. Lim using publicly available code (thanks to B. Schenke)

Longitudinal dynamics in small systems

Phys. Rev. Lett. 121, 222301 (2018)



p+Al, p+Au, d+Au, ^{3}He+Au

Good agreement with wounded quark model
(M. Barej et al, Phys. Rev. C 97, 034901 (2018))

Good agreement with 3D hydro
(P. Bozek et al, Phys. Lett. B 739, 308 (2014))

Longitudinal dynamics in small systems

Phys. Rev. Lett. 121, 222301 (2018)



- v_2 vs η in p+Al, p+Au, d+Au, and ^{3}He+Au
- Good agreement with 3D hydro for p+Au and d+Au (Bozek et al, PLB 739, 308 (2014))

Longitudinal dynamics in small systems

Phys. Rev. Lett. 121, 222301 (2018)



- v_2 vs η in p+Al, p+Au, d+Au, and ^{3}He+Au
- Good agreement with 3D hydro for p+Au and d+Au (Bozek et al, PLB 739, 308 (2014))
- Prevalence of non-flow near the EP detector ($-3.9 < \eta < -3.1$)

Longitudinal dynamics in small systems

Phys. Rev. Lett. 121, 222301 (2018)



- v_2 vs η in p+Al, p+Au, d+Au, and ^{3}He+Au
- Good agreement with 3D hydro for p+Au and d+Au (Bozek et al, PLB 739, 308 (2014))
- Prevalence of non-flow near the EP detector ($-3.9 < \eta < -3.1$)

Testing hydro by controlling system size and life time

J.D. Orjuela Koop et al
Phys. Rev. C 93, 044910 (2016)



Geometry in d+Au collisions dominated by deuteron shape, thus largely independent of collision energy

Spacetime volume of system in QGP phase decreases with decreasing collision energy

d+Au beam energy scan

Phys. Rev. C 96, 064905 (2017)



- Hydro theory agrees with higher energies very well, underpredicts lower energies
- Likely need different EOS for lower energies; influence of conserved charges likely more important at lower energies (see e.g. M. Martinez et al, arXiv:1911.10272, 1911.12454)
- Nonflow likelier to be an issue due to lower multiplicity at lower energies

d+Au beam energy scan

Phys. Rev. Lett. 120, 062302 (2018)



d+Au beam energy scan

Phys. Rev. Lett. 120, 062302 (2018)

200 GeV | 62.4 GeV | 39 GeV | 19.6 GeV



- Measurement of $v_2\{6\}$ in d+Au at 200 GeV and $v_2\{4\}$ in d+Au at all energies

d+Au beam energy scan

Phys. Rev. Lett. 120, 062302 (2018)



- Measurement of $v_2\{6\}$ in d+Au at 200 GeV and $v_2\{4\}$ in d+Au at all energies
- Multiparticle correlations can be a good indicator of collectivity

Summary

- Collective motion translates initial geometrical shape into final state azimuthal anisotropies
- Evidence of this translation is seen in small and large systems
- The v_2 and v_3 in p+Au, d+Au, and ^{3}He+Au qualitatively follow the geometrical ordering of the initial state
- The v_2 and v_3 in p+Au, d+Au, and ^{3}He+Au quantitatively agree with multiple hydrodynamical calculations
- Several internal cross-checks of published PHENIX small system results give confidence to the measurements, which have remained unchanged
- A variety of collective signatures are seen in the d+Au beam energy scan
  —v_2 vs p_T agrees with hydro at the higher two energies
  —Observation of multiparticle correlations at all energies

Extra material

Components and cumulants in p+Au and d+Au at 200 GeV

Phys. Rev. Lett. 120, 062302 (2018)

$$v_2\{4\} = (-c_2\{4\})^{1/4}$$

Negative $c_2\{4\}$ means real $v_2\{4\}$



Components and cumulants in p+Au and d+Au at 200 GeV

Phys. Rev. Lett. 120, 062302 (2018)



Components and cumulants in p+Au and d+Au at 200 GeV

Phys. Rev. Lett. 120, 062302 (2018)



$c_2\{4\}$ is positive in p+Au

Can we blame this on nonflow?

Components and cumulants in p+Au and d+Au at 200 GeV

Phys. Rev. Lett. 120, 062302 (2018)





Components and cumulants in p+Au and d+Au at 200 GeV

Use of subevents further suppresses nonflow

Positive $c_2\{4\}$ in p+Au doesn't seem to be related to nonflow



Cumulants in p+Au and d+Au at 200 GeV

B. Schenke et al, Phys. Lett. B 803, 135322



Cumulants are computationally expensive in hydro theory, so not as well-studied

This particular calculation doesn't show the strong geometry dependence seen in the data

Important to note this is 2+1D hydro, so the kinematics can't match the data

Particle species dependence of “Cronin enhancement”

PHENIX, Phys. Rev. C 88, 024906 (2013)



π^+, π^-, π^0,
K^+, K^-,
p, $\bar{p}$,
ϕ

Protons much more strongly modified than pions
ϕ mesons similar to pions

Photons in small systems



Photons in small systems





- Thermal(-ish) photons in p+Au?

Photons in small systems



- Thermal(-ish) photons in p+Au? Theory from Phys. Rev. C 95, 014906 (2017)

Photon yields



Common scaling for Au+Au and Pb+Pb at different energies; very different from N_{coll}-scaled $p+p$

Photon yields



Common scaling for Au+Au and Pb+Pb at different energies; very different from N_{coll}-scaled p+p

p+Au and d+Au in between

Comparisons with STAR

Phys. Rev. C 96, 064905 (2017)



FVTX EP: $0.65 < \Delta\eta < 3.35$

MPC EP: $2.75 < \Delta\eta < 4.05$

2PC: 3-sub event method with BBC, FVTX, CA

- Nonflow is kinematically suppressed in PHENIX

Comparisons with STAR

Phys. Rev. C 96, 064905 (2017)





- Nonflow is kinematically suppressed in PHENIX
- STAR measurement uses kinematic range with more nonflow
  —Subtracted result matches PHENIX
- For highest p_T points, oversubtraction is an issue
  —See S. Lim et al, Phys. Rev. C 100, 024908 (2019)

Comparisons with STAR





- Nonflow is kinematically suppressed in PHENIX
- We can also choose a different set of detectors to better (NOT exactly) match the STAR acceptance

  —Good agreement with STAR unsubtracted results in this configuration (more nonflow)